



Double Disaster: Mental Health Of Survivors of Wildfires and Earthquake in a Part of Greece

PDF (Size: 79KB) PP. 132-137 DOI: 10.4236/psych.2011.22021

Author(s)

Vicky Papanikolaou, Dimitrios Adamis, Robert C. Mellon, Gerasimos Prodromitis, John Kyriopoulos

ABSTRACT

This paper investigates stress related psychological morbidity in individuals who experienced two disasters 11 months apart (wildfire and earthquake) in a rural area of Greece. A sample of 150 participants has been assessed after the wildfires and after the earthquake using the Symptom Checklist 90-Revised. Survivors had elevated levels of psychopathology in all subscales of the SCL-90-R after the earthquake. Significant risk factors for further development of psychopathology were damages to property and complete loss of property from both disasters. Double disasters can cause considerable psychological symptoms in victims and there are reasons for policy makers to create services in order to help and improve the mental health of those affected but also to help them rebuild their property.

KEYWORDS

Double Disaster, Greece, Psychopathology, Psychological Distress, Adults

Cite this paper

Papanikolaou, V. , Adamis, D. , Mellon, R. , Prodromitis, G. & Kyriopoulos, J. (2011). Double Disaster: Mental Health Of Survivors of Wildfires and Earthquake in a Part of Greece. *Psychology*, 2, 132-137. doi: 10.4236/psych.2011.22021.

References

- [1] Clayer, J. R., Bookless-Pratz, C., & Harris, R. L. (1985). Some health consequences of a natural disaster. *Medical Journal of Australia*, 143, 182-184.
- [2] Den Ouden, D. J., Van Der Velden, P. G., Grevink, L., Morren, M., Dirkzwager, A. J. E., & Yzermans, C. J. (2007). Use of mental health services among disaster survivors: Predisposing factors. *BMC Public Health*, 7, 173. doi:10.1186/1471-2458-7-173
- [3] Derogatis, L. R. & Psychometric, R. C. (1992). SCL-90-R: Administration, scoring & procedures manual-II, for the revised version and other instruments of the psychopathology rating scale series. *Clinical Psychometric Research*, Towson.
- [4] Donias, S., Karastergiou, A., & Manos, N. (1991). Standardization of the symptom checklist 90 rating scale in a Greek population. *Psychiatriki*, 2, 42-48.
- [5] EM-DAT. (2008). The OFDA/CRED International Disaster Database PreventionWeb, Louvain Centre for Research on the Epidemiology of Disasters, Ecole de Santé Publique, Université Catholique de Louvain. <http://www.preventionweb.net/english/countries/statistics/index.php?cid=68>. Retrieved 15/9/2010..
- [6] Freedy, J. R., Saladin, M. E., Kilpatrick, D. G., Resnick, H. S., & Saunders, B. E. (1994) Understanding acute psychological distress following natural disaster. *Journal of Trauma Stress*, 7, 257-273. doi:10.1002/jts.2490070207
- [7] Gibbs, M. S. (1989). Factors in the victim that mediate between disaster and psychopathology: A review. *Journal of Traumatic Stress*, 2, 489- 514. doi:10.1002/jts.2490020411

• Open Special Issues

• Published Special Issues

• Special Issues Guideline

PSYCH Subscription

Most popular papers in PSYCH

About PSYCH News

Frequently Asked Questions

Recommend to Peers

Recommend to Library

Contact Us

Downloads: 258,591

Visits: 569,301

Sponsors, Associates, and Links >>

- [8] Heir, T. & Weisaeth, L. (2008). Acute disaster exposure and mental health complaints of Norwegian tsunami survivors six months post disaster. *Psychiatry*, 71, 266-276. doi:10.1521/psyc.2008.71.3.266
- [9] Hussain, A., Weisaeth, L., & Heir, T. (2010, In Press) Psychiatric disorders and functional impairment among disaster victims after exposure to a natural disaster: A population based study. *Journal of Affective Disorders*.
- [10] Mellon, R. C., Papanikolau, V., & Prodromitis, G. (2009). Locus of control and psychopathology in relation to levels of trauma and loss: Self-reports of Peloponnesian wildfire survivors. *Journal of Trauma Stress*, 22, 189-196. doi:10.1002/jts.20411
- [11] Norris, F. H., Friedman, M. J., Watson, P. J., Byrne, C. M., Diaz, E., & Kaniasty, K. (2002). 60,000 disaster victims speak: Part I. An empirical review of the empirical literature, 1981-2001. *Psychiatry*, 65, 207-239. doi:10.1521/psyc.65.3.207.20173
- [12] Norris, F. H., Friedman, M. J., & Watson, P. J. (2002). 60,000 disaster victims speak: Part II. Summary and implications of the disaster mental health research. *Psychiatry*, 65, 240-260. doi:10.1521/psyc.65.3.240.20169
- [13] O'Neill, H. K., Evans, B. A., Bussman, M. D., & Strandberg, D. K. (1999). Psychological distress during the Red River flood: Predictive utility of the Conservation of Resources model. *Applied Behavioral Science Review*, 7, 159-169. doi:10.1016/S1068-8595(00)80016-1
- [14] Smith, B. W. and Freedy, J. R. (2000) Psychosocial resource loss as a mediator of the effects of flood exposure on psychological distress and physical symptoms. *Journal of Trauma Stress*, 13, 349. doi:10.1023/A:1007745920466
- [15] Solomon, S., Smith, E. M., Robins, L., & Fischbach, R. L. (1987) Social involvement as a mediator of disaster-induced stress. *Journal of Applied Social Psychology*, 17, 1092-1112. doi:10.1111/j.1559-1816.1987.tb02349.x
- [16] Weiss, M. G., Saraceno, B., Saxena, S., & van Ommeren, M. (2003). Mental health in the aftermath of disasters: Consensus and controversy. *Journal of Nervous and Mental Disease*, 191, 611-615. doi:10.1097/01.nmd.0000087188.96516.a3