



Mathematics > Differential Geometry

A remark on weighted Bergman kernels on orbifolds

[Xianzhe Dai](#), [Kefeng Liu](#), [Xiaonan Ma](#)

(Submitted on 23 Jul 2011)

In this note, we explain that Ross-Thomas' result on the weighted Bergman kernels on orbifolds can be directly deduced from our previous result. This result plays an important role in the companion paper to prove an orbifold version of Donaldson Theorem.

Subjects: **Differential Geometry (math.DG)**; Complex Variables (math.CV)

Cite as: **arXiv:1107.4677 [math.DG]**
(or **arXiv:1107.4677v1 [math.DG]** for this version)

Submission history

From: Xiaonan Ma [[view email](#)]
[v1] Sat, 23 Jul 2011 10:47:30 GMT (7kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.DG

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math.CV](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

