



On removability properties of ψ -uniform domains in Banach spaces

M. Huang, M. Vuorinen, X. Wang

(Submitted on 1 Jun 2012)

Suppose that E and E' denote real Banach spaces with the same dimension at least 2. The main aim of this paper is to show that a domain D in E is a ψ -uniform domain if and only if $D \setminus P_D$ is a ψ_1 -uniform domain, and a domain D in E is a uniform domain if and only if $D \setminus P_D$ is also a uniform domain, where P_D denotes a countable set in D with the property that the quasihyperbolic distance between each pair of distinct points in it has a lower bound greater than or equal to $1/2$.

Subjects: **Complex Variables (math.CV)**

Cite as: **arXiv:1206.0128 [math.CV]**

(or **arXiv:1206.0128v1 [math.CV]** for this version)

Submission history

From: Manzi Huang [[view email](#)]

[v1] Fri, 1 Jun 2012 09:33:16 GMT (14kb)

Which authors of this paper are endorsers?

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.CV

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1206](#)

Change to browse by:

[math](#)

References & Citations

- [NASA ADS](#)

Bookmark ([what is this?](#))

