



Families of Integrable Equations

[Pavlos Kassotakis](#), [Maciej Nieszporski](#)

(Submitted on 3 Jun 2011 (v1), last revised 28 Oct 2011 (this version, v3))

We present a method to obtain families of lattice equations. Specifically we focus on two of such families, which include 3-parameters and their members are connected through Bäcklund transformations. At least one of the members of each family is integrable, hence the whole family inherits some integrability properties.

Subjects: **Exactly Solvable and Integrable Systems (nlin.SI);**
Mathematical Physics (math-ph)

MSC classes: 82B20, 37K35, 39A05

Journal reference: SIGMA 7 (2011), 100, 14 pages

DOI: [10.3842/SIGMA.2011.100](https://doi.org/10.3842/SIGMA.2011.100)

Cite as: [arXiv:1106.0636 \[nlin.SI\]](https://arxiv.org/abs/1106.0636)
(or [arXiv:1106.0636v3 \[nlin.SI\]](https://arxiv.org/abs/1106.0636v3) for this version)

Submission history

From: [Maciej Nieszporski \[view email\]](#)

[\[v1\]](#) Fri, 3 Jun 2011 13:36:05 GMT (12kb)

[\[v2\]](#) Fri, 30 Sep 2011 01:58:54 GMT (15kb)

[\[v3\]](#) Fri, 28 Oct 2011 05:51:31 GMT (28kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nlin.SI

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[math](#)

[math-ph](#)

[nlin](#)

References & Citations

- [NASA ADS](#)

Bookmark (what is this?)

