



High Energy Physics - Theory

Swaying oscillons in the signum-Gordon model

H. Arodz, Z. Swierczynski

(Submitted on 16 Jun 2011)

We present a new class of oscillons in the (1+1)-dimensional signum-Gordon model. The oscillons periodically move to and fro in the space. They have finite total energy, finite size, and are strictly periodic in time. The corresponding solutions of the scalar field equation are explicitly constructed from the second order polynomials in the time and position coordinates.

Comments: 12 pages, 3 figures

Subjects: **High Energy Physics - Theory (hep-th)**; Mathematical Physics (math-ph); Exactly Solvable and Integrable Systems (nlin.SI)

Cite as: **arXiv:1106.3169 [hep-th]**
(or **arXiv:1106.3169v1 [hep-th]** for this version)

Submission history

From: Henryk Arodz [[view email](#)]

[v1] Thu, 16 Jun 2011 08:40:15 GMT (86kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

hep-th

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[math](#)

[math-ph](#)

[nlin](#)

[nlin.SI](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

