

A New Preconditioner with Two Variable Relaxation Parameters for Saddle Point Linear Systems with Highly Singular (1,1) Blocks

Yuping Zeng, Chenliang Li

School of Mathematics and Computing Science, Guilin University of Electronic Technology, Guilin, China

E-mail: chenliang_li@hotmail.com

Received July 24, 2011; revised August 26, 2011; accepted September 6, 2011

Abstract

In this paper, we provide new preconditioner for saddle point linear systems with (1,1) blocks that have a high nullity. The preconditioner is block triangular diagonal with two variable relaxation parameters and it is extension of results in [1] and [2]. Theoretical analysis shows that all eigenvalues of preconditioned matrix is strongly clustered. Finally, numerical tests confirm our analysis.

Keywords: Saddle Point Linear Systems, Block Triangular Preconditioner, Krylov Subspace Methods

1. Introduction

Consider the following saddle point linear system

$$Ax \equiv \begin{pmatrix} F & B^T \\ B & 0 \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix} = \begin{pmatrix} f \\ g \end{pmatrix} \equiv b, \quad (1)$$

where $F \in R^{n \times n}$ is symmetric and positive semidefinite with nullity r , $B \in R^{m \times n}$ ($m \leq n$) has full row rank, $f \in R^n$ and $g \in R^m$, u and p are unknown. Note that the assumption that A is nonsingular, *i.e.*, the system (1) has a unique solution implies that $\text{null}(F)/\text{null}(B) = 0$, which we use in our analysis below. Saddle point linear systems of form (1) can arise, for example, from constraint optimization [3], mixed finite element formulation for the Stokes problem [4], and discrete time harmonic Maxwell equations in mixed form [5].

There has many techniques for solving Saddle point linear systems of form (1), see [6] for a comprehensive survey. However, when F is singular, it cannot be inverted and the Schur complement does not exist. In this case, one possible way of dealing with system is by augmentation [7]. Another way we can refer to [8] where Grief and Schötzau exploited a preconditioning technique for solving time-harmonic Maxwell equations in mixed form.

Recently, Rees and Grief [2] extend the work by Grief and Schötzau [8] to interior point methods for optimization problems. The preconditioner has attractive property of improved eigenvalue clustering with ill-conditioned

the (1,1) block of saddle point systems. Based on the basis of above work, Huang *et al.* constructed two block triangular preconditioners for solving saddle point systems (1) [1].

In this paper we are devoted to give new block triangular preconditioner for solving saddle point systems of (1) with an ill-conditioned (1,1) blocks. The preconditioner is involving two parameters, and they are extension of recent work in Grief and Schötzau [8], Rees and Grief [2], and Cheng *et al.* [1].

2. New Preconditioner and Spectral Analysis

Rees and Grief [9] provided the following preconditioner for the symmetric saddle point systems (1)

$$M = \begin{pmatrix} F + B^T W^{-1} B & tB^T \\ 0 & W \end{pmatrix},$$

where t is a scalar and $W \in R^{m \times m}$ is symmetric positive weight matrix.

Recently, Huang *et al.* [6] established the following preconditioners for the saddle point systems:

$$M_t = \begin{pmatrix} F + B^T W^{-1} B & (1-t)B^T \\ 0 & tW \end{pmatrix},$$

where $t \neq 0$ is a parameter and

$$\hat{M}_t = \begin{pmatrix} F + tB^T W^{-1} B & tB^T \\ 0 & \frac{(1-t)}{t}W \end{pmatrix},$$

where $1 \neq t > 0$.

In this section, we introduce the following preconditioner involving two parameters:

$$M_{\eta,\varepsilon} = \begin{pmatrix} F + \eta B^T W^{-1} B & (1 - \eta\varepsilon) B^T \\ 0 & \varepsilon W \end{pmatrix}, \quad (2)$$

for the saddle point systems (1), where $\eta > 0$ and $\varepsilon \neq 0$. We note that when the parameter $\eta = 1$, and $\varepsilon = t$, $M_{\eta,\varepsilon}$ reduce to M_t ; when $\eta = t$ and $\varepsilon = \frac{1-t}{t}$, $M_{\eta,\varepsilon}$ reduce to M_t .

Theorem 2.1. The matrix $M_{\eta,\varepsilon}^{-1} A$ has two distinct eigenvalues, given by

$$\lambda_1 = 1 \text{ and } \lambda_2 = -1/\eta\varepsilon,$$

with the algebraic multiplicities n and r , respectively. The remaining $m - r$ eigenvalues satisfy the relation

$$\lambda = -\frac{\mu}{\varepsilon(\eta\mu + 1)}, \quad (3)$$

where μ are positive generalized eigenvalues of

$$B^T W^{-1} B u = \mu F u, \quad (4)$$

Let $\{x_i\}_{i=1}^r$ be a basis of the null space of F , $\{z_i\}_{i=1}^{n-m}$ be a basis of the null space of B , and $\{y_i\}_{i=1}^{n-m}$ a set of linearly independent vectors that complete $\text{null}(F) \cup \text{null}(B)$ to a basis of R^n . Then the r vectors

$\left(x_i; \frac{1}{\varepsilon} W^{-1} B x_i\right)$, the $n - m$ vectors $(z_i; 0)$, and the $m - r$

vectors $\left(y_i, \frac{1}{\varepsilon} W^{-1} B y_i\right)$ are linearly independent eigenvec-

tors associated with $\lambda = 1$, and the r vectors $(x_i; -\eta W^{-1} B x_i)$ are eigenvectors associated with

$$\lambda = -\frac{1}{\eta\varepsilon}.$$

Proof: Suppose that λ is an eigenvalue of $\hat{M}_{\eta,\varepsilon}^{-1} A$, whose eigenvector is $(u^T, p^T)^T$. Then the corresponding eigenvalue problem is

$$\begin{pmatrix} F & B^T \\ B & 0 \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix} = \lambda \begin{pmatrix} F + \eta B^T W^{-1} B u & (1 - \eta\varepsilon) B^T \\ 0 & \varepsilon W \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix}.$$

From the second row we can obtain $p = \frac{1}{\varepsilon\lambda} W^{-1} B u$.

By substituting it into the first row we have

$$(1 - \lambda) \left[\lambda F u + \left(\frac{1}{\varepsilon\lambda} + \lambda \eta \right) B^T W^{-1} B u \right] = 0. \quad (5)$$

If $\lambda = 1$, then (5) is satisfied for any $u \in R^n$, and hence $(u; W^{-1} B u)$ is an eigenvector of $M_{\eta,\varepsilon}^{-1} A$.

If $u \in \text{null}(F)$, then from (5) we obtain

$$(1 - \lambda) \left(\frac{1}{\varepsilon\lambda} + \lambda \eta \right) B^T W^{-1} B u = 0.$$

From which it follows that $\left(u; \frac{1}{\varepsilon} W^{-1} B u\right)$ and $(u; -\eta W^{-1} B u)$ are eigenvectors associated $\lambda = 1$ and $\lambda = -\frac{1}{\eta\varepsilon}$.

Next, suppose $\lambda = 1$ and $\lambda = -1/\eta\varepsilon$. We divide (5) by $(1 - \lambda)(1/\varepsilon + \lambda\eta)$, which yields (3), with u defined in (4).

Now we can find a specific set of linearly independent eigenvectors for $\lambda = 1$ and $\lambda = -1/\eta\varepsilon$. Since $\text{null}(F) \cap \text{null}(B) = 0$, the vectors $\{x_i\}_{i=1}^r$ and $\{z_i\}_{i=1}^{n-m}$ defined above are linearly independent and form a subspace of R^n of dimension $n - m + r$. Let $\{y_i\}_{i=1}^{n-m}$ complete this set to a basis of R^n . It follows that $(x_i, \varepsilon^{-1} W^{-1} B x_i)$, $(z_i, 0)$, and $(y_i, \varepsilon^{-1} W^{-1} B y_i)$ are eigenvectors associated with $\lambda = 1$. The r vectors $(x_i, -\eta W^{-1} B x_i)$ are eigenvectors associated with $\lambda = -1/\eta\varepsilon$.

When the parameters satisfy $-1/\eta\varepsilon = 1$, we can obtain the following corollary from Theorem 2.1.

Corollary 2.2. Let $-1/\eta\varepsilon = 1$. Then the matrix $M_{\eta,\varepsilon}^{-1} A$ has one eigenvalue which given by $\lambda = 1$ with algebraic multiplicity $n + r$. The remaining $m - r$ eigenvalues satisfy the relation

$$\lambda = -\eta\mu / (\eta\mu + 1),$$

where μ are positive generalized eigenvalues of

$$B^T W^{-1} B u = \mu F u.$$

Theorem 2.3. The matrix $M_{\eta,\varepsilon}^{-1} A$ has two distinct eigenvalues, given by

$$\lambda_1 = 1 \text{ and } \lambda_2 = -1/\eta\varepsilon,$$

with the algebraic multiplicities n and r , respectively. The remaining $m - r$ eigenvalues lie in the interval

$$(0, -1/\eta\varepsilon)(\varepsilon < 0), \text{ or } (-1/\eta\varepsilon, 0)(\varepsilon > 0).$$

Proof: From Theorem 2.1 we obtain that the matrix $M_{\eta,\varepsilon}^{-1} A$ has two distinct eigenvalues, given by

$$\lambda_1 = 1 \text{ and } \lambda_2 = -1/\eta\varepsilon,$$

with the algebraic multiplicities n and r , respectively.

Taking inner product of (5) with u , we can obtain that the remaining $m - r$ eigenvalues satisfy

$$\lambda = \frac{-\langle B^T W^{-1} B u, u \rangle / (\eta\varepsilon)}{\langle F u, u \rangle / \eta + \langle B^T W^{-1} B u, u \rangle}, \quad (6)$$

where $\langle \cdot, \cdot \rangle$ denotes the standard Euclidean inner product, $u \notin \text{null}(F)$ and $u \notin \text{null}(B)$. By (6), we have that the remaining $m - r$ eigenvalues lie in interval

$$(0, -1/\eta\varepsilon)(\varepsilon < 0), \text{ or } (-1/\eta\varepsilon, 0)(\varepsilon > 0).$$

When the parameters satisfy $-1/\eta\varepsilon = 1$, we can obtain the following corollary from Theorem 2.3.

Corollary 2.4. Let $-1/\eta\varepsilon = 1$. Then the matrix $M_{\eta\varepsilon}^{-1}A$ has one eigenvalue which given by $\lambda = 1$ with algebraic multiplicity $n + r$. The remaining $m - r$ eigenvalues lie in the interval $(0, 1)$.

3. Numerical Experiments

We consider the following finite element discretization of the time-harmonic Maxwell equations ($k^2 = 0$) [5,8]. The following two-dimensional Model problem is considered: find u and p that satisfy

$$\begin{aligned} \nabla \times \nabla \times u + \nabla p &= f \text{ in } \Omega \\ \nabla \cdot u &= 0 \text{ in } \Omega \\ u \times n &= 0 \text{ on } \partial\Omega \\ p &= 0 \text{ on } \partial\Omega \end{aligned} \tag{7}$$

Here $\Omega \in R^2$ is a simply connected polyhedron domain with a connected boundary $\partial\Omega$, and $\sim n$ denotes the outward unit normal on $\partial\Omega$. The datum f is a given source (not necessarily divergence free). Using the lowest order N'ed'elec elements of first kind [9,10] for the approximation of the vector field and standard nodal elements for multiplier yields the following saddle-point linear system

$$Ax \equiv \begin{pmatrix} F & B^T \\ B & 0 \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix} = \begin{pmatrix} g \\ 0 \end{pmatrix} \equiv b.$$

Experiments were done in a square domain ($0 \leq x \leq 1$; $0 \leq y \leq 1$). And we set the right-hand side function so that the exact solution is given by

$$u(x, y) = (y(1 - y), x(1 - x))^T.$$

In our numerical experiments the matrix W in the augmentation block preconditioner is taken as $W = I$.

We consider three meshes with different values of n and m in **Table 1**. **Table 2** shows iteration counts for different η and meshes, applying BiCGSTAB of block-triangular preconditioner, and $-1/\eta\varepsilon = 1$. We ob-

Table 1. Values of n and m and size of the linear systems for three meshes.

h	n	m	$n + m$
$\frac{1}{8}$	176	49	225
$\frac{1}{16}$	225	736	961
$\frac{1}{32}$	961	3008	3969

Table 2. Iteration counts for different and meshes, using BiCGSTAB for solving the saddle point system with preconditioner $\mathcal{M}_{\eta\varepsilon}$, the iteration was stopped once

$$\|r^{(k)}\| / \|r^{(0)}\| \leq 10^{-13}.$$

h	0.1	0.5	2	4	6
$h = \frac{1}{8}$	3	3	1	1	0.5
$h = \frac{1}{16}$	3	2	2	1	1
$h = \frac{1}{32}$	1	0.5	0.5	0.5	0.5

serve that for a fixed mesh, When $\eta = 6$, We spent the least iteration counts. In particular, when $\eta > 2$, the iteration we spent are less than $\eta = 2$ (corresponding to M_2 [1]). It shows that preconditioner $\lambda = M_{\eta\varepsilon}$ is more efficient than M_t and \hat{M}_t .

4. Acknowledgements

The authors thanks the Chinese National Science Foundation Project (11161014), the Science and Technology Development Foundation of Guangxi (Grant No. 0731 018), Innovation Project of Guangxi Graduate Education (Grant No. ZYC0430).

5. References

- [1] T. Z. Huang, G. H. Cheng and S. Q. Shen, "New Block Triangular Preconditioners for Saddle Point Linear Systems with Highly Sigular (1,1) Blocks," *Compututer Physics Communications*, Vol. 180, No. 2, 2009, pp. 192-196.
- [2] T. Rees and C. Grief, "A Preconditioner for Linear Systems Arising from Interios Point Optimization Methods," *SIAM Journal of Scientific Computing*, Vol. 29, No. 5, 2007, pp. 1992-2007. [doi:10.1137/060661673](https://doi.org/10.1137/060661673)
- [3] S. Wright, "Stability of Augmented System Factorizations in Interior-Point Methods," *SIAM Journal of Matrix Analysis and Applications*, Vol. 18, No. 1, 1997, pp. 191-222. [doi:10.1137/S0895479894271093](https://doi.org/10.1137/S0895479894271093)
- [4] V. Girault and P. Raviart, "Finite Elment methods for Navier-Stokes Equations," Springer-Verlag, Berlin, 1986. [doi:10.1007/978-3-642-61623-5](https://doi.org/10.1007/978-3-642-61623-5)
- [5] C. Grief and D. Schöttau, "Preconditioners for Discretized Time-Harmonic Maxwell Equations in Mixed Form," *Numerical Linear Algebra with Applications*, Vol. 14, No. 4, 2007, pp. 281-297. [doi:10.1002/nla.515](https://doi.org/10.1002/nla.515)
- [6] M. Benzi, G. H. Golub and J. Lieson, "Numerical Solution of Saddle Point Problems," *Acta Numerica*, Vol. 14, 2005, pp.1-137. [doi:10.1017/S0962492904000212](https://doi.org/10.1017/S0962492904000212)
- [7] G. H. Golub and C. Grief, "On Solving Block Structured Indefinite Linear Systems," *SIAM Journal of Scientific Computing*, Vol. 24, No. 6, 2003, pp. 2076-2092. [doi:10.1137/S1064827500375096](https://doi.org/10.1137/S1064827500375096)

- [8] C. Grief and D. Schötzau, "Preconditioners for Saddle Point Linear Systems with Highly Singular (1,1) Blocks," *Electronic Transactions on Numerical Analysis*, Vol. 22, 2006, pp. 114-121.
- [9] P. Monk, "Finite Elements for Maxwell's Quations," Oxford University Press, New York, 2003.
- [10] J. C. Nédélec, "A New Family of Mixed Finite Elements in \mathbb{R}^3 ," *Numerische Mathematik*, Vol. 50, No. 1, 1986, pp. 57-81. [doi:10.1007/BF01389668](https://doi.org/10.1007/BF01389668)