

## 包含纯净两因子交互作用的分区组折衷设计的某些结果

卢海威<sup>1</sup>, 赵胜利<sup>2</sup>, 卢星辰<sup>3</sup>

1. 泰山医学院基础医学院, 泰安 271016;
2. 曲阜师范大学数学科学学院, 曲阜 273165;
3. 泰山医学院外国语学院, 泰安 271016

## Some Results on Blocked Compromise Plans with Clear Two-factor Interactions

LU Haiwei<sup>1</sup>, ZHAO Shengli<sup>2</sup>, LU Xingchen<sup>3</sup>

1. School of Basic Medical Sciences, Taishan Medical University, Taian 271016;
2. School of Mathematical Sciences, Qufu Normal University, Qufu 273165;
3. Foreign Languages School, Taishan Medical University, Taian 271016

- [摘要](#)
- [参考文献](#)
- [相关文章](#)

全文: [PDF \(287 KB\)](#) [HTML \(1 KB\)](#) 输出: [BibTeX](#) | [EndNote \(RIS\)](#) [背景资料](#)

## 服务

- ▶ [把本文推荐给朋友](#)
- ▶ [加入我的书架](#)
- ▶ [加入引用管理器](#)
- ▶ [E-mail Alert](#)
- ▶ [RSS](#)

## 作者相关文章

- ▶ [卢海威](#)
- ▶ [赵胜利](#)
- ▶ [卢星辰](#)

**摘要** 纯净的分区组折衷设计可以估计所有的因子主效应和一部分特殊的两因子交互作用, 而且不需要假设其他的两因子交互作用是可以忽略的. 本文主要研究了几类纯净的分区组折衷设计存在的必要条件, 并给出了一些纯净的分区组设计的构造方法.

**关键词:** 折衷设计 纯净 分辨率

**Abstract:** The clear blocked compromise plans allow estimation of all main effects and some specified two-factor interactions without assuming the remaining two-factor interactions are negligible. This paper gets some necessary conditions of the existence of some kinds of clear blocked compromise plans and provides some methods for constructing clear blocked compromise plans.

**Key words:** compromise plan clear resolution





收稿日期: 2010-03-20;






基金资助: 国家自然科学基金(10826059, 10901092), 山东省自然科学基金(Q2007A05), 山东省博士基金(BS2011SF006), 山东省软科学(2010RKGA1048) 资助项目.

引用本文:

卢海威, 赵胜利, 卢星辰. 包含纯净两因子交互作用的分区组折衷设计的某些结果[J]. 应用数学学报, 2012, 35(5): 892-900.

LU Haiwei, ZHAO Shengli, LU Xingchen. Some Results on Blocked Compromise Plans with Clear Two-factor Interactions[J]. Acta Mathematicae Applicatae Sinica, 2012, 35(5): 892-900.

- [1] Zhang R C, Park D K. Optimal Blocking of Two-level Fractional Factorial Designs. *Journal of Statistical Planning and Inference*, 2002, 91: 107-121
- [2] Wu C F J, Chen Y. A Graph-aided Method for Planning Two-level Experiments When Certain Interactions are Important. *Technometrics*, 1992, 34: 162-175 
- [3] Chen B J, Li P F, Liu M Q, Zhang R C. Some Results on Blocked Regular 2-level Fractional Factorial Designs with Clear Effects. *Journal of Statistical Planning and Inference*, 2006, 136(12): 4436-4449 
- [4] Ai M Y, He S Y. An Efficient Method for Identifying Clear Effects in Blocked Fractional Factorial Designs. *Statistics and Probability Letters*, 2006, 76: 1889-1894 
- [5] Tang B. Orthogonal Arrays Robust to Nonnegligible Two-factor Interactions. *Biometrika*, 2006, 93: 137-146 

- [6] Xu H Q. A Catalogue of Three-level Regular Fractional Factorial Designs. *Metrika*, 2005, 62: 259-281 
- [7] Ai M Y, Zhang R C.  $s^{n-m}$  Designs Containing Clear Main Effects or Clear Two-factor Interactions. *Statistics and Probability Letters*, 2004, 69: 151-160 
- [8] Chen H, Hedayat A S.  $2^{n-m}$  Designs With Resolution III and IV Containing Clear Two-factor Interactions. *Journal of Statistical Planning and Inference*, 1998, 75: 147-158 
- [9] Ke W, Tang B, Wu H. Compromise Plans with Clear Two-factor Interactions. *Statistica Sinica*, 2005, 15: 709-715
- [10] Li P F, Chen B J, Liu M Q, Zhang R C. A Note on Minimum Aberration and Clear Criteria. *Statistics and Probability Letters*, 2006, 76: 1007-1011 
- [11] Tang B, Ma F, Ingram D, Wang H. Bounds on the Maximum Number of Clear Two-factor Interactions for  $2^{m-p}$  Designs of Resolution III and IV. *Canadian Journal of Statistics*, 2002, 30: 127-136 
- [12] Wu H Q, Wu C F J. Clear Two-factor Interaction and Minimum Aberration. *Annals of Statistics*, 2002, 30: 1496-1511 
- [13] Yang J F, Li P F, Liu M Q, Zhang R C.  $2^{(n_1+n_2)-(k_1+k_2)}$  Fractional Factorial Split-plot Designs Containing Clear Effects. *Journal of Statistical Planning and Inference*, 2006, 136(12): 4450-4458 
- [14] Yang G J, Liu M Q. A Note on the Lower Bounds on Maximum Number of Clear Two-factor Interactions for  $2_{III}^{m-p}$  and  $2_{IV}^{m-p}$  Designs. *Communications in Statistics—theory and Methods*, 2006, 35: 849-860 
- [15] Yang G J, Liu M Q, Zhang R C. Weak Minimum Aberration and Maximum Number of Clear Two-factor Interactions in  $2_{IV}^{m-p}$  Designs. *Science in China (Series A)*, 2005, 48(11): 1479-1487
- [16] Zhao S L, Zhang R C.  $2^{m4^n}$  Designs with Resolution III or IV Containing Clear Two-factor Interaction Components. *Statistical Papers*, 2008, 49: 441-454
- [17] Zhao S L, Zhang R C. Compromise  $4^m 2^n$  Plans with Clear Two-factor Interactions. *Acta Mathematicae Applicatae Sinica*, 2010, 26(1): 99-106 
- [18] Zhao S L, Zhang R C, Liu M Q. Some Results On  $4^{m2^n}$  Designs with Clear Two-factor Interaction Components. *Science in China (Series A)*, 2008, 51(7): 1297-1314
- [19] Zi X M, Liu M Q, Zhang R C. Asymmetrical Designs Containing Clear Effects. *Metrika*, 2007, 65: 123-131
- [20] Zi X M, Zhang R C, Liu M Q. Bounds On the Maximum Numbers of Clear Two-factor Interactions for  $2^{(n_1+n_2)-(k_1+k_2)}$  Fractional Factorial Split-plot Designs. *Science in China (Series A)*, 2006, 49: 1816-1829 
- [1] 杨贵军, 刘民千. 一类包含最大数量纯净两因子交互效应的  $2_{IV}^{m-p}$  设计[J]. 应用数学学报, 2005, 28(1): 143-150.