



华东师范大学学报(自然科学版) » 2013, Vol. 2013 » Issue (1): 76-90 DOI:

应用数学与基础数学

最新目录 | 下期目录 | 过刊浏览 | 高级检索

◀◀ Previous Articles | Next Articles ▶▶

## $E_6$ 型Weyl群中左胞腔的左连通性

米倩倩, 时俭益

华东师范大学--数学系, 上海 200241

Left-connectedness of left cells in the Weyl Group of type  $E_6$

MI Qian-qian, SHI Jian-yi

Department of Mathematics, East China Normal University, Shanghai 200241, China

- 摘要
- 参考文献
- 相关文章

全文: [PDF](#) (265 KB) [HTML](#) (1 KB) 输出: [BibTeX](#) | [EndNote \(RIS\)](#) [背景资料](#)

摘要 首先通过计算机编程找出 $E_6$ 型Weyl群左胞腔的所有极短元,利用这些极短元证明了 $E_6$ 型Weyl群的所有左胞腔都是左连通的,从而证明了Lusztig关于左胞腔左连通性的一个猜想在 $E_6$ 型Weyl群中是成立的.

关键词: [Weyl群](#) [左胞腔](#) [双边胞腔](#) [左连通性](#)

Abstract: We showed that all the left cells in the Weyl group  $E_6$  were left-connected, verifying a conjecture of Lusztig in our case.

Key words: [Weyl group](#) [left cells](#) [two-sided cells](#) [left-connectedness](#)

收稿日期: 2012-02-01; 出版日期: 2013-01-25

引用本文:

.  $E_6$ 型Weyl群中左胞腔的左连通性[J]. 华东师范大学学报(自然科学版), 2013, 2013(1): 76-90.

. Left-connectedness of left cells in the Weyl Group of type  $E_6$ [J]. Journal of East China Normal University(Natural Sc, 2013, 2013(1): 76-90.

### 服务

- ▶ 把本文推荐给朋友
- ▶ 加入我的书架
- ▶ 加入引用管理器
- ▶ E-mail Alert
- ▶ RSS

### 作者相关文章

- [1] {1}
- [2] KAZHDAN D, LUSZTIG G. Representation of Coxeter groups and Hecke algebras[J]. Invent Math 1979, 53: 165-184.
- [3] {2}
- [4] ASAI T, KAWANAKAN, SPALTENSTEIN N, et al. Open Problems in algebraic Groups [C]// Problems from the Conference on Algebraic Groups and Representations Held at Katata. Katata: Taniguchi Foundation, 1983.
- [5] {3}
- [6] SHI J Y. The Kazhdan-Lusztig cells in certain affine Weyl groups[M]. Lecture Notes in Math 1179. Berlin: Springer, 1986.
- [7] {4}
- [8] SHI J Y. A two-sided cell in an affine Weyl group, II[J]. J London Math Soc, 1988, 37(2): 253-264.

- [14] {5}
- [15] SHI J Y. Left cells containing a fully commutative element[J]. J Comb Theory Series A, 2005, 113: 556-565. 
- [17] {6}
- [18] SHI J Y. Left-connectedness of some left cells in certain Coxeter
- [19] groups of simply-laced type[J]. J Algebra, 2008, 319(6): 2410-2413.
- [20] {7}
- [21] TONG C Q. Left cells in Weyl group of type \$E\_6\$[J]. Comm in
- [22] Algebra, 1995, 23(13): 5031-5047.
- [23] {8}
- [24] SHI J Y. A new algorithm for finding an I.c.r set in certain
- [25] two-sided cells[J]. Pacific J Math, 2012, 256(1): 235--252.
- [26] {9}
- [27] LUSZTIG G. Cells in affine Weyl group [C]// Algebraic Groups and
- [28] Related Topics. Advanced Studies in Pure Math, 1985, 6: 255-287.
- [29] {10}
- [30] BARBASCH D, VOGAN D. Primitive ideals and orbital integrals in
- [31] complex classical groups[J]. Math Ann, 259: 153-199.
- [32] {11}
- [33] BARBASCH D, VOGAN D. Primitive ideals and orbital integrals in
- [34] complex exceptional groups[J]. J Algebra, 1983, 80: 350-382. 
- [35] {12}
- [36] LUSZTIG G. Characters of Reductive Groups over a Finite Field [M].
- [37] Ann Math Studies 107, Princeton: Princeton University Press, 1984.
- [38] {13}
- [39] LUSZTIG G. Cells in affine Weyl group, IV [J]. J Fac Fci Univ Tokyo
- [40] Sect IA Math, 1989, 36: 297-328. 
- [41] {14}
- [42] SHI J Y. Left cells in affine Weyl groups[J] T\$\ddot{o}\$hoku J.
- [43] Math, 1994, 46: 105-124.
- [44] {15}
- [45] CARTER R W. Finite Groups of Lie Type: Conjugacy Classes and Complex
- [46] Characters[M]. Wiley Series in Pure and Applied Mathematics. London:
- [47] John Wiley, 1985. 
- [1] 岳明仕. 拟分裂情形下仿射\,Weyl 群\,\$\widetilde{\{ \bm{C} \}}\_{-\{ \bm{4} \}}\$~的胞腔[J]. 华东师范大学学报(自然科学版), 2013, 2013(1): 61-75.
- [2] 黄 谦. 加权的~Coxeter~群~\$\widetilde{\{ \bm{C} \}}\_{-\{ \bm{n} \}}\$~的左胞腔[J]. 华东师范大学学报(自然科学版), 2013, 2013(1): 91-103, 114.

版权所有 © 2011 《华东师范大学学报(自然科学版)》编辑部  
本系统由北京玛格泰克科技发展有限公司设计开发 技术支持: support@magtech.com.cn