

General Relativity and Quantum Cosmology

Poincare algebra realized in Hamiltonian formalism of the Relativistic Theory of Gravitation

V.O.Soloviev, M.V.Tchichikina

(Submitted on 13 Jan 2010)

We obtain the Poincare group generators by proper choice of arbitrary functions present in the Relativistic Theory of Gravitation (RTG) Hamiltonian. Their Dirac brackets give the Poincare algebra in accordance with the fact that RTG has 10 integrals of motion.

Comments: 8 pages, Russian

Subjects: **General Relativity and Quantum Cosmology (gr-qc)**Cite as: [arXiv:1001.2099v1](#) [gr-qc]

Submission history

From: Vladimir O. Soloviev [[view email](#)]

[v1] Wed, 13 Jan 2010 08:39:08 GMT (6kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PostScript](#)
- [PDF](#)
- [Other formats](#)

Current browse context:

gr-qc

[< prev](#) | [next >](#)[new](#) | [recent](#) | [1001](#)

References & Citations

- [SLAC-SPIRES HEP](#)
([refers to](#) | [cited by](#))
- [CiteBase](#)

Bookmark([what is this?](#))

 [CiteULike logo](#) [Connotea logo](#) [BibSonomy logo](#) [Mendeley logo](#) [Facebook logo](#) [del.icio.us logo](#) [Digg logo](#) [Reddit logo](#)