



### 发光学应用及交叉前沿

#### 基于拉曼光谱的微重力三维培养条件下的人HepG2细胞的研究

王雁军<sup>1</sup>, 王桂文<sup>2</sup>, 姚辉璐<sup>2</sup>

1. 中国科学院动物研究所膜工程与生物膜国家重点实验室 细胞生物学系, 北京 100101;

2. 广西科学院 生物物理实验室, 广西 南宁 530003

PDF 下载

引用本文

摘要：采用一个波长为785 nm的半导体激光束来囚禁人的肝癌细胞并激发人的肝癌细胞的拉曼光谱,分析模拟微重力三维培养条件下的人肝癌细胞的DNA、蛋白和脂类的生物学物质的变化和表达情况。结果显示,微重力三维培养条件下和平面培养下的人肝癌细胞的拉曼光谱在527.1 367.1 438.1 659 cm<sup>-1</sup>处有明显的差异。

关键词：人肝癌细胞 微重力 三维培养 拉曼光谱

#### Study on Liver Cancer Cells (HepG2) Under Simulated Microgravity Based on Raman Spectroscopy

WANG Yan-jun<sup>1</sup>, WANG Gui-wen<sup>2</sup>, YAO Hui-lu<sup>2</sup>

1. State Key Laboratory of Biomembrane and Membrane Biotechnology, Institute of Zoology, Chinese Academy of Sciences (CAS), Beijing 100101, China;

2. Laboratory for Biophysics, Guangxi Academy of Sciences, Nanning 530003, China

Abstract: In our work, HepG2 was cultured onto biodegradable polyglycolic acid(PGA) polymer scaffolds, which was cultured in a rotating cell culture system (RCCS) to form a three-dimensional (3D) and human liver cancer cells cultured in common plate. We detected the spectra of HepG2 cells under two culture conditions and compared the spectra changes of the different cells under two culture conditions. The results show that Raman spectroscopy can differentiate among HepG2 cells under two culture conditions. 527 cm<sup>-1</sup> and 1 357 cm<sup>-1</sup> peaks of HepG2 cells under simulated microgravity are higher than that of HepG2 cells in common plate. 1 438 cm<sup>-1</sup> and 1 659 cm<sup>-1</sup> peaks of HepG2 cells under simulated microgravity are lower than that of HepG2 cells in common plate. It is known that 527 cm<sup>-1</sup> represents Fe(II), N-N-Melm, 1 357 cm<sup>-1</sup> represents Guanine, porphyrins (haemoglobin), lipids, 1 438 cm<sup>-1</sup> represents CH<sub>2</sub> deformation (lipids), cholesterol, and 1 659 cm<sup>-1</sup> represents Amide I: proteins. These biomaterials which induce these peak changes are possibly related to 3D culture.

Keywords: liver cancer cells simulated microgravity three dimensional culture Raman spectroscopy

收稿日期 2013-07-01 修回日期 2013-08-22 网络版发布日期

基金项目:

国家自然科学基金(60868002) 资助项目

通讯作者: 姚辉璐, E-mail: huiluy@gmail.com

作者简介: 王雁军(1976-), 女, 辽宁丹东人, 主要从事肝脏病和激光拉曼光谱的研究。E-mail: wangyanjun117@163.com

作者Email: huiluy@gmail.com

#### 参考文献:

- [1] Nakamura K, Kuga H, Morisaki T, *et al.* Simulated microgravity culture system for a 3-D carcinoma tissue model [J]. *Biotechniques*, 2002, 33(5):1068-1070.
- [2] Wang X, Wei G, Yu W, *et al.* Scalable producing embryoid bodies by rotary cell culture system and constructing engineered cardiac tissue with ES-Derived cardiomyocytes in vitro [J]. *Biotechnol. Prog.*, 2006, 22(3):811-818.
- [3] Martin A, Zhou A, Gordon R E, *et al.* Thyroid organoid formation in simulated microgravity: Influence of keratinocyte growth factor [J]. *Thyroid*, 2000, 10(6):481-487.
- [4] Draux F, Jeannesson P, Beljebbar A, *et al.* Raman spectral imaging of single living cancer cells: A preliminary study [J]. *Analyst*, 2009, 134(3):542-548.
- [5] Kang L L, Huang Y X, Luo M. Confocal Raman microscopy on single living young and old erythrocytes [J]. *Spectrosc. Spect. Anal.* (光谱学与光谱分析), 2008, 28(10):2343-2347 (in Chinese).
- [6] Schrader B, Schulz H, Andreev G N, *et al.* On-destructive NIR-FT-Raman spectroscopy of plant and animal tissues, of food and works of art [J]. *Talanta*, 2000, 53(1):35-45.
- [7] Xie C A, Li Y Q. Confocal micro-Raman spectroscopy of single biological cells using optical trapping and shifted excitation difference techniques [J]. *Appl. Phys.*, 2003, 93(5):2982-2986.
- [8] Xie C A, Li Y Q, Tanq W J. Study of dynamical process of heat denaturation in optically trapped single

#### 本刊中的类似文章

1. MgZnO半导体材料光致发光以及共振拉曼光谱研究[J]. 2013,34(9): 1149-1154
2. 高质量ZnO及BeZnO薄膜的发光性质[J]. 2013,34(8): 1035-1039
3. 肿节风的表面增强拉曼光谱检测[J]. 2013,34(8): 1084-1088
4. 不同发酵时间普洱茶的表面增强拉曼光谱鉴别分析[J]. 2013,34(2): 230-234
5. 溶液浓度对CCl<sub>4</sub> 费米共振的影响[J]. 2012,33(8): 812-816
6. 富硅氮化硅薄膜的制备及其光学带隙研究[J]. 2012,(6): 596-600
7. AlN插入层对α-AlGaIn的外延生长的影响[J]. 2012,(5): 519-524
8. (Cu,Al)掺杂ZnO薄膜表面缺陷的拉曼光谱研究[J]. 2012,33(1): 109-113
9. 心肌梗死患者红细胞的光镊拉曼光谱[J]. 2011,32(10): 1088-1092
10. 并五噻吩分子光谱和激发态的密度泛函理论[J]. 2008,29(5): 821-826
11. 沉积工艺参数对碳纳米管薄膜场发射性能的影响[J]. 2006,27(1): 129-133
12. 聚合物前驱体衍生SiC纳米棒的光学性质[J]. 2005,26(5): 641-646
13. 刻线镍膜上沉积的碳纳米管场发射特性[J]. 2004,25(6): 743-747

microorganisms by near-infrared Raman spectroscopy [J]. *Appl. Phys.*, 2003, 94(9):6138-6142.

- [9] Potten C S, Loeffler M. Stem cells: Attributes, cycles, spirals, pitfalls and uncertainties. Lessons for and from the crypt [J]. *Development*, 1990, 110(4):1001-1020.
- [10] Fischbach C, Kong H J, Hsiong S X, *et al.* Cancer cell angiogenic capability is regulated by 3D culture and integrin engagement [J]. *Proc. Natl. Acad. Sci. USA*, 2009, 106(2):399-404.
- [11] Kinasiewicz A, Kawiak J, Werynski A. 3D matrigel culture improves differentiated functions of HepG2 cells in vitro [J]. *Biocybern. Biomed. Eng.*, 2006, 26(4):47-55.
- [12] Fisher K E, Pop A, Koh W, *et al.* Tumor cell invasion of collagen matrices requires coordinate lipid agonist-induced G-protein and membrane-type matrix metalloproteinase-1-dependent [J]. *Mol. Cancer*, 2006, 5:69-72.