



High Energy Physics - Phenomenology

Nuclear parton distributions and deviations from DGLAP at an Electron Ion Collider

[Alberto Accardi](#), [Vadim Guzey](#), [Juan Rojo](#)

(Submitted on 20 Jun 2011)

We explore the potential of an Electron Ion Collider to determine nuclear modifications of PDFs. We find that gluon shadowing can be accurately measured down to $x=0.001$, and discuss the possibility of detecting non-linear QCD effects with inclusive measurements.

Comments: 4 pages, 6 figures, to appear in the proceedings of XIX International Workshop on Deep Inelastic Scattering and Related Subjects (DIS 2011), Newport News, VA USA, April 11-15, 2011

Subjects: **High Energy Physics - Phenomenology (hep-ph)**; High Energy Physics - Experiment (hep-ex); Nuclear Theory (nucl-th)

Cite as: [arXiv:1106.3839 \[hep-ph\]](#)
(or [arXiv:1106.3839v1 \[hep-ph\]](#) for this version)

Submission history

From: [Juan Rojo \[view email\]](#)

[v1] Mon, 20 Jun 2011 08:31:40 GMT (135kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

hep-ph

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[hep-ex](#)

[nucl-th](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

