



Nuclear Experiment

Anisotropic flow of charged particles at $\sqrt{s_{NN}} = 2.76$ TeV measured with the ALICE detector

[Ante Bilandzic](#)

(Submitted on 30 Jun 2011)

Measurements of anisotropic flow in heavy-ion collisions provide evidence for the creation of strongly interacting matter which appears to behave as an almost ideal fluid. Anisotropic flow signals the presence of multiple interactions and is very sensitive to the initial spatial anisotropy of the overlap region in non-central heavy-ion collisions. In this article we report measurements of elliptic v_2 , triangular v_3 , quadrangular v_4 and pentagonal v_5 flow. These measurements have been performed with 2- and multi-particle correlation techniques.

Comments: 4 pages, 4 figures, Quark Matter 2011 proceedings

Subjects: **Nuclear Experiment (nucl-ex)**

Cite as: [arXiv:1106.6209v1](#) [nucl-ex]

Submission history

From: Ante Bilandzic [[view email](#)]

[v1] Thu, 30 Jun 2011 12:49:18 GMT (37kb,D)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [Other formats](#)

Current browse context:

nucl-ex

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

