

# 正相液相色谱-串联质谱法分离普萘洛尔对映体

张娟红<sup>1,2</sup>, 王荣<sup>1,2\*</sup>, 谢华<sup>1</sup>, 孟宪栋<sup>1</sup>, 贾正平<sup>1,2\*</sup>, 马骏<sup>1</sup>, 张军莉<sup>1</sup>, 王娟<sup>1</sup>

1. 兰州军区兰州总医院全军临床药理基地, 甘肃 兰州 730050; 2. 兰州大学药学院, 甘肃 兰州 730000

## Enantiomeric separation of propranolol by normal phase chiral liquid chromatography coupled with tandem mass spectrometry

ZHANG Juanhong<sup>1,2</sup>, WANG Rong<sup>1,2\*</sup>, XIE Hua<sup>1</sup>, MENG Xiandong<sup>1</sup>, JIA Zhengping<sup>1,2\*</sup>, MA Jun<sup>1</sup>, ZHANG Junli<sup>1</sup>, WANG Juan<sup>1</sup>

1. Base for Clinical Pharmacology, Lanzhou General Hospital of Lanzhou Military Region, Lanzhou 730050, China; 2. Pharmacy College, Lanzhou University, Lanzhou 730000, China

摘要

参考文献

相关文章

Download: PDF (174KB) [HTML 1KB](#) Export: BibTeX or EndNote (RIS) Supporting Info

**摘要** 建立正相液相色谱-串联质谱(LC-MS/MS)分离普萘洛尔对映体的方法,并用于盐酸普萘洛尔片对映体含量测定。样品使用甲醇进行简单提取,采用Chiralcel OD-H手性柱,以正己烷-乙醇-氨水(70:30:0.4, v/v/v)为流动相,流速为0.4 mL/min。在正离子模式下,通过电喷雾离子化(ESI+),采用多反应监测(MRM)方式进行检测,用于定量分析检测的离子对为m/z 260.2→116.0,在20 min内完成普萘洛尔对映体定量分析。盐酸普萘洛尔对映体在2.5~1000 μg/L质量浓度范围内线性关系良好,定量限为2.5 μg/L;日内及日间测定的相对标准偏差小于2.64%。两种对映体的加样回收率范围分别为99.08%~102.58%和100.21%~103.16%。该方法准确、简便、可靠、有效,可用于盐酸普萘洛尔片对映体的质量控制。

关键词: 正相液相色谱 手性分离 测定 普萘洛尔对映体

**Abstract:** A rapid and sensitive method was developed and validated using a normal phase liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS) for determination of propranolol enantiomers in pharmaceuticals. Sample preparation involved a single extraction step by the addition of methanol. Separation of propranolol enantiomers was achieved on a Chiralcel OD-H chiral column using a mobile phase consisting of n-hexane-ethanol-ammonia (70:30:0.4, v/v/v), and the flow rate was 0.40 mL/min for 20 min. The analyte was monitored by tandem mass spectrometry with electrospray positive ionization in multiple reaction monitoring (MRM) mode, using the transitions of m/z 260.2→116.0. Propranolol enantiomers can be completely separated. The linear range was 2.5~1000 μg/L, and the limit of quantification (LOQ) was 2.5 μg/L. The values for within day and between day precisions and accuracies were well within the generally accepted criteria for analytical methods. The relative standard deviations (RSDs) were less than 2.64%, and the recoveries of the two enantiomers were 99.08%~102.58% and 100.21%~103.16%, respectively. The separation method is accurate, convenient, reliable, efficient, and can be subsequently used for quality control of propranolol enantiomers in pharmaceuticals.

**Keywords:** normal phase liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS) chiral separation determination propranolol enantiomers

Received 2010-09-05; published 2011-01-21

Fund:

国家自然科学基金项目(No. 20775089)和甘肃省科技支撑计划项目(No. 1011FKCA144)。

Corresponding Authors: .1123.2011.00026 \*通讯联系人: 王荣,教授,主要研究方向为手性药物分析及体内药物分析. E-mail: wangrong-69@163.com. 贾正平,教授,博士生导师,主要研究方向为手性药物分析及分子药理学. Email: wangrong-69@163.com

引用本文:

张娟红<sup>1,2</sup>, 王荣<sup>1,2\*</sup>, 谢华<sup>1</sup>, 孟宪栋<sup>1</sup>, 贾正平<sup>1,2\*</sup>, 马骏<sup>1</sup>, 张军莉<sup>1</sup>, 王娟<sup>1</sup>.正相液相色谱-串联质谱法分离普萘洛尔对映体[J] 色谱, 2011,V29(01): 26-30

ZHANG Juanhong<sup>1,2</sup>, WANG Rong<sup>1,2\*</sup>, XIE Hua<sup>1</sup>, MENG Xiandong<sup>1</sup>, JIA Zhengping<sup>1,2\*</sup>, MA Jun<sup>1</sup>, ZHANG Junli<sup>1</sup>, WANG Juan<sup>1</sup>.Enantiomeric separation of propranolol by normal phase chiral liquid chromatography coupled with tandem mass spectrometry[J] Chinese Journal of Chromatography, 2011,V29(01): 26-30

链接本文:

<http://www.chrom-china.com/CN/10.3724/SP.J.1123.2011.00026> 或 <http://www.chrom-china.com/CN/Y2011/V29/I01/26>

## Service

- ▶ 把本文推荐给朋友
- ▶ 加入我的书架
- ▶ 加入引用管理器
- ▶ Email Alert
- ▶ RSS

## 作者相关文章

- ▶ 王荣