

#### arXiv.org > astro-ph > arXiv:1010.5782

Search or Article-id

(Help | Advanced search) All papers - Go!

## **Download:**

- PDF
- PostScript
- Other formats

Current browse context: astro-ph.IM

< prev | next >

new | recent | 1010

Change to browse by:

astro-ph astro-ph.CO

#### **References & Citations**

- SLAC-SPIRES HEP (refers to | cited by)
- NASA ADS

Bookmark(what is this?)

Astrophysics > Instrumentation and Methods for Astrophysics

# The first release of data from the Herschel ATLAS: the SPIRE images

E. Pascale, R. Auld, A. Dariush, L. Dunne, S. Eales, S. Maddox, P.
Panuzzo, M. Pohlen, D. J. B. Smith, S. Buttiglione, A. Cava, D. L.
Clements, A. Cooray, S. Dye, G. de Zotti, J. Fritz, R. Hopwood, E. Ibar,
R. J. Ivison, M. J. Jarvis, L. Leeuw, M. López-Caniego, E. Rigby, G.
Rodighiero, D. Scott, M. W. L. Smith, P. Temi M. Vaccari, I. Valtchanov
(Submitted on 27 Oct 2010)

We have reduced the data taken with the Spectral and Photometric Imaging Receiver (SPIRE) photometer on board the Herschel Space Observatory in the Science Demonstration Phase (SDP) of the Herschel Astrophysical Terahertz Large Area Survey (H-ATLAS). We describe the data reduction, which poses specific challenges, both because of the sheer size of the data, and because only two scans are made for each region. We implement effective solutions to process the bolometric timelines into maps, and show that correlations among detectors are negligible, and that the photometer is stable on time scales up to 250 s. This is longer than the time the telescope takes to cross the observed sky region, and it allows us to use naive binning methods for an optimal reconstruction of the sky emission. The maps have equal contribution of confusion and white instrumental noise, and the latter is estimated to 5.3, 6.4, and 6.7 mJy/beam (1-{\sigma}), at 250, 350, and 500 \mu{m}, respectively. This pipeline is used to reduce other H-ATLAS observations, as they became available, and we discuss how it can be used with the optimal map maker implemented in the Herschel Interactive Processing Environment (HIPE), to improve computational efficiency and stability. The SDP dataset is available from this http URL

Comments:8 pages, 4 figures, submitted to MNRASSubjects:Instrumentation and Methods for Astrophysics (astro-ph.IM);<br/>Cosmology and Extragalactic Astrophysics (astro-ph.CO)Cite as:arXiv:1010.5782v1 [astro-ph.IM]

### **Submission history**

From: Enzo Pascale [view email] [v1] Wed, 27 Oct 2010 20:00:03 GMT (246kb) Link back to: arXiv, form interface, contact.