



Nuclear Theory

Nuclear structure in strong magnetic fields: nuclei in the crust of a magnetar

[Daniel Pena Arteaga](#), [Marcella Grasso](#), [Elias Khan](#), [Peter Ring](#)

(Submitted on 26 Jul 2011)

Covariant density functional theory is used to study the effect of strong magnetic fields, up to the limit predicted for neutron stars (for magnetars $B \approx 10^{18} \text{G}$), on nuclear structure. All new terms in the equation of motion resulting from time reversal symmetry breaking by the magnetic field and the induced currents, as well as axial deformation, are taken into account in a self-consistent fashion. For nuclei in the iron region of the nuclear chart it is found that fields in the order of magnitude of 10^{17}G significantly affect bulk properties like masses and radii.

Comments: 9 pages, 6 figures

Subjects: **Nuclear Theory (nucl-th)**; Solar and Stellar Astrophysics (astro-ph.SR)

Cite as: [arXiv:1107.5243 \[nucl-th\]](#)
(or [arXiv:1107.5243v1 \[nucl-th\]](#) for this version)

Submission history

From: Daniel Pena Arteaga [[view email](#)]

[v1] Tue, 26 Jul 2011 15:28:46 GMT (837kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nucl-th

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[astro-ph](#)

[astro-ph.SR](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark ([what is this?](#))

