

Browse

or

Submit

Why isn't the universe as bright as it should be?

Study explains why galaxies don't churn out as many stars as they should.

Jennifer Chu | MIT News Office
March 4, 2015

Press Inquiries

PRESS MENTIONS

A handful of new stars are born each year in the Milky Way, while many more blink on across the universe. But astronomers have observed that galaxies should be churning out millions more stars, based on the amount of interstellar gas available.

Now researchers from MIT, Columbia University, and Michigan State University have pieced together a theory describing how clusters of galaxies may regulate star formation. They describe their framework this week in the journal *Nature*.

When intracluster gas cools rapidly, it condenses, then collapses to form new stars. Scientists have long thought that something must be keeping the gas from cooling enough to generate more stars — but exactly what has remained a mystery.

MIT researchers have found that the high temperature of intracluster gas, which condenses to form stars, may be hindering the development of new stars, reports Jonathan O'Callaghan for the *Daily Mail*. The researchers hope to use the new findings to better understand how stars form in surrounding galaxies.

Daily Mail

RELATED

Michael McDonald

For some galaxy clusters, the researchers say, the intracluster gas may simply be too hot — on the order of hundreds of millions of degrees Celsius. Even if one region experiences some cooling, the intensity of the surrounding heat would keep that region from cooling further — an effect known as conduction.

“It would be like putting an ice cube in a boiling pot of water — the average temperature is pretty much still boiling,” says Michael McDonald, a Hubble Fellow in MIT’s Kavli Institute for Astrophysics and Space Research. “At super-high temperatures, conduction smooths out the temperature distribution so you don’t get any of these cold clouds that should form stars.”

For so-called “cool core” galaxy clusters, the gas near the center may be cool enough to form some stars. However, a portion of this cooled gas may rain down into a central black hole, which then spews out hot material that serves to reheat the surroundings, preventing many stars from forming — an effect the team terms “precipitation-driven feedback.”

“Some stars will form, but before it gets too out of hand, the black hole will heat everything back up — it’s like a thermostat for the cluster,” McDonald says. “The combination of conduction and precipitation-driven feedback provides a simple, clear picture of how star formation is governed in galaxy clusters.”

Crossing a galactic threshold

Throughout the universe, there exist two main classes of galaxy clusters: cool core clusters — those that are rapidly cooling and forming stars — and non-cool core clusters — those have not had sufficient time to cool.

The Coma cluster, a non-cool cluster, is filled with gas at a scorching 100 million degrees Celsius. To form any stars, this gas would have to cool for several billion years. In contrast, the nearby Perseus cluster is a cool core cluster whose intracluster gas is a relatively mild several million degrees Celsius. New stars occasionally emerge from the cooling of this gas in the Perseus cluster, though not as many as scientists would predict.

“The amount of fuel for star formation outpaces the amount of stars 10 times, so these clusters should be really star-rich,” McDonald says. “You really need some mechanism to prevent gas from cooling, otherwise the universe would have 10 times as many stars.”

McDonald and his colleagues worked out a theoretical framework that relies on two anti-cooling mechanisms.

The group calculated the behavior of intracluster gas based on a galaxy cluster’s radius, mass, density, and temperature. The researchers found that there is a critical temperature threshold below which the cooling of gas accelerates significantly, causing gas to cool rapidly enough to form stars.

MIT Kavli Institute for Astrophysics and Space Research

School of Science


ARCHIVES



Most massive and luminous galaxy cluster identified



Researchers detect brightest pulsar ever recorded



Researchers identify one of the earliest stars in the universe



When the first stars blinked on

According to the group's theory, two different mechanisms regulate star formation, depending on whether a galaxy cluster is above or below the temperature threshold. For clusters that are significantly above the threshold, conduction puts a damper on star formation: The surrounding hot gas overwhelms any pockets of cold gas that may form, keeping everything in the cluster at high temperatures.

"For these hotter clusters, they're stuck in this hot state, and will never cool and form stars," McDonald says. "Once you get into this very high-temperature regime, cooling is really inefficient, and they're stuck there forever."

For gas at temperatures closer to the lower threshold, it's much easier to cool to form stars. However, in these clusters, precipitation-driven feedback starts to kick in to regulate star formation: While cooling gas can quickly condense into clouds of droplets that can form stars, these droplets can also rain down into a central black hole — in which case the black hole may emit hot jets of material back into the cluster, heating the surrounding gas back up to prevent further stars from forming.

"In the Perseus cluster, we see these jets acting on hot gas, with all these bubbles and ripples and shockwaves," McDonald says. "Now we have a good sense of what triggered those jets, which was precipitating gas falling onto the black hole."

On track

McDonald and his colleagues compared their theoretical framework to observations of distant galaxy clusters, and found that their theory matched the observed differences between clusters. The team collected data from the Chandra X-ray Observatory and the South Pole Telescope — an observatory in Antarctica that searches for far-off massive galaxy clusters.

The researchers compared their theoretical framework with the gas cooling times of every known galaxy cluster, and found that clusters filtered into two populations — very slowly cooling clusters, and clusters that are cooling rapidly, closer to the rate predicted by the group as a critical threshold.

By using the theoretical framework, McDonald says researchers may be able to predict the evolution of galaxy clusters, and the stars they produce.

"We've built a track that clusters follow," McDonald says. "The nice, simple thing about this framework is that you're stuck in one of two modes, for a very long time, until something very catastrophic bumps you out, like a head-on collision with another cluster."

The researchers hope to look deeper into the theory to see whether the mechanisms regulating star formation in clusters also apply to individual galaxies. Preliminary evidence, he says, suggests that is the case.

“If we can use all this information to understand why or why not stars form around us, then we’ve made a big step forward,” McDonald says.

“[These results] look very promising,” says Paul Nulsen, an astronomer at the Harvard-Smithsonian Center for Astrophysics who was not involved in this research. “More work will be needed to show conclusively that precipitation is the main source of the gas that powers feedback. Other processes in the feedback cycle also need to be understood. For example, there is still no consensus on how the gas falling into a massive black hole produces energetic jets, or how they inhibit cooling in the remaining gas. This is not the end of the story, but it is an important insight into a problem that has proved a lot more difficult than anyone ever anticipated.”

This research was funded in part by the National Science Foundation and NASA.

Topics: [Kavli Institute](#) [Astronomy](#) [Astrophysics](#) [Physics](#)

[Black holes](#) [Stars](#) [Research](#) [space](#)

[Space, astronomy and planetary science](#)

[National Science Foundation \(NSF\)](#) [School of Science](#)

Comments

[View the discussion thread.](#)

About This Website

This Website is maintained by the MIT News Office.

[About the MIT News Office](#)

[MIT News Press Center](#)

[Press Inquiries](#)

[Filming Guidelines](#)

[Contact Us](#)

[Terms of Use](#)

[RSS](#)

[Twitter](#)

[Facebook](#)

[Google+](#)

[Instagram](#)

[Flickr](#)

[MIT Homepage](#)

[MIT Video](#)

[MIT Connect](#)

[MIT Resources](#)

[Events Calendar](#)

Resources

[Submit Campus News](#)

[Pitch a Story](#)

[Sign Up for Newsletter](#)

[YouTube](#)

[About MIT](#)

[Admissions](#)

[Alumni](#)

[Education](#)

[Labs and Centers](#)

[Subscribe to Press Releases](#)

[Terms of Use](#)

MIT News Office • Building 11-400

Massachusetts Institute of Technology • Cambridge, MA 02139-4307

[GIVE TO MIT](#)