

Science News

from research organizations

The environment for permafrost in Daisetsu Mountains in Japan is projected to decrease significantly

Date: July 30, 2021

Source: National Institute for Environmental Studies

Summary: Areas with climatic conditions suitable for sustaining permafrost in the Daisetsu Mountains are projected. The size of the area in the Daisetsu Mountains where climatic conditions were suitable for permafrost were estimated to be approximately 150 km² in 2010. Under the business-as-usual scenario, this area is projected to disappear by around 2070. Under the low-carbon scenario consistent with Paris target scenario, the area is projected to decrease to approximately 13% of 2010 by 2100.

Share: [!\[\]\(17413706fd4997a1a4bdf85c6864eee1_img.jpg\)](#) [!\[\]\(f419710cbe076aa30a9c6c031b5cbe84_img.jpg\)](#) [!\[\]\(2726020a4107bdc9042b257034f90eb3_img.jpg\)](#) [!\[\]\(9459655bf14a84f4d775e8d814cca8c9_img.jpg\)](#) [!\[\]\(de47dbdca34225b222a4a87ac0e499b3_img.jpg\)](#)

FULL STORY

Areas with ground temperatures that remain below 0°C for more than two years are referred to as permafrost, and approximately one quarter of the Northern Hemisphere and 17% of the Earth's exposed land surface is permafrost. Permafrost is found in mountainous areas as well as in high-latitude tundra and taiga regions. Recent observations have shown that the permafrost in mountainous areas is thawing in the European Alps, Scandinavia and on the Tibetan Plateau.

The islands that make up Japan form a long arc that extends from tropical regions in the south to Palearctic regions in the north. Permafrost has been reported to exist in the Daisetsu Mountains, Mt. Fuji and Mt. Tateyama. The Daisetsu Mountains on the northern island of Hokkaido, which are known to support a variety of alpine plants and animals, are also a valuable natural resource that is visited by numerous climbers every year. However, there is some concern that the thawing of permafrost will have a major impact on this mountain ecosystem. No projections of future permafrost distribution have been undertaken in East Asia, including in Japan, to date.

In this study, by applying the statistical method developed in a previous study for estimating the permafrost distribution to outputs from bias-corrected and downscaled climate model with a 1 km resolution for the area of interest, we investigated the climatic conditions required to sustain permafrost in the Daisetsu Mountains.

Researchers have studied the climate conditions suitable to maintain permafrost in the Daisetsu Mountains. The region of the climate conditions suitable to maintain permafrost starts to decrease significantly in size from around 2000, and under both the low-carbon scenario consistent with Paris target (RCP2.6) and business-as-

usual scenario (RCP8.5). The steep decline in the region at the end of the 20th century occurs because the surface air temperature increases significantly from around the year 2000 onward. The size of the area in the Daisetsu Mountains where climatic conditions were suitable for permafrost were estimated to be approximately 150 km² in 2010. Under the RCP8.5 scenario, this area is projected to disappear by around 2070. Under the RCP2.6 scenario, the area is projected to decrease to approximately 20 km² by 2100.

The finding that the climatic conditions of the Daisetsu Mountains are projected to shift towards conditions where permafrost will disappear regardless of the climate scenario used, indicates the importance of impact assessments and developing adaptation measures to climate change. Thawing of permafrost can have a significant impact on the surface vegetation and alpine ecosystems. Previous studies have shown that the thawing of permafrost can have a variety of adverse effects, including lowering groundwater levels and replacement of alpine meadows by steppes, extensive desertification such as that in the eastern and western parts of the Tibetan Plateau, and a decrease in species diversity due to warming. Moreover, thawing of the frozen ground reduces the stability of the ground in mountain regions, potentially increasing the frequency and magnitude of rock falls and landslides, which may affect the safety of trekkers that visit the Daisetsu Mountains annually. In order to deal with these problems, it is very important to monitor the environmental changes in mountainous areas. In addition to accurately monitoring changes in mountain environments, providing local governments with appropriate measures to prepare for major future environmental changes, as shown in this study, is an important issue for the future.

Story Source:

Materials provided by **National Institute for Environmental Studies**. *Note: Content may be edited for style and length.*

Journal Reference:

1. Tokuta Yokohata, Go Iwahana, Toshio Sone, Kazuyuki Saito, Noriko N. Ishizaki, Takahiro Kubo, Hiroyuki Oguma, Masao Uchida. **Projections of surface air temperature required to sustain permafrost and importance of adaptation to climate change in the Daisetsu Mountains, Japan.** *Scientific Reports*, 2021; 11 (1) DOI: 10.1038/s41598-021-94222-4

Cite This Page:

MLA	APA	Chicago
-----	-----	---------

National Institute for Environmental Studies. "The environment for permafrost in Daisetsu Mountains in Japan is projected to decrease significantly." ScienceDaily. ScienceDaily, 30 July 2021. <www.sciencedaily.com/releases/2021/07/210730165447.htm>.

RELATED STORIES

- Climate Warming to Increase Carbon Loss in Canadian Peatland by 103 Per Cent
June 7, 2021 — Carbon loss in Canadian peatland is projected to increase by 103 per cent under a high emission scenario, according to new ...
- Colombia Could Lose 60% of Land Suitable for Irrigated Rice Due to Climate Change
May 29, 2019 — Without significant global reductions in greenhouse gas emissions, Colombia will have 60% less land suitable for rice production by the 2050s. Due primarily to increased temperatures and decreased ...

Degrading Permafrost Puts Arctic Infrastructure at Risk by Mid-Century

Dec. 11, 2018 — Seventy percent of the current infrastructure in the Arctic has a high potential to be affected by thawing permafrost in the next 30 years. Even meeting the climate change targets of the Paris ...

Warmer Future for the Pacific Northwest If Carbon Dioxide Levels Rise, Climate Projections Show

Feb. 20, 2018 — In the midst of an unseasonably warm winter in the Pacific Northwest, a comparison of four publicly available climate projections has shown broad agreement that the region will become considerably ...

FROM AROUND THE WEB

ScienceDaily shares links with sites in the TrendMD network and earns revenue from third-party advertisers, where indicated.

Climatic suitable area analysis and response to climate change of *Actinidia arguta* in China*

Jinpeng ZHAO et al., Chinese Journal of Eco-Agriculture

Impact of climate change on allowable bearing capacity on the Qinghai-Tibetan Plateau

Xiao-Ming Xu et al., Advances in Climate Change Research, 2019

Temperature and precipitation projection at 1.5 and 2°C increase in global mean temperature

Ting HU et al., Chinese Science Bulletin, 2017

Future Changes in the Impact of North Pacific Midlatitude Oceanic Frontal Intensity on the Wintertime Storm Track in CMIP5 Models

Yao Yao et al., Journal of Meteorological Research, 2020

Responses and changes in the permafrost and snow water equivalent in the Northern Hemisphere under a scenario of 1.5 °C warming

Ying Kong et al., Advances in Climate Change Research, 2017

Akonni Biosystems Submits Multiplex Diagnostics System to FDA

360Dx, 2019

Modeling the carbon dynamics of alpine grassland in the Qinghai-Tibetan Plateau under scenarios of 1.5 and 2 °C global warming

Shu-Hua Yi et al., Advances in Climate Change Research, 2019

MFB Fertility Progesterone Ovulation Test Gets FDA Clearance

Doug Macron et al., 360Dx, 2020

Powered by **TREND MD**

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

✉ Email Newsletters

📡 RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

 Facebook

 Twitter

 LinkedIn

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 Leave Feedback

 Contact Us

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2021 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

— [CCPA: Do Not Sell My Information](#) — — [GDPR: Privacy Settings](#) —