

## Science News

from research organizations

### What's killing coral reefs in Florida is also killing them in Belize Study of Belize Barrier Reef Shows Nitrogen Enrichment from Land-based Sources Doubled in Four Decades

*Date:* August 3, 2021

*Source:* Florida Atlantic University

*Summary:* Only 17 percent of live coral cover remains on fore-reefs in Belize. A study finds new evidence that nitrogen enrichment from land-based sources like agriculture run-off and sewage, are significantly driving macroalgal blooms to increase on the Belize Barrier Reef and causing massive decline in hard coral cover. With only 2 percent of hard coral cover remaining in the Florida Keys National Marine Sanctuary, it's too late to save that reef, but there's still hope for the Belize Barrier Reef.

*Share:* [!\[\]\(17413706fd4997a1a4bdf85c6864eee1\_img.jpg\)](#) [!\[\]\(f419710cbe076aa30a9c6c031b5cbe84\_img.jpg\)](#) [!\[\]\(2726020a4107bdc9042b257034f90eb3\_img.jpg\)](#) [!\[\]\(9459655bf14a84f4d775e8d814cca8c9\_img.jpg\)](#) [!\[\]\(de47dbdca34225b222a4a87ac0e499b3\_img.jpg\)](#)

#### FULL STORY

In 1842, Charles Darwin described the Belize Barrier Reef as "the most remarkable reef in the West Indies." Fast forward to 2021, only about 17 percent of live coral cover remains on fore-reefs in Belize. Overfishing, resulting in reduced grazing of algae, has long been blamed for adversely impacting this globally significant ecosystem along the Caribbean coast of Central America. Designated in part as a UNESCO World Heritage Site in 1996, the Mesoamerican reef is second in length only to the iconic Great Barrier Reef in Australia.

A first-of-its-kind study provides evidence that reduced grazing is not solely responsible for coral death in this barrier reef and the proof is in the chemistry. Instead, nitrogen enrichment from land-based sources is significantly driving macroalgal blooms to increase on the Belize Barrier Reef and causing massive decline in hard coral cover. With only 2 percent of hard coral cover remaining in the Florida Keys National Marine Sanctuary, it's too late to save that reef, but there is still hope for the Belize Barrier Reef.

To assess the level of nutrient enrichment and eutrophication in the coastal waters of the Belize Barrier Reef due to human impacts on watersheds and subsequent management of these nutrients, researchers used benthic macroalgae as cumulative bioindicators of nutrient enrichment.

Brian Lapointe, Ph.D., co-author and a senior research professor from Florida Atlantic University's Harbor Branch Oceanographic Institute, and collaborators from the Wildlife Conservation Society, Belize Program, analyzed the carbon (C), nitrogen (N) and phosphorus (P) content and molar C:N:P ratios of baseline tissue

from the 1980s and compared them with more recent samples collected in 2017 and 2019, with the recent data also being analyzed for stable isotopes of C and N.

Results of the study, published in the journal *Marine Pollution Bulletin*, suggest that N:P ratios have doubled over the past 35 years within the main Belize Barrier Reef lagoon, similar to the three-fold increases in N:P of macroalgae at Looe Key reef in the lower Florida Keys since the 1980s. The study provides an expansive geographic scope of potential nutrient impacts in the northern and central lagoon, while also using a unique, local historical baseline in the central lagoon to gauge the change in nutrient availability across the Belize Barrier Reef over the past four decades.

"Our study is the first to suggest that nutrient stress caused by nitrogen enrichment and elevated N:P ratios, both detrimental to hard coral physiology, is a primary mechanism by which human activities within watersheds and coastal environments of Belize are negatively impacting the barrier reef," said Lapointe. "This is perhaps most acute for the health and biodiversity of hard corals that are the critical ecosystem engineers of tropical reefs. The impacts of deforestation, agriculture run-off, sewage, increasing numbers and capacity of cruise ships and tourism infrastructure, effluent from large-scale aquaculture production and large amounts of nutrient-rich ash from forest fires in the region are contributing to overall declining water quality and incidents of hyper-eutrophication in areas such as the Placencia Lagoon and the New River, Belize."

Lapointe and co-authors Alexander Tewfik, Ph.D., a marine ecologist who served as senior conservation scientist from 2014 to 2019 for the Wildlife Conservation Society and is currently a consultant for the organization; and Myles Phillips, MSc., technical coordinator, Wildlife Conservation Society, examined the relationships between the C, N, P contents, stoichiometry (C:N:P) and stable isotopes of C and N found within the tissue of a variety of common macroalgae species sampled in the lower riverine, estuarine and shallow marine environments from around Belize City to the offshore fore-reef habitats of the Belize Barrier Reef.

They also compared their Belize City transect data from 2019, which included the Swallow Caye Wildlife Sanctuary declared to protect the endangered West Indian manatee and lying just 3 kilometers from Belize City, to similar elemental content (N, P) and stoichiometry data collected from macroalgae by the team in the South Water Caye Marine Reserve in 2017 and during the 1980s by Lapointe.

Researchers found that increasing nutrients, especially N, from the Belize River and Haulover Creek (a side channel and outfall of the Belize River), sustain macroalgal blooms from nearshore areas of the main lagoon to the offshore fore-reef. Significant offshore gradients of C:N:P in macroalgae all indicated land-based sources of these key nutrients; C:N and C:P ratios of macroalgae were lower in nearshore waters with lower salinity values, pointing to freshwater runoff as the source of nutrient enrichment; and N values of macroalgae were overall enriched well above values for N<sub>2</sub> fixation, especially in near-shore waters, where values matched those reported for sewage pollution.

N:P ratios of macroalgae were elevated across the entire Belize City nearshore-to-offshore gradient to values of approximately 70:1, some four-fold higher than the Redfield Ratio (16:1) for oceanic waters and more than two-fold higher than values for macroalgae on the Belize Barrier Reef in the mid-1980s (about 30:1). This indicates widespread N-enrichment, increasing N:P ratios and a strengthening of P-limitation, which also is consistent with patterns of pelagic *Sargassum* across large areas of the Atlantic basin and Caribbean Sea, which have plagued the nearshore waters and beaches of the region with increased biomass since 2011.

"The mouth of Haulover Creek was clearly the most polluted site examined in our study, as it drains directly through the commercial center of the Belize City watershed that is heavily impacted by untreated sewage," said Tewfik. "In addition, effluent from the processing of hundreds of metric tons of queen conch and Caribbean spiny lobster, the two most important fisheries exports in Belize, entering Haulover Creek is contributing to the negative effects of poor water quality in the lagoon."

To mitigate the environmental impacts on the Belize Barrier Reef, the researchers say that a number of actions could be taken to reduce nutrient loading from urban effluents and agricultural run-off to improve water quality and restore the shallow coastal ecosystems of this globally significant coral reef ecosystem. They recommend to continue efforts in compiling ecological assessments of impacts, standardize appropriate management

practices and water quality monitoring protocols, and adopt a focused ridge-to-reef conservation strategy as well as increasing awareness of often "invisible" nutrient contamination among the general public and political bodies.

"Eutrophication is an important and widespread problem that is recognized by coral reef scientists around the world," said Lapointe. "Our latest data provide further evidence that it is unlikely that simply controlling fishing practices, even in light of climate change, will enhance resilience of Belize's coral reefs or any other coral reefs being impacted by eutrophication. The reality of this large-scale coastal pollution must be rigorously confronted by scientists and resource managers alike in Belize, the United States and world-wide."

This research was supported by NASA's Biological Diversity and Ecological Forecasting Program under contract number 80NSSC19K0200 (Climate-influenced Nutrient Flows and Threats to the Biodiversity of the Belize Barrier Reef Reserve System), Coastal Association of Science and Technology (COAST) and FAU Harbor Branch. A number of additional staff of Wildlife Conservation Society, Belize aided with all field logistics, initial sample preparations and project support.

---

### Story Source:

Materials provided by **Florida Atlantic University**. Original written by Gisele Galoustian. *Note: Content may be edited for style and length.*

---

### Journal Reference:

1. Brian E. Lapointe, Alexander Tewfik, Myles Phillips. **Macroalgae reveal nitrogen enrichment and elevated N:P ratios on the Belize Barrier Reef**. *Marine Pollution Bulletin*, 2021; 171: 112686 DOI: 10.1016/j.marpolbul.2021.112686

---

### Cite This Page:

MLA

APA

Chicago

---

Florida Atlantic University. "What's killing coral reefs in Florida is also killing them in Belize: Study of Belize Barrier Reef Shows Nitrogen Enrichment from Land-based Sources Doubled in Four Decades." ScienceDaily. ScienceDaily, 3 August 2021. <[www.sciencedaily.com/releases/2021/08/210803105534.htm](http://www.sciencedaily.com/releases/2021/08/210803105534.htm)>.

---

### RELATED STORIES

#### Turning the Tables -- How Table Corals Are Regenerating Reefs

June 1, 2021 — New research shows table corals can regenerate coral reef habitats on the Great Barrier Reef decades faster than any other coral type. The research suggests overall reef recovery would slow ...

#### Acropora Spp. Coral Still Thrives in the Holdout Refuge of Coral Gardens, Belize

Sep. 30, 2020 — Coral Gardens Reef in Belize remains a refuge for Acropora spp. coral despite widespread devastation in other areas of the western North Atlantic/Caribbean, according to a new ...

#### Climate Change, Human Activity Lead to Nearshore Coral Growth Decline

Aug. 28, 2019 — New research compares the growth rates between nearshore and offshore corals in the Belize Mesoamerican Barrier Reef System, the world's second-largest reef system. While nearshore corals have ...

#### Thirty Years of Unique Data Reveal What's Really Killing Coral Reefs

July 15, 2019 — Coral bleaching is not just due to a warming planet, but also a planet that is simultaneously being enriched with reactive nitrogen from sources like improperly treated sewage, and fertilizers. ...

## FROM AROUND THE WEB

---

*ScienceDaily shares links with sites in the TrendMD network and earns revenue from third-party advertisers, where indicated.*

## Free Subscriptions

---

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 [Email Newsletters](#)

 [RSS Feeds](#)

## Follow Us

---

Keep up to date with the latest news from ScienceDaily via social networks:

 [Facebook](#)

 [Twitter](#)

 [LinkedIn](#)

## Have Feedback?

---

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 [Leave Feedback](#)

 [Contact Us](#)

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2021 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

— [CCPA: Do Not Sell My Information](#) — — [GDPR: Privacy Settings](#) —