Your source for the latest research news

Science News

from research organizations

Semi-natural habitat patches complement flower strips in protecting pollinators

Date:	August 3, 2021
Source:	University of Freiburg
Summary:	Ecologists observe the diversity of insects on the edge of apple orchards on Lake Constance.
Share:	f 🌶 🦻 in 🗖
FULL STORY	

At the moment, many flower strips are buzzing and humming: cornflowers, poppies, wild carrots and many other flowers attract numerous insects. The field edges covered by these flowers typically bloom between mid-May and mid-August. Complementary habitats are needed to support pollinator insects in agricultural landscapes throughout the year. Semi-natural small structures, such as ditches, banks, hedgerows, or overgrown fences, could provide such a complement. "Researchers have already shown many times how important natural habitats are for pollinators. Almost always, however, only large-scale structures have been researched for this purpose, for example, wide meadows or pastures. Studies on what small structures mean for pollinators and which species particularly benefit from them are rare," says Vivien von Königslöw from the Institute of Earth and Environmental Sciences at the University of Freiburg. As a result, together with Dr. Anne-Christine Mupepele and Prof. Dr. Alexandra-Maria Klein, she studied flower strips as well as semi-natural habitat patches in the Lake Constance area over a period of two years, a place in which there is a particular interest in pollinating insects due to large-scale fruit cultivation. The researchers published their results in the journal Biological Conservation.

Semi-natural habitats attract more bees

"Our goal was to find out how the diversity of wild bees and hoverflies can be promoted in the vicinity of largescale orchards," says von Königslöw. To do this, their study compared the occurrence of bees and hoverflies in flower strips and in existing flower-rich habitats, each located on the edge of conventional apple orchards in southern Germany. Their analysis showed that the different flowering times and plant species in the seminatural habitats, such as hedgerows and small groves, mainly benefit solitary and oligolectic bees, i.e. those that collect only one pollen species. The existing biotope areas attracted bee species with a different pollen specialization than the sown flower strips. At the same time, the researchers found a greater number of pollinators in the flower strips and counted more species than in the small structures. "Thus, semi-natural habitats complement existing flower strips," von Königslöw concludes.

For their research, the ecologists established flower strips at the edge of private orchards in 2018. Seminatural small structures, including drainage ditches, embankments and overgrown fences, were already in place. The researchers monitored the bees and hoverflies at least once a month from spring to late summer.

Effective and cost-efficient

"Semi-natural habitat patches can play an important role in protecting pollinators because they help ensure that flowers are available all year round," says Klein, head of the Chair of Nature Conservation and Landscape Ecology at the University of Freiburg. They also provide potential retreats and nesting sites, which are important for overwintering bumblebees, for example. "For effective and cost-efficient protection of pollinating insects, the focus should not only be on flower strips," Klein concludes. "Existing small structures of spontaneous vegetation, plant species that grow on their own from existing seeds in the soil, are also attractive to insects and should be preserved." The Freiburg scientist explains that at present, however, there are hardly any incentives for farmers to develop and preserve small semi-natural habitat patches.

Story Source:

Materials provided by University of Freiburg. Note: Content may be edited for style and length.

Journal Reference:

 Vivien von Königslöw, Anne-Christine Mupepele, Alexandra-Maria Klein. Overlooked jewels: Existing habitat patches complement sown flower strips to conserve pollinators. *Biological Conservation*, 2021; 261: 109263 DOI: 10.1016/j.biocon.2021.109263

Cite This Page:

APA Chicago

MLA

University of Freiburg. "Semi-natural habitat patches complement flower strips in protecting pollinators." ScienceDaily. ScienceDaily, 3 August 2021. <www.sciencedaily.com/releases/2021/08/210803121337.htm>.

RELATED STORIES

Plant Provenance Influences Pollinators

Apr. 22, 2021 — Insect decline poses challenges for the pollination of wild and cultivated plants. Landscape ecologists have now discovered that the diversity of insects that interact with plant communities is ...

Cichlid Fishes from African Lake Tanganyika Shed Light on How Organismal Diversity Arises Nov. 18, 2020 — Lake Tanganyika in Africa is a true hotspot of organismal diversity. Approximately 240 species of cichlid fishes have evolved in this lake in less than 10 million years. A research team has ...

Newly Discovered Pathogen in NY Apples Causes Bitter Rot Disease

July 6, 2020 — In a study of New York state apple orchards, plant pathologists have identified a new fungal pathogen that causes bitter rot disease in ...

Orchards in Natural Habitats Draw Bee Diversity, Improve Apple Production

Jan. 17, 2019 — Apple orchards surrounded by agricultural lands are visited by a less diverse collection of bee species than orchards surrounded by natural ...

FROM AROUND THE WEB

ScienceDaily shares links with sites in the TrendMD network and earns revenue from third-party advertisers, where indicated.

Geographic mosaic theory of coevolution in plant and pollinator interaction: Process, evidence, prospect

Xiaoyue Wang et al., Chinese Science Bulletin, 2019

Distribution patterns of pollination insect community under different agricultural landscape context

Yameng LIU et al., Acta Ecologica Sinica, 2018

Progress in pollination ecology at the community level Qiang FANG et al., Chinese Science Bulletin, 2014

Study on the features of butterfly pollinator communities in crucial ecological regions of the Pearl River Delta 徐安乐 et al., Journal of Environmental Entomology, 2020

Reproduction biology of mimosa bimucronata – a ruderal species L.A Silva et al., Advances in Weed Science, 2011

Acute Lymphoblastic Lukemia: Outcomes for Adolescence MRP, 2020

Addressing Lung Cancer Screening Disparities: Panel Issues Recommendations Cardiology Advisor, 2020

Nectar Secretion Characteristics and Their Effects on Insect Pollination of Goodyera foliosa, A Nectar Rewarding Orchid

Nan-nan LIU et al., Journal of Tropical and Subtropical Botany

Powered by TREND MD

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

Email Newsletters

A RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

- f Facebook
- ✓ Twitter
- in LinkedIn

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

- Leave Feedback
- Contact Us

About This Site | Staff | Reviews | Contribute | Advertise | Privacy Policy | Editorial Policy | Terms of Use

Copyright 2021 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners. Financial support for ScienceDaily comes from advertisements and referral programs, where indicated. — CCPA: Do Not Sell My Information — — GDPR: Privacy Settings —