

Science News

from research organizations

Butterflies and moths have difficulty adjusting to a rapidly changing climate

Date: June 10, 2021

Source: University of Helsinki

Summary: Climate change exerts great pressure for change on species and biodiversity. A recent study indicates that the few moth and butterfly species (Lepidoptera) capable of adjusting to a changing climate by advancing their flight period and moving further north have fared the best in Finland. In contrast, roughly 40% of Lepidoptera species have not been able to respond in either way, seeing their populations decline.

Share: [!\[\]\(17413706fd4997a1a4bdf85c6864eee1_img.jpg\)](#) [!\[\]\(f419710cbe076aa30a9c6c031b5cbe84_img.jpg\)](#) [!\[\]\(2726020a4107bdc9042b257034f90eb3_img.jpg\)](#) [!\[\]\(9459655bf14a84f4d775e8d814cca8c9_img.jpg\)](#) [!\[\]\(de47dbdca34225b222a4a87ac0e499b3_img.jpg\)](#)

FULL STORY

Climate change exerts great pressure for change on species and biodiversity. A recent study conducted by the University of Helsinki and the Finnish Environment Institute indicates that the few moth and butterfly species (Lepidoptera) capable of adjusting to a changing climate by advancing their flight period and moving further north have fared the best in Finland. In contrast, roughly 40% of Lepidoptera species have not been able to respond in either way, seeing their populations decline.

Climate change is bringing about rapid change in Finnish nature -- can species keep up with the pace? Adjusting to climate change can manifest through earlier phenology such as moth and butterfly flight periods, bird nesting, or plant flowering taking place earlier than before. Species can also adjust by shifting their range further north, as individuals relocate to new areas where conditions have become favourable.

The researchers emphasise that, to preserve biodiversity as climate change intensifies, it is of utmost importance to ensure sufficiently extensive, interconnected and habitats of high-quality which make it possible for species to adjust to the challenges generated by climate change.

The study carried out by the University of Helsinki and the Finnish Environment Institute compared temporal shifts in the flight period of 289 moths and butterflies and spatial shifts in their northern range boundary, as well as changes in abundance over a roughly 20-year period.

"Roughly 45% of the species that we studied had either moved northward or advanced their flight period," says Postdoctoral Researcher Maria Hällfors from the University of Helsinki. "They fared much better than the 40% of the species which had not responded in either way. On average, the populations of these poorly responding

species had declined. The largest increase in abundance was seen in the 15% of the species that both moved northward and advanced their flight. This demonstrates that the ability to respond to a changing environment is vital for species."

Few species have advanced their phenology

Another interesting finding was the fact that while nearly half of the species had moved northward, only 27% had advanced their flight period.

"This finding deviates from observations made elsewhere in Europe where advancing the flight period has been much more common among Lepidoptera," says Senior Researcher Juha Pöyry from the Finnish Environment Institute.

In Finland, the species that have advanced their flight the most are the ones that overwinter as adults, including the European peacock butterfly. In fact, it appears that Lepidoptera species living in Finland respond more readily by expanding their ranges northward compared to advancing their flight. Species which are found further north than before include the antler moth and the scarce copper.

"It may be that the increasing light in the spring is a more important cue for butterflies and moths to start their flight than temperature on its own," Pöyry adds.

Sufficient habitats are vital

A potential explanation for the infrequency of species responding optimally, that is, by both advancing their flight and moving northward, could be the scarcity of suitable habitats.

"For organisms to be able to respond to climate change by shifting their range further north, sufficient amounts of suitable habitats of high-quality are needed," says Mikko Kuussaari, Senior Researcher at the Finnish Environment Institute.

The amount of habitats important for many moth and butterfly species has decreased, resulting in population decline for many of them. For example, many butterfly species have suffered from a decrease in meadows.

"Declining populations are usually not able to provide a sufficient basis for the species to spread to new areas. Small populations also contain less genetic diversity that could help the local populations adjust by changing the timing of their flight." Kuussaari adds. Indeed, safeguarding biodiversity requires, above all, the maintenance of sufficiently large and interconnected habitats of high-quality .

Long-term monitoring enables research

The study utilised data on Lepidoptera flight periods collected in two long-term monitoring projects coordinated by the Finnish Environment Institute. Of the two, the Finnish national moth monitoring scheme was launched in 1993 and the butterfly monitoring scheme in agricultural landscapes in 1999. A dataset of citizen observations openly available through the Finnish Biodiversity Information Facility was utilised to calculate species range boundary shifts.

"Without such long-term monitoring schemes and the great contribution of volunteer butterfly and moth enthusiasts in collecting observations, as well as collaboration between different research organisations, it would be impossible to carry out these kinds of analyses encompassing hundreds of species," says Associate Professor Marjo Saastamoinen from the University of Helsinki.

Story Source:

Materials provided by **University of Helsinki**. *Note: Content may be edited for style and length.*

Journal Reference:

1. Maria H. Hällfors, Juha Pöyry, Janne Heliölä, Ilmari Kohonen, Mikko Kuussaari, Reima Leinonen, Reto Schmucki, Pasi Sihvonen, Marjo Saastamoinen. **Combining range and phenology shifts offers a winning strategy for boreal Lepidoptera.** *Ecology Letters*, 2021; DOI: 10.1111/ele.13774

Cite This Page:

MLA

APA

Chicago

University of Helsinki. "Butterflies and moths have difficulty adjusting to a rapidly changing climate." ScienceDaily. ScienceDaily, 10 June 2021. <www.sciencedaily.com/releases/2021/06/210610162410.htm>.

RELATED STORIES

Provide Shady Spots to Protect Butterflies from Climate Change

Sep. 24, 2020 — Researchers have discovered significant variations in the ability of different UK butterfly species to maintain a suitable body temperature. Species that rely most on finding a suitably shady ...

Secrets to Climate Change Adaptation Uncovered in the European Corn Borer Moth

Oct. 10, 2019 — Biologists have found two genes that may permit some insect species to survive climate change by adjusting their biological annual clocks while others succumb. The researchers looked at the European ...

Populations of Widely Spread Tree Species Respond Differently to Climate Change

June 8, 2018 — A new study shows that not all populations of a single, widely spread tree species respond the same to climate change, something scientists will need to consider when making climate change ...

How Land Use and Climate Change Are Driving Species Distribution Shifts

May 7, 2018 — Climate change is altering where species live all over the planet. With global warming, species are moving towards the poles and up elevation where temperature is lower. However, along with global ...

FROM AROUND THE WEB

ScienceDaily shares links with sites in the TrendMD network and earns revenue from third-party advertisers, where indicated.

Distributional responses to climate change for alpine species of *Cyananthus* and *Primula* endemic to the Himalaya-Hengduan Mountains

Xie He et al., *Plant Diversity*, 2019

Advancements of the researches on biodiversity loss mechanisms

FuWen WEI et al., *Chinese Science Bulletin*, 2014

'Natural selection in real time': Which species are most likely to adapt to climate change?

McGill University et al., Genetic Literacy Project

Viewpoint: Conservation isn't enough. We need technology to blunt the impacts of climate change | Genetic Literacy Project

Lauren Anderson, Genetic Literacy Project

Projected impacts of climate change on protected birds and nature reserves in China

Xueyan Li et al., *Science Bulletin*, 2015

Panel Issues Recommendations to Address Lung Cancer Screening Disparities

Cardiology Advisor, 2020

Viewpoint: Conservation isn't enough. We need technology to blunt the impacts of climate change

Lauren Anderson, Genetic Literacy Project

High Anesthesiologist Volume Tied to Better Outcomes for Complex GI Surgical Cases

Endocrinology Advisor, 2020

Powered by **TREND MD**

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 [Email Newsletters](#)

 [RSS Feeds](#)

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

 [Facebook](#)

 [Twitter](#)

 [LinkedIn](#)

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 [Leave Feedback](#)

 [Contact Us](#)

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2021 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.