

地球动力学★地震学★地磁学

扬子褶皱带侏罗纪砂岩古地磁及其褶皱带弧形弯曲的成因

韩玉林^{1,2}, 许长海³, 周祖翼³, 谈晓冬¹

1 中国科学院边缘海地质重点实验室, 中国科学院南海海洋研究所, 广州 510301

2 中国科学院研究生院地球科学学院, 北京 100049

3 同济大学海洋与地球科学学院, 上海 200092

收稿日期 2009-2-3 修回日期 2009-11-26 网络版发布日期 2009-12-20 接受日期

摘要 为了更好地认识上扬子褶皱带和中扬子褶皱带走向差异的机制, 我们对中、上扬子褶皱带过渡的关键地区重庆市万州和云阳两个地区的中、晚侏罗世砂岩进行了古地磁研究. 逐步热退磁分离出两个组分, 低温组分(LTC)在所有样品中均分离出来, 为现代地磁场的重磁化; 中侏罗世样品和万州地区的晚侏罗世样品分离出来的高温组分(HTC)也为现代地磁场的重磁化. 云阳地区晚侏罗世样品分离出来的高温组分通过逐步展平褶皱检验显示: 在褶皱展平至33.8%时, 精度参数达到最大, 相应的古地磁方向为 $D=19.1^\circ$, $I=48.9^\circ$ ($a_{95}=6.3^\circ$), 古地磁极为 73.5°N , 198.2°E ($dp=5.5^\circ$, $dm=8.3^\circ$), 与白垩纪参考古地磁极对比, 此高温组分揭示云阳地区在褶皱变形的后期经历了 $7.7^\circ \pm 6.1^\circ$ 的顺时针旋转. 结合前人的数据, 我们认为中扬子褶皱带普遍存在弯山构造(orocline), 这可能与华北板块向华南板块的挤入作用有关; 但是中、上扬子褶皱带过渡地区的弧形弯曲总体上不是由弯山构造形成的, 很可能是在太平洋板块向北西方向俯冲的宏观板块构造背景下的应变分异作用形成的.

关键词 扬子褶皱带 侏罗纪 古地磁 弯山构造

分类号 P318

DOI: 10.3969/j.issn.0001-5733.2009.12.016

Paleomagnetism of Jurassic sandstones from Yangtze fold belt and its implications for the [JP2] fold belt curvature

HAN Yu-Lin^{1,2}, XU Chang-Hai³, ZHOU Zu-Yi³, TAN Xiao-Dong¹

1 CAS Key Laboratory of Marginal Sea Geology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China

2 College of Earth Sciences, Graduate University of Chinese Academy of Sciences, Beijing 100049, China

3 School of Ocean and Earth Sciences, Tongji University, Shanghai 200092, China

Received 2009-2-3 Revised 2009-11-26 Online 2009-12-20 Accepted

Abstract To promote our knowledge about the origin of the difference in fold axis strikes between upper and middle Yangtze fold belt, we conducted a paleomagnetic study of Middle and Late Jurassic sandstones from Wanzhou and Yunyang County, Chongqing City, a transition zone between the upper and middle Yangtze fold belt. Stepwise thermal demagnetization isolated two components. The low temperature components (LTC) were isolated from all the samples and overprinted by recent geomagnetic field. The high temperature components (HTC) isolated from the Middle and Late Jurassic samples from Wanzhou County were also overprinted by recent geomagnetic field. Stepwise unfolding indicates that the maximum precision parameter of HTC isolated from the Late Jurassic samples from Yunyang County is achieved at 33.8% unfolding. The best-clustered HTC mean direction is $D=19.1^\circ$, $I=48.9^\circ$ ($a_{95}=6.3^\circ$), corresponding to a paleopole at 73.5°N , 198.2°E ($dp=5.5^\circ$, $dm=8.3^\circ$). Compared with the Cretaceous reference paleopole, it reveals that Yunyang County has experienced a clockwise rotation of $7.7^\circ \pm 6.1^\circ$ at the late stage of folding. Combined with published data, our new results support a general orocline in the middle Yangtze fold belt, most probably caused by indentation of NCB into SCB. However, the curvature in the transition zone of upper and middle Yangtze fold belt could be due to strain partitioning in the course of NW directed subduction of the Pacific plate rather than oroclinal bedding.

Key words Yangtze fold belt; Jurassic; Paleomagnetism; Orocline

通讯作者:

谈晓冬 xdtan@scsio.ac.cn

作者个人主页: 韩玉林^{1,2}; 许长海³; 周祖翼³; 谈晓冬¹

扩展功能

本文信息

▶ [Supporting info](#)

▶ [PDF](#) (2887KB)

▶ [\[HTML全文\]](#) (0KB)

▶ [参考文献](#)

服务与反馈

▶ [把本文推荐给朋友](#)

▶ [加入我的书架](#)

▶ [加入引用管理器](#)

▶ [引用本文](#)

▶ [Email Alert](#)

▶ [文章反馈](#)

▶ [浏览反馈信息](#)

相关信息

▶ [本刊中包含“扬子褶皱带”的相关文章](#)

▶ 本文作者相关文章

· [韩玉林](#)

· [许长海](#)

· [周祖翼](#)

· [谈晓冬](#)