



地理学报(英文版) 2003年第13卷第4期

C02 processes in an alpine grassland ecosystem on the Tibetan Plateau

作者: PEI Zhiyong OUYANG Hua

In this paper, the CO₂ concentrations profile from 1.5 m depth in soil to 32 m height in atmosphere were measured from July 2000 to July 2001 in an alpine grassland ecosystem located in the permafrost area on the Tibetan Plateau, which revealed that CO₂ concentrations varied greatly during this study period. Mean concentrations during the whole experiment in the atmosphere were absolutely lower than the CO₂ concentrations in soil, which resulted in CO₂ emissions from the alpine steppe soil to the atmosphere. The highest CO₂ concentration was found at a depth of 1.5 m in soil while the lowest CO₂ concentration occurred in the atmosphere. Mean CO₂ concentrations in soil generally increased with depth. This was the compositive influence of the increasing soil moistures and decreasing soil pH, which induced the increasing biological activities with depth. Temporally, the CO₂ concentrations at different layers in air remained a more steady state because of the atmospheric turbulent mixing. During the seasonal variations, CO₂ concentrations at surface soil interface showed symmetrical patterns, with the lowest accumulation of CO₂ occurring in the late winter and the highest CO₂ concentration in the growing seasons.

CO₂ processes in an alpine grassland ecosystem on the Tibetan Plateau PEI Zhiyong, OUYANG Hua, ZHOU Caiping, XU Xingliang (Inst. of Geographic Sciences and Natural Resources Research, CAS, Beijing 100101, China) 1 Introduction The current concern about global climate change has made it of great interest to find out the real causes of air temperature rising. Observations and further analyses suggest that greenhouse gas increases are responsible for the climate change (Tett et al., 1999; Crowley, 2000). Among all the greenhouse gases in atmosphere, the increasing concentrations of carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) contribute more than 70% of the global warming (Lashof et al., 1990; Rodhe, 1990). Carbon dioxide is the primary gas involved in the exchange for C between the atmosphere and the Earth, and it is responsible for 50% of all greenhouse forcing (Rodhe, 1990). The concentration of atmospheric CO₂ has increased from 280 p.p.m.v. since pre-industrial period (pre-1800) to current about 380 p.p.m.v. which is still increasing at a rate of 3 p.p.m.v. per year (Neftel et al., 1985; Friedli et al., 1986; Fan et al., 1998; Monnin et al., 2001). Beside anthropogenic changes, a large amount of C is returned to the biosphere from the atmosphere by plant photosynthesis and subsequently released from biota to the atmosphere by respiration or burning of plants, so the release of CO₂ from terrestrial biota has contributed significantly to the present atmospheric CO₂ concentration (Sommerfeld, 1993). Carbon exchange between the terrestrial biosphere and the atmosphere is one of the key processes that need to be assessed in the context of the Tokyo Protocol (IGPB Terrestrial Carbon Working Group, 1998). Overview of the current state of the knowledge of global and regional patterns of C exchange by terrestrial ecosystems showed that much of the exchange is affected by human activities (including changing land use and more subtle management effects, such as reduced fire frequency leading to woody encroachment) and is coupled to other climatological and biogeochemical processes (Caspersen et al., 2000; Falkowski et al., 2000; Schimel et al., 2001). The net C exchange of terrestrial ecosystems is the result of a delicate balance between uptake (photosynthesis) and loss (respiration), and shows strong diurnal, seasonal and annual variabilities (Valentini et al., 2000). Soils store two or three times more C than exists in the atmosphere as CO₂, and it is thought that the temperature sensitivity of decomposing organic matter in soil partly determines how much C will be transferred to the atmosphere as a result of global warming (Davidson et al., 2000). Environmental factors such as soil moistures and temperatures influence soil biological activity and CO₂ diffusion, and therefore they have pronounced influences on the seasonal dynamics of C exchange (Keeling et al., 1995; Davidson et al., 1998; Tufekcioglu et al., 2001). According to the previous studies, some other fact

ors such as nitrogen deposition, the availability of soil organic matter, soil pH, and density of plant roots, providing the substrates for soil biological activity, may also control the overall magnitudes of the soil-atmospheric CO₂ exchange (Kelting et al., 1998; Nadelhoffer et al., 1999; Tufekcioglu et al., 1999; Rosenfeld, 2001). The above studies were all limited to the C sequences in soil, plant or atmosphere respectively. Few researches have been done about the C processes within the soil-biosphere-atmosphere profile in one unique site point. It is of great importance to improve the general understanding of C exchange in an integrated ecosystem throughout soil, biosphere and atmosphere. As "the third pole" of the earth, the Tibetan Plateau is one and the only active continental collision area in the world, with a mean altitude of the plateau is more than 4000 m above the sea level and an area of about 2,500,000 km². Great uplift of the plateau since Late Cenozoic has been strongly affecting the physical environment of the plateau itself and its neighboring regions. Meanwhile, the plateau is also a sensitive monitor of climate change in the Asian monsoon region, which is closely related to the global change (Zheng et al., 2000). Due to the topographic features and the characteristics of the atmospheric circulation, typical alpine zones of forest, meadow, grassland and desert appear in succession from southeast to northwest in the plateau (Zheng et al., 1979). Alpine grassland is one of the most important ecosystems on the Tibetan Plateau, which occupied almost 1/3 of the whole plateau area. Besides, the area is special for its lacking of human activities, so this is an ideal place to examine the C exchange within the whole soil-biosphere-atmosphere system. The aim of the present study was (1) to measure CO₂ concentrations within the layers near ground to determine the C exchange in the alpine grassland ecosystem on the Tibetan Plateau; and (2) to analyse the relationships between C exchange and environmental factors.

2 Experimental design

We hypothesized that C exchange would happen between different layers near ground, and the exchange should be controlled by the environmental factors, such as temperatures, moistures and soil pH and so on. To check up the above hypotheses, we carried out the CO₂ concentration profile measurements in a typical alpine grassland ecosystem on the Tibetan Plateau.

2.1 Sampling site

2.2 Materials and methods

2.3 Sample analyses

3 Results

3.1 Soil and vegetation characteristics

The alpine steppe soil at our study site had a lighter texture (sandy loam), and the other physical and chemical characteristics of surface soil were showed in Table 1. Higher C storage was found in 20-30 cm depth of this sandy loam soil. Soil moisture increased with depth, while soil pH decreased gradually with depth. Deeper layer's soil temperature ranged from about -10 to 11 °C, and the variation of upper layer's temperature varied more widely (Figure 2). Among these three layers, the lowest temperature was found in January at a depth of 50 cm and the highest in July at the same depth. The variation of deeper layer temperatures had a distinct time lag following the temperature variations of the upper horizons. These deeper soil temperatures indicated that soil remained frozen throughout the year at a depth of 1.5 m. The vegetation biomass of the study site was slightly lower than other grasslands in the plain areas (Table 2). The ratio of biomass between below-ground and above-ground was about 16:1, which was higher than those of plain areas (Chen et al., 2000). The vegetation root system here was likely stronger than the plain areas due to the frigid climate.

3.2 CO₂ concentration profile

CO₂ concentrations showed great variations within the soil-atmosphere profile during the study period. The lowest CO₂ concentration occurred in the atmosphere. In general, the mean concentrations in the atmosphere were all much lower than the CO₂ concentrations in soil. The largest accumulation of CO₂ was found at a depth of 1.5 m in soil. Variation of soil CO₂ concentrations showed a significant pattern with depth. Soil CO₂ concentrations in our study increased with depth, which is similar to the studies at the Canadian Agriculture Research Station, Delhi, Ontario (Burton et al., 1994). Furthermore, the standard deviations of CO₂ concentrations in atmosphere were all much lower than those in soil, so the CO₂ concentrations in air showed better stabilities than CO₂ concentrations below-ground (Figure 3).

3.3 Temporal variations of CO₂ concentrations

Soil-atmospheric CO₂ concentration profiles varied significantly during the study period, both temporally and with depth. Seasonal variation of CO₂ concentrations at upper layers in soil showed very distinct patterns (Figure 4). Soil CO₂ concentrations at upper layers decreased gradually in the fall, and the lowest accumulation of CO₂ occurred in the late winter. Soil CO₂ concentrations increased to the highest point in the growing season. This phenomenon was similar with the results at the Canadian Agriculture Research Station, Delhi, Ontario (Burton et al., 1994). However, the accumulation of CO₂ concentrations at 1.5 m depth did not exhibit the same trend, the high accumulations of CO₂ were found during the non-growing season. CO₂ concentrations at different heights in air showed more steady-states, but the variations at all horizons did not show any distinct trends during the study period.

3.4 CO₂ fluxes between soil and atmosphere

The mean CO₂ concentrations in the atmosphere were all much lower than the CO₂ concentrations in the soil, which would introduce CO₂ emissions from the alpine steppe soil to the atmosphere in our study area. Measured fluxes of CO₂ in site from the soil to the atmosphere was about 0.17 mol·m⁻²·s⁻¹. The CO₂ contribution from soil to the atmospheric CO₂ was lower than other grasslands in plain areas (Dugas et al., 1997; Dong et al., 2000; Mielnick et al., 2001). The CO₂ emissions s

showed a very distinct trend which decreased in the autumn and increased in the spring (Figure 5). CO₂ emissions during the growing seasons were much higher than those during the non-growing seasons, which accounted for almost 90% of the whole year emissions. The highest mean CO₂ emission occurred in August.

4 Discussion

The net carbon exchange of terrestrial ecosystems is the result of a delicate balance between uptake (photosynthesis) and loss (respiration), and shows a strong diurnal, seasonal and annual variability (Valentini et al., 2000). Soils store two or three times more carbon than exists in the atmosphere as CO₂, and it is thought that the temperature sensitivity of decomposing organic matter in soil partly determines how much carbon will be transferred to the atmosphere as a result of global warming (Davidson et al., 2000). Generally, concentrations of CO₂ in the soil atmosphere are controlled primarily by the rate of soil respiration (including root respiration and microbial respiration), which is also modified by the rate of CO₂ production by biota within soils (Raich et al., 1992; Kelting et al., 1998).

4.1 Variations between CO₂ concentrations and fluxes

Although mean CO₂ concentrations varied greatly with depth in soil, the seasonal variations of CO₂ concentrations showed significant correlations ($R^2 > 0.85$, $p < 0.001$) between the upper layers in the soil. The variation of soil CO₂ concentrations at 1.5 m depth exhibited a different seasonal pattern with those of the other horizons, and the correlation between 1 m and 1.5 m was relatively weak. A significant correlation ($R^2 = 0.63$, $p < 0.001$) was also found between CO₂ at soil surface fluxes and soil CO₂ concentrations at -0.2 m depth during the study period (Figure 6). This implies the variation of CO₂ concentrations in the surface horizon was the most direct driving force of CO₂ emissions from the soil to the atmosphere. The variations of atmospheric CO₂ concentrations at 4 m, 8 m, 16 m and 32 m heights above ground did not show any distinct relationships ($R^2 < 0.4$) between each other. CO₂ transferred quickly and unpredictably between different layers in the air due to the atmospheric turbulence. The atmospheric CO₂ concentrations varied randomly during the experimental period, and their variations between layers hardly have any significant correlations in the atmosphere. Compared with the turbulent concentrations in the atmosphere, soil CO₂ diffusions from deeper layers to upper layers were generally slow and durative.

4.2 Relationships between temperature and CO₂ variations

Soil temperature has often been described as the dominant independent variable determining the carbon uptake and ecosystem respiration rate (Braswell et al., 1997; Tian et al., 1998; Valentini et al., 2000). In the present study, soil CO₂ concentrations at the upper horizons had a seasonal pattern that closely resembled that of the soil temperature (Figures 2 and 4). The temporal variations of CO₂ concentrations at upper layers can be well explained by the seasonal pattern of soil temperature in our study. Significant correlations between CO₂ concentrations and soil temperatures were also found at depths of 0.5 m and 1.0 m ($R^2 = 0.70$, $p < 0.001$) (Figure 7). The annual soil respiration rates were closely correlated with soil temperatures, which agrees well with results of several previous studies that warmer temperatures enhance CO₂ production in different types of soil (Rochette et al., 1999; Waddington et al., 2001). In our study, soil CO₂ efflux showed an asymmetric pattern during the whole year experiment, which was almost the same as the pattern in northern California forest (Xu et al., 2001). The maximum CO₂ emission occurred in summer, and CO₂ emission decreased to the minimum in the winter. Regression analyses showed that the variation in CO₂ fluxes was positively related to air temperature (Figure 8). Root respiration and microbial respiration were the main source of soil CO₂. According to Zogg's study (1996), root respiration rates increased sharply with soil temperature in northern hardwood forests, and the similar relationship between root respiration rates and soil temperatures was also found in many kinds of ecosystems (Bowden et al., 1993; Kelting et al., 1998; Tufekcioglu et al., 2001; Burton et al., 2002). As another important component of soil respiration, microbial decomposition and respiration rates were also strongly affected by soil temperature (Whalen et al., 1990; Updegraff et al., 1998). Microbial populations may adapt better to changing conditions and warmer temperatures. This is supported by studies that have shown that microbial efficiency is affected by the response of the micro-organisms to temperature and that different microbial communities dominate CO₂ production as temperature increases (Lekkerkerk et al., 1990; Zak et al., 1999; Waddington et al., 2001). So soil temperature was the most important factor driving seasonal variation in soil respiration, and this temperature sensitivity of soil respiration has a profound effect on seasonal variation in soil CO₂ concentrations and soil surface CO₂ fluxes.

4.3 Influence of soil moisture

It is well known that soil moisture plays a major role in regulating soil atmosphere CO₂ concentrations through the influence on rates of biological activity and gas diffusion (Parfitt et al., 1997). As documented by Tufekcioglu et al. (2001), a positive linear relationship between soil moistures and soil respiration rates was found among sites. It means that CO₂ production rates generally increased with soil moisture. Orchard et al. (1992) attributed this relationship to the effects of water content on soil microbial communities. While soil temperature was the most important factor driving seasonal variation in soil CO₂ concentrations, it was not significant in terms of explaining variations in depths. Soil CO₂ concentrations increased with depth in our study site, while the soil moistures also increased with depth due to the high surface evaporation in the study site.

(Gao et al., 1995). According to the previous studies, available moisture exerts a great influence on soil microbial activity, and low soil moisture probably reduces microbial populations (Tester, 1988; Ellert et al., 1999; Xu et al., 2001). Furthermore, gas transfer coefficient was determined by soil physical properties, while the increasing soil moisture represented as a diffusion barrier of soil gas (Ellert et al., 1999). So the concentration gradient of soil atmosphere CO₂ could be explained by the increasing of soil moisture with depth in our study. During the seasonal pattern, variation of CO₂ concentration at 1.5 m depth showed a contrary trend to the other horizons (Figure 4). Tufekcioglu et al. (1999) has studied the root biomass distribution in a private farm, and no roots were observed below 1.25 m depth in dry grassland ecosystems. So the microbial respiration occupied all the soil respiration at 1.5 m depth. Why should soil respiration be higher during the cold winter? According to Grace's explanation, perhaps it was because the soil at 1.5 m depth was wetter for longer, and so microbes that were adapted to work at low temperatures were active for most of the year (Grace et al., 2000). Acknowledgements We would like to thank Liu Yunfen, Ma Zhixue and Shi Buhong for their assistance in field sampling, and we are grateful to Qi Yuchun for the help in laboratory. Thanks also should be given to Luo Tianxiang and Niu Haishan for their suggestions during the data analyses.

关键词: CO₂; soil; atmosphere; alpine grassland; Tibetan Plateau