




News



News

[News From the Field](#)

[For the News Media](#)

[Special Reports](#)

[Research Overviews](#)

[NSF-Wide Investments](#)

[Speeches & Lectures](#)

[NSF Current Newsletter](#)

[Multimedia Gallery](#)

[News Archive](#)

News by Research Area

[Arctic & Antarctic](#)

[Astronomy & Space](#)

[Biology](#)

[Chemistry & Materials](#)

[Computing](#)

[Earth & Environment](#)

[Education](#)

[Engineering](#)

[Mathematics](#)

[Nanoscience](#)

[People & Society](#)

[Physics](#)

Press Release 09-189

Ancient China's Sand Dunes Reveal Unexpected Dryness During Heavy Monsoon Rains



Dry sand dunes and monsoon rains in ancient China are out of sync, scientists have found.

[Credit and Larger Version](#)

October 6, 2009

The windswept deserts of northern China might seem an odd destination for studying the heavy monsoon rains that routinely drench the more tropical regions of Southeast Asia. But the sandy dunefields that mark the desert margin between greener pastures to the south and the Gobi Desert to the north are a rich source of information about past climates in Asia, says University of Wisconsin-Madison geographer Joseph Mason.

Wetter periods allow vegetation to take root on and stabilize sand dunes. During dry spells, plants die off and the dunes are more active, constantly shifting as sand is blown away and replenished.

Such patterns of dune activity provide a history of the area's climate--if one can read them, Mason says. "When did those periods of stability or activity occur and from that, what can we infer about climate change?"

As reported in a paper in the October issue of the journal *Geology*, Mason and colleagues mapped sand dune activity across northern China and found unexpectedly high levels of mobility and change 8,000 to 11,500 years ago, a time period generally thought to have a wetter climate.

"The result challenges existing ideas about the monsoon's regional influence, and could impact future climate predictions," says David Verardo, program director in the National Science Foundation (NSF)'s Division of Atmospheric and Geospace Sciences, which funded the research.

Today, the dunes are at the edge of the monsoon region and the scientists expected to find close correlation between precipitation in the dunefields and the strength of the monsoon.

What they found instead was rather surprising. "They turn out to be almost completely out of phase," Mason says. "Where we find lots of active dunes turns out to be a time

when the monsoon system is supposed to have been stronger in southern and central China."

Part of the explanation may lie in local patterns of atmospheric circulation. At the peak of the summer monsoon, central China experiences both heavy summer rainfall and strong upward airflow.

That upward flow tends to be balanced out by more downward air motion--which suppresses precipitation--in areas north and west of the monsoon core.

Regional climate modeling data from the University of Wisconsin-Madison Center for Climatic Research, led by co-author and UW-Madison geoscientist Zhengyu Liu, shows that this pattern may have been strengthened between 8,000 and 11,500 years ago.

The models also show high summer temperatures at that time, which would have increased evaporation and further reduced the moisture that supports dune-stabilizing plants.

This pattern of climate change had been described for areas distant from the monsoon, like Central Asia around the Caspian and Aral Seas and in northern Mongolia.

However, Mason says, "it hasn't really been recognized that this effect could be going on in northern China, which is where our study sites are. What it means is there's much more of a contrast in climate change across a fairly short distance."

The findings relied on a technique called optically stimulated luminescence (OSL), which dates the last time the sand was exposed to sunlight. Radiocarbon dating methods are of limited use since sand typically contains little or no organic material. The OSL method identifies time periods when the sand was actively moving around, indicating little precipitation, and times when dunes were stable.

Mason's previous work in the area suggests that moisture and precipitation are the most significant factors in determining the activity of the Chinese dunes.

The results mean that common assumptions about the effects of future climate changes--including the increased monsoon rainfall predicted by many climatologists--may be incorrect.

"If monsoon rainfall increases in southern China over the next century, the logical assumption would be that these dunes would become more stable as more precipitation also reaches the dune fields and increases vegetation cover," Mason says. "That may not be true. The dunes can become active and the climate there can become drier even when the monsoon is getting stronger."

Even if future rainfall in northern China isn't reduced by changing air circulation patterns as it was in the past, rising temperatures will undoubtedly increase evaporation, he says, exacerbating the water shortages that already plague the area.

An accompanying increase in sand dune activity would reduce available grazing land and worsen air quality.

"If it's drier you have less vegetation and the dunes are active," says Mason. "There will almost certainly be more dust produced, which is a major environmental hazard. Some of the dust from northern China actually reaches Korea, Japan, and even the western U.S."

The paper is co-authored by scientists in China, the Illinois State Geological Survey, and the University of Nebraska-Lincoln. Funding also was provided by the National Natural Science Foundation of China, the China National S&T Basic Work Program, and Nanjing University.

-NSF-

Media Contacts

Cheryl Dybas, NSF (703) 292-7734 cdybas@nsf.gov

Jill Sakai, University of Wisconsin (608) 262-9772 jasakai@wisc.edu

The National Science Foundation (NSF) is an independent federal agency that supports fundamental research and education across all fields of science and engineering. In fiscal year (FY) 2009, its budget is \$9.5 billion, which includes \$3.0 billion provided through the American Recovery and Reinvestment Act. NSF funds reach all 50 states through grants to over 1,900 universities and institutions. Each year, NSF receives about 44,400 competitive requests for funding, and makes over 11,500 new funding awards. NSF also awards over \$400 million in professional and service contracts yearly.

 [Get News Updates by Email](#)

Useful NSF Web Sites:


NSF Home Page: <http://www.nsf.gov>

NSF News: <http://www.nsf.gov/news/>

For the News Media: <http://www.nsf.gov/news/newsroom.jsp>

Science and Engineering Statistics: <http://www.nsf.gov/statistics/>

Awards Searches: <http://www.nsf.gov/awardsearch/>

 [Print this page](#)

[↑ Top](#)

[Web Policies and Important Links](#)

[Privacy](#)

[FOIA](#)

[Help](#)

[Contact NSF](#)

[Contact Webmaster](#)

[SiteMap](#)



The National Science Foundation, 4201 Wilson Boulevard, Arlington, Virginia 22230, USA
Tel: (703) 292-5111, FIRS: (800) 877-8339 | TDD: (800) 281-8749

Last Updated:
October 6, 2009
[Text Only](#)

Last Updated: October 6, 2009