

[ENVIRONMENT](#) | [NEWS RELEASES](#) | [RESEARCH](#) | [SCIENCE](#)


October 1, 2015

Simulating path of 'magma mush' inside an active volcano

Hannah Hickey

UW News

Months of warning signs from Mauna Loa, on Hawaii's Big Island, prompted the U.S. Geological Survey to recently start releasing weekly [updates](#) on activity at the world's largest active volcano. For now, such warning signs can only rely on external clues, like earthquakes and gas emissions. But a University of Washington simulation has managed to demonstrate what's happening deep inside the volcano. The study, published Sept. 7 in [Nature Geoscience](#), is the first to simulate the individual crystals' movement in the magma chamber to better understand the motion of the magma and buildup of pressure.



Doctoral student Jillian Schleicher and professor George Bergantz study volcanoes using computer models and rock samples, like this sample from Mauna Loa. *Dennis Wise/University of Washington*

"The thing about studying volcanoes is we can't really see inside of them to know what's going on," said co-author [Jillian Schleicher](#), a UW doctoral student in Earth and space sciences. "Whenever there's unrest, like earthquakes, gas emissions or surface deformation, it's really difficult to know what processes are taking place inside the volcano."

Each volcano has a unique personality. Volcanologists use the remains of past eruptions and previous warning signs to predict when it might blow. But those predictions are based on only vague understanding of the system's inner workings.

The idealized UW computer simulation could help volcanologists better understand how energy builds up inside a system like Mauna Loa, which is a focus of the UW group's research, to predict when it will erupt.

"This tool is novel because it lets us explore the mechanics," said first author [George Bergantz](#), a UW professor of Earth and space sciences. "It creates an interpretive framework for what controls the movement, and what might produce the signals we see on the outside."


The team used a computer model originally developed by the U.S. Department of Energy to model fuel combustion. The UW group previously adapted the code to simulate volcanic eruptions and ash plumes; this paper is the first time it's been used to go down inside the volcano and examine the movement of each individual crystal.

A volcano is filled with "magma mush," a slushy material that is part magma, or liquid rock, and part solid crystal. Previous studies approximated it as a thick fluid. But capturing its true dual nature makes a difference, since the crystals interact in ways that matter for its motion.

"If we see earthquakes deep inside Mauna Loa, that tells us that there's magma moving up through the volcano," Bergantz said. "But how can we better understand its progress up through that plumbing system?"

The simulation shows the magma has three circulation states: slow, medium and fast. In the slow state, new magma just percolates up through the crystal pores. As the rate of injected magma increases, however, it creates a "mixing bowl" region where older crystals get mixed in with the new material.

"In these crystal-rich mushes, we know that we have magma going in and sometimes it might punch through [the layer of crystals at the bottom]," Schleicher said. "But we don't know how the mixing is happening or the timescales involved."



New magma injected from below (red) combines with older magma (blue) and solid crystals (black and white) in a "mixing bowl" region discovered by the simulation. *George Bergantz/University of Washington*

The current model is an idealized magma chamber, but with more computing power it could be expanded to reproduce a particular volcano's internal structure.

Now that researchers can simulate what happens inside a magma chamber, Schleicher will look at rock samples from Mauna Loa and analyze the layers in the crystals. Crystals preserve chemical clues as they grow, similar to tree rings. Matching the model with the crystals' composition will help recreate the rock's history and track how magma has moved inside Mauna Loa.

"Mauna Loa is a terrific place to study because it's very active, and the rocks contain a single type of crystal," Bergantz said. "What we learn at Mauna Loa will allow us to make headway on other places, like Mount St. Helens, that are intrinsically more difficult."

The research was funded by the National Science Foundation. The other co-author is [Alain Burgisser](#) at the University of Savoy Mont Blanc in France.

###

For more information, contact Bergantz at 206-685-4972 or bergantz@uw.edu and Schleicher at jmschl@uw.edu. Click [here](#) for more photos.

NSF grants: EAR-1049884, EAR01447266, DGE-1256082, TG-EAR140013

[Facebook](#)

[Twitter](#)

[Email](#)

Tag(s): [College of the Environment](#) • [Department of Earth and Space Sciences](#) • [George Bergantz](#) • [volcanoes](#)

SEARCH UW NEWS

Search for:



CATEGORIES

[Browse](#)

UW EXPERTS

[Hurricanes and weather](#)

[Flooding and landslides](#)

[Wildfires](#)

[Public policy and politics](#)

[More topics](#)

Latest news



Mobile health has power to transform HIV/AIDS nursing

2 days ago



Study brings new climate models of small star TRAPPIST 1' s seven intriguing worlds

2 days ago



UW political scientist Mark Smith asks: How do we know what' s true?

3 days ago

More

NEWSLETTER SIGNUP

UW Today Daily

UW Today Week in Review