



A size-structured food-web model for the global ocean

B. A. Ward, S. Dutkiewicz, O. Jahn and M. J. Follows

Limnol. Oceanogr., 57(6), 2012, 1877-1891 | DOI: 10.4319/lo.2012.57.6.1877

ABSTRACT: We present a model of diverse phytoplankton and zooplankton populations embedded in a global ocean circulation model. Physiological and ecological traits of the organisms are constrained by relationships with cell size. The model qualitatively reproduces global distributions of nutrients, biomass, and primary productivity, and captures the power-law relationship between cell size and numerical density, which has realistic slopes of between -1.3 and -0.8 . We use the model to explore the global structure of marine ecosystems, highlighting the importance of both nutrient and grazer controls. The model suggests that zooplankton : phytoplankton ($Z : P$) biomass ratios may vary from an order of 0.1 in the oligotrophic gyres to an order of 10 in upwelling and high-latitude regions. Global estimates of the strength of bottom-up and top-down controls within plankton size classes suggest that these large-scale gradients in $Z : P$ ratios are driven by a shift from strong bottom-up, nutrient limitation in the oligotrophic gyres to the dominance of top-down, grazing controls in more productive regions.

Article Links

[Download Full-text PDF](#)

[Web Appendix](#)

[Return to Table of Contents](#)

Please Note

Articles in L&O appear in PDF format. Open access articles may be freely downloaded by anyone. Other articles are available for download to subscribers only, or may be purchased for \$10 per article. All L&O articles are moved into Open Access after three years.