

Mitochondrial ATP-Sensitive Potassium Channel Activation Protects Cerebellar Granule Neurons From Apoptosis Induced by Oxidative Stress

Yasushi Teshima, MD, PhD; Masaharu Akao, MD, PhD; Ronald A. Li, PhD; Tae H. Chong, BA; William A. Baumgartner, MD; Michael V. Johnston, MD; Eduardo Marbán, MD, PhD

Background and Purpose—Mitochondrial ATP-sensitive potassium (mitoK_{ATP}) channels are present in the brain, and several reports have shown that mitoK_{ATP} channel openers protect the brain against ischemic injury. However, the precise mechanisms of this protection are not well established. We hypothesized that mitoK_{ATP} channel openers prevent apoptosis by preserving mitochondrial membrane potential.

Methods—We investigated the effect of mitoK_{ATP} channel openers on apoptosis induced by oxidative stress using cultured cerebellar granule neurons.

Results—The mitoK_{ATP} channel opener diazoxide (100 μmol/L) significantly suppressed the number of cells with terminal deoxynucleotidyl transferase–mediated dUTP nick end-labeling (TUNEL)–positive nuclei and the increase in caspase-3 activity induced by 20 μmol/L H₂O₂. Diazoxide and another opener, pinacidil, prevented the loss of mitochondrial inner membrane potential (ΔΨ_m) induced by H₂O₂. These effects were abolished by 5-hydroxydecanoate (500 μmol/L), a mitoK_{ATP} channel blocker. Cyclosporin A and bongkreic acid, inhibitors of the mitochondrial permeability transition pore, also prevented ΔΨ_m loss, confirming the involvement of the mitochondrial permeability transition in the apoptotic cascade in neurons. Furthermore, diazoxide prevented the increase in extracellular glutamate concentration induced by H₂O₂, but this effect was not attributable to activation of surface K_{ATP} channels.

Conclusions—MitoK_{ATP} channel openers inhibited apoptosis by preserving mitochondrial inner membrane potential. These beneficial effects may suggest a possible new target for neuroprotection. (*Stroke*. 2003;34:1796-1802.)

Key Words: apoptosis ■ mitochondria ■ neurons ■ oxidative stress ■ potassium channels

Opening of mitochondrial ATP-sensitive potassium (mitoK_{ATP}) channels is protective against cardiac ischemia.^{1,2} The mitoK_{ATP} channel opener diazoxide inhibits activation of the mitochondrial apoptotic pathway induced by hydrogen peroxide (H₂O₂) in cardiac myocytes, and this protective effect is abolished by the mitoK_{ATP} channel blocker 5-hydroxydecanoate (5HD).³ These facts suggest that the prevention of apoptosis may be an important mechanism of mitoK_{ATP}-related cardioprotection. Despite these observations in the heart, less is known about the role of mitoK_{ATP} channels in other organs.

Apoptosis contributes to ischemic injury in the brain,⁴ and brain mitochondria contain 7 times more mitoK_{ATP} channels than heart mitochondria.⁵ Several reports have shown that mitoK_{ATP} channel openers protect the brain against ischemic injury in vivo: diazoxide preserved neuronal-vascular response after cerebral ischemia/reperfusion in piglets,⁶ mitigated canine brain injury induced by hypothermic circulatory arrest,⁷ and decreased cortical infarct size in a murine stroke

model.⁸ However, the mechanisms of these beneficial effects are still obscure. We hypothesized that mitoK_{ATP} channels may inhibit apoptosis by preserving mitochondrial membrane potential in neurons, as they do in cardiomyocytes.

In the present study we used H₂O₂ to simulate neuronal apoptosis induced by oxidative stress⁹ in cultured cerebellar granule neurons, cells that depend on high K⁺ for survival.¹⁰

Materials and Methods

All procedures were performed in accordance with the Johns Hopkins University animal care guidelines, which conform to the *Guide for the Care and Use of Laboratory Animals*, published by the National Institutes of Health.

Primary Culture of Cerebellar Granule Neurons

Cerebellar granule neurons were prepared from 7- to 8-day-old Sprague-Dawley rats as previously described.¹¹ Cells were kept for at least 7 days in culture to differentiate.

Experimental Protocol

Cultured cerebellar granule neurons were randomly assigned to 1 of 4 groups, as follows: (1) control; (2) incubation with H₂O₂ (30

Received July 24, 2002; final revision received January 16, 2003; accepted February 11, 2003.

From the Institute of Molecular Cardiobiology (Y.T., M.A., R.A.L., E.M.), Kennedy Krieger Institute (T.H.C., M.V.J.), and Department of Surgery (W.A.B.), The Johns Hopkins University, Baltimore, Md.

Correspondence to Eduardo Marbán, MD, PhD, Institute of Molecular Cardiobiology, The Johns Hopkins University, 720 Rutland Ave, 844 Ross Bldg, Baltimore, MD 21205. E-mail marban@jhmi.edu

© 2003 American Heart Association, Inc.

Stroke is available at <http://www.strokeaha.org>

DOI: 10.1161/01.STR.0000077017.60947.AE




Figure 1. Effects of diazoxide and 5HD on cell injury induced by H₂O₂. A, Phase-contrast microscope images. B, Representative images of TUNEL staining. C, Quantitative percentages of the cells with TUNEL-positive nuclei. D, Effects of diazoxide and 5HD on caspase-3 activity at 4, 8, and 16 hours after H₂O₂ application. Data of each time point are relative percentage of caspase-3 activity to control at the same time point. CON indicates control; H₂O₂, 20 μ mol/L H₂O₂; DZ, H₂O₂+100 μ mol/L diazoxide; 5HD, H₂O₂+diazoxide+500 μ mol/L 5HD; and Caspase inhibitor, H₂O₂+50 μ mol/L z-VAD-fmk. Data are mean \pm SEM (n=4 per group in TUNEL staining; n=3 per group in caspase-3 activity assay). * P <0.001 vs CON, ** P <0.001 vs H₂O₂, # P <0.05 vs CON, DZ at the same time point.

μ mol/L for time-lapse confocal microscopy, and 20 μ mol/L for the other experiments) (3) 100 μ mol/L diazoxide with H₂O₂; and (4) 100 μ mol/L diazoxide and 500 μ mol/L 5HD with H₂O₂. In the appropriate groups, diazoxide and 5HD were applied 20 minutes before H₂O₂ application. In the experiments with mitochondrial permeability transition pore (PTP) inhibition, 0.2 or 1 μ mol/L cyclosporin A (CsA) or 50 μ mol/L bongkreikic acid (BA) was used instead of diazoxide. Appropriate H₂O₂ concentrations and the length of stimulation were used in each experiment to induce the similar effects on the cells in different conditions.

Terminal Deoxynucleotidyl Transferase-Mediated dUTP Nick End-Labeling Staining

Terminal deoxynucleotidyl transferase-mediated dUTP nick end-labeling (TUNEL) staining was performed 16 hours after H₂O₂ exposure (or at the equivalent time in controls) according to the manufacturer's protocol (Roche), as previously described.³

Caspase-3 Activity Assay

Caspase-3 activity was measured by detection of the cleavage of a colorimetric caspase-3 substrate, *N*-acetyl-Asp-Glu-Val-Asp-p-nitroaniline (DEVD-pNA), with the use of an assay kit, ApoAlert CPP32 (Clontech). The assay was performed according to the manufacturer's protocol at 4, 8, and 16 hours of stimulation.

Flow Cytometry Analysis of Mitochondrial Membrane Potential

Loss of mitochondrial membrane potential ($\Delta\Psi_m$) was assessed by a fluorescent indicator, tetramethylrhodamine ethyl ester (TMRE) (Molecular Probes) (100 nmol/L), with the use of flow cytometry. Cells were treated with drugs for 120 minutes at 37°C and analyzed by FACScan (20 000 cells per sample) (Becton Dickinson). Data were analyzed with the use of WinMDI analysis software (<http://facs.scripps.edu>).

Time-Lapse Analysis of $\Delta\Psi_m$ Loss

Cells were placed in Hank's salt-based MEM with 25 mmol/L HEPES buffer to avoid pH change in non-CO₂ equilibrated environment.

After 30-minute incubation in medium containing 100 nmol/L TMRE at 37°C, time-lapse scanning was started with a 20 \times objective lens. During the scanning, cells were maintained at 37°C with a heater platform installed on the microscope stage. Thirty cells were randomly and prospectively selected in each scan, and TMRE fluorescence intensity was sequentially monitored every 5 minutes.

Assays for Extracellular Glutamate Concentration

Glutamate concentrations in the culture medium from each experimental group were assayed by high-performance liquid chromatography with electrochemical detection, as described previously.¹²

Electrophysiology

Electrophysiological recordings were performed at room temperature with the whole-cell patch clamp technique. Pipette electrodes had final tip resistances of 5 to 7 mol/L Ω . The internal pipette solution contained the following (mmol/L): potassium glutamate 120, KCl 25, ATP (magnesium salt) 1, EGTA 10, MgCl₂ 0.5, HEPES 10 [pH 7.2]. The external bath solution contained the following (mmol/L): NaCl 140, KCl 5, MgCl₂ 1, CaCl₂ 1, HEPES 10 [pH 7.4]. Designated amounts of drugs were added to the bath when required. Currents were acquired and analyzed with custom-written software.

Statistical Analysis

All quantitative data are presented as mean \pm SEM. Multiple comparisons among groups were performed by 1-way ANOVA with Fisher's least significant difference as the post hoc test. A level of P <0.05 was accepted as statistically significant.

Results

Effects of Diazoxide on Apoptosis in Cerebellar Granule Neurons

Morphology of cultured cells was assessed by transmitted phase-contrast microscopy (Figure 1A). The control group showed round cell bodies with clear edges and fine dendritic

network. Incubation with 20 $\mu\text{mol/L}$ H_2O_2 for 16 hours decreased the number of cells and induced shrinkage of cell bodies and disruption of dendritic networks. Diazoxide mitigated the morphological manifestations of cell damage, and these protective effects were abolished by 5HD. Next we performed TUNEL staining as an indicator of apoptosis. Figure 1B shows representative staining in each group, and Figure 1C shows quantitative results. Control cells showed few TUNEL-positive nuclei ($3.6 \pm 0.84\%$). The number of TUNEL-positive nuclei was increased to $47.7 \pm 4.7\%$ in the cells exposed to H_2O_2 . Diazoxide significantly decreased the number of TUNEL-positive nuclei to $7.7 \pm 2.7\%$. Treatment with 5HD partially abolished these protective effects of diazoxide ($27.5 \pm 2.1\%$).

Effects of Diazoxide on Caspase-3 Activity in Cerebellar Granule Neurons

Since activation of apoptosis-effector caspases such as caspase-3 plays an important role in neuronal cell death after cerebral ischemia,¹³ we next assayed caspase-3 activity. Figure 1D shows relative caspase-3 activity in each group at various times after H_2O_2 application. At 8 and 16 hours, caspase-3 activity was significantly increased in the H_2O_2 group (124% and 161% of control group, respectively), and diazoxide suppressed these increases. 5HD completely abolished the effects of diazoxide.

Effects of Diazoxide on $\Delta\Psi_m$

Dissipation of $\Delta\Psi_m$ is a critical event early in the process of apoptosis.¹⁴ To examine whether preservation of $\Delta\Psi_m$ is associated with the antiapoptotic effect of mitoK_{ATP} channel activation, we assessed the effect of diazoxide on $\Delta\Psi_m$ using flow cytometry. Incubation with 20 $\mu\text{mol/L}$ H_2O_2 for 2 hours decreased TMRE fluorescence and shifted the distribution curve leftward to a level similar to that obtained with the cells treated with mitochondrial uncoupler carbonyl cyanide *p*-(trifluoromethoxy) phenylhydrazone (FCCP), indicating the depolarization of $\Delta\Psi_m$ (Figure 2A). Diazoxide remarkably decreased the fraction of neurons that underwent H_2O_2 -induced dissipation of $\Delta\Psi_m$, as shown by the preservation of cells with normal $\Delta\Psi_m$ (Figure 2A). 5HD abolished the salutary effect of diazoxide, reverting the distribution to that of the H_2O_2 group (Figure 2A). Quantitative summary data from flow cytometry are shown in Figure 2B; here, we evaluated the percentage of cells that exhibited a high level of TMRE fluorescence (defined as $>10^2$ of fluorescence intensity). Diazoxide inhibited the $\Delta\Psi_m$ loss in a concentration-dependent manner, and this effect reached saturation at a concentration of 100 $\mu\text{mol/L}$ (Figure 2B, left). Furthermore, pinacidil, which activates both mitochondrial and sarcolemmal K_{ATP} channels,² also prevented $\Delta\Psi_m$ loss in a 5HD-inhibitable manner (Figure 2B, left). Diazoxide or 5HD alone did not measurably affect TMRE fluorescence (Figure 2B, right).

Time-Lapse Analysis of $\Delta\Psi_m$ Loss

To examine time-dependent changes of $\Delta\Psi_m$ on a single-cell basis, confocal microscopy was performed with the use of cells loaded with TMRE. Figure 3A shows sequential images of TMRE fluorescence in each group. Figure 3C demon-




Figure 2. Effects of diazoxide and 5HD on $\Delta\Psi_m$ in H_2O_2 -stimulated cells. A, Representative histograms in each group obtained from flow cytometry. CON indicates control; H_2O_2 , 20 $\mu\text{mol/L}$ H_2O_2 ; DZ, H_2O_2 +100 $\mu\text{mol/L}$ diazoxide; 5HD, H_2O_2 +diazoxide+500 $\mu\text{mol/L}$ 5HD; and FCCP, 10 $\mu\text{mol/L}$ FCCP. B, Quantitative results of flow cytometry. Data are mean \pm SEM ($n=3$). * $P<0.05$, # $P<0.01$ vs DZ 0 $\mu\text{mol/L}$. Pina indicates pinacidil.

strates time course of TMRE fluorescence in 30 cells randomly and prospectively selected in each group. TMRE fluorescence did not change during 90 minutes of scanning in the control group. In cells treated with H_2O_2 , fluorescence intensity began to decrease within 10 minutes of H_2O_2 application and progressively decreased thereafter. Diazoxide prevented $\Delta\Psi_m$ loss in the majority of cells and delayed the onset of mitochondrial depolarization in those cells that were not fully protected. These effects of diazoxide were cancelled by 5HD. The averaged intensity data from 30 cells are shown in Figure 3B.

Effects of CsA and BA on $\Delta\Psi_m$

PTP is a nonspecific large-conductance channel in the inner mitochondrial membrane. Opening of PTP dissipates $\Delta\Psi_m$ by




Figure 3. Time-lapse analysis of loss of $\Delta\Psi_m$. A, Representative images of sequential changes of TMRE fluorescence in each group. B, Mean fluorescence intensity of each group. Results are mean \pm SEM. C, Time course of TMRE fluorescence of 30 cells randomly and prospectively selected in each group. CON indicates control; H₂O₂, 30 μ mol/L H₂O₂; DZ, H₂O₂+100 μ mol/L diazoxide; and 5HD, H₂O₂+diazoxide+500 μ mol/L 5HD. Similar results were obtained from 3 independent experiments.

increasing inner membrane permeability.¹⁵ We next addressed the question of whether PTP may account for the $\Delta\Psi_m$ dissipation induced by H₂O₂ in cerebellar granule neurons. The effects of CsA, a PTP inhibitor, on $\Delta\Psi_m$ loss by H₂O₂ were investigated with flow cytometry (Figure 4A) as well as time-lapse confocal microscopy (Figure 4B) of cerebellar neurons. Incubation with 20 μ mol/L H₂O₂ for 2 hours resulted in $\Delta\Psi_m$ depolarization to a level similar to that of with FCCP-treated cells, as shown by the decrease of TMRE fluorescence. CsA remarkably prevented H₂O₂-induced dissipation of $\Delta\Psi_m$, as shown by the preservation of cells with intact $\Delta\Psi_m$. Observation of fluorescence intensity in individual cells revealed that CsA inhibited H₂O₂-induced dissipation of $\Delta\Psi_m$ in a manner similar to that of diazoxide, decreasing the number of cells undergoing $\Delta\Psi_m$ depolarization and delaying the onset of $\Delta\Psi_m$ depolarization in unprotected cells. **Importantly, 5HD did not inhibit the effect of CsA.** BA, an inhibitor of adenine nucleotide translocase (a putative component of the PTP), prevented $\Delta\Psi_m$ loss induced by H₂O₂ in a manner similar to that of CsA, and 5HD did not inhibit this effect of BA. **These results indicate the involvement of PTP in $\Delta\Psi_m$ loss induced by H₂O₂ and may suggest that mitoK_{ATP} channels work upstream of PTP.** Figure 4C shows pooled flow cytometry data. We evaluated the percentage of cells that exhibited a high level of TMRE fluorescence

(defined as $>10^2$ of fluorescence intensity). The pooled data confirm the representative results.

Effects of Diazoxide on Extracellular Glutamate Concentration

Given the well-recognized role of glutamate in excitotoxicity and neuronal apoptosis,¹⁶ we assayed extracellular glutamate concentration in each experimental group. H₂O₂ stimulation significantly increased glutamate concentration in the culture medium, and diazoxide suppressed this increase in a 5HD-inhibitable manner (Figure 5A).

Effects of Diazoxide on Surface K_{ATP} Channels

In some cells, K_{ATP} channels exist both in the surface membrane and in the inner mitochondrial membrane. If surface K_{ATP} channels are present in cerebellar granule neurons, they may be activated by diazoxide; the resultant hyperpolarization could underlie the inhibition of glutamate release and the neuroprotective effects. To determine whether the antiapoptotic effects of diazoxide in cerebellar granule neurons are mediated by mitoK_{ATP} channels rather than surface K_{ATP} channels, we measured surface membrane current in cultured cerebellar granule neurons. Figure 5B shows that application of the surface K_{ATP} channel agonist pinacidil² (100 μ mol/L) or its potent derivative P-1075¹⁷ (100 μ mol/L)




Figure 4. Effects of CsA on $\Delta\Psi_m$ in H_2O_2 -stimulated cells. **A**, Representative histograms obtained by flow cytometry. Cells were stimulated by $20 \mu\text{mol/L}$ H_2O_2 . FCCP indicates $10 \mu\text{mol/L}$ FCCP. **B**, Time-dependent alterations of TMRE fluorescence in 30 randomly selected cells. Cells were stimulated by $30 \mu\text{mol/L}$ H_2O_2 . **C**, Quantitative data of flow cytometry. Data are expressed as percentage of cells with high level of TMRE fluorescence. Data are mean \pm SEM ($n=3$ per group). * $P<0.01$ vs CsA $0 \mu\text{mol/L}$.

resulted in modest, statistically insignificant ($P>0.05$) increases in the steady state outward currents. No hint of an increase, however, was observed with diazoxide ($100 \mu\text{mol/L}$). Taken collectively, the results indicate that surface K_{ATP} channels are sparsely distributed in rat cerebellar granule neurons; the few channels that may be present are not sensitive to diazoxide. Thus, the protective effects we have described are not attributable to surface K_{ATP} channel activation.

Discussion

The principal findings of our study are as follows. (1) Diazoxide, a mitoK_{ATP} channel opener, inhibited the apoptotic cascade and alleviated cell damage induced by oxidative stress. (2) Diazoxide and pinacidil prevented $\Delta\Psi_m$ loss induced by oxidative stress in a 5HD-inhibitable manner. (3) CsA or BA, an inhibitor of PTP, prevented $\Delta\Psi_m$ loss induced by oxidative stress, and this effect was not inhibited by 5HD. (4) Diazoxide prevented the increase in the concentration of extracellular glutamate induced by H_2O_2 stimulation. These results indicate that mitoK_{ATP} channel openers protect neurons against oxidative stress by preserving $\Delta\Psi_m$. Inhibition of PTP opening likewise suppresses H_2O_2 -induced apoptosis. Inhibition of glutamate release may be one mechanism contributing to the neuroprotective effects of diazoxide.

Figure 6 shows a schematic diagram of our working hypothesis to explain the neuroprotection by activation of mitoK_{ATP} channels. Opening of PTP is considered to be a critical event in apoptosis^{14,18} and possibly a trigger of cell death in ischemia/reperfusion injury in the brain.¹⁹ When one considers the present results that both CsA and BA prevented $\Delta\Psi_m$ loss induced by H_2O_2 , PTP opening is closely involved in the process of apoptosis in neurons. In fact, several reports showed that CsA protected the brain against ischemia/reperfusion injury in rodents.^{20,21} Although we did not directly prove the prevention of PTP opening by diazoxide, the present observations that diazoxide and CsA inhibited $\Delta\Psi_m$ loss in a similar manner may indicate that these 2 drugs affect the same pathway. Furthermore, the results that 5HD did not cancel the effects of CsA and BA suggest that mitoK_{ATP} channels presumably come into play upstream of PTP. Diazoxide was reported to partially depolarize $\Delta\Psi_m$ in neuronal cells,²² which may attenuate Ca^{2+} accumulation in mitochondria by decreasing the driving force for Ca^{2+} uptake. In fact, diazoxide attenuates the accumulation of mitochondrial Ca^{2+} during simulated ischemia/reperfusion injury in cardiomyocytes.²³ Matrix Ca^{2+} overload, in turn, is known to induce opening of PTP.²⁴ Hence, activation of mitoK_{ATP} channels may prevent PTP opening by reducing matrix Ca^{2+} overload, thereby inhibiting apoptosis. Reactive oxygen spe-




Figure 5. A, Effects of diazoxide on extracellular glutamate concentration. Data are relative percentage of glutamate concentration to control. CON indicates control; H₂O₂, 20 μ mol/L H₂O₂; DZ, H₂O₂+100 μ mol/L diazoxide; and 5HD, H₂O₂+diazoxide+500 μ mol/L 5HD. Data are mean \pm SEM (n=4 per group). * P <0.05 vs CON. B, Distribution of raw (open squares) and averaged (solid circles) steady state outward current densities measured at 300 ms after depolarizing cells to +20 mV from a holding potential of -80 mV preceded by a 100-ms prepulse to 0 mV in the absence and presence of 100 μ mol/L pinacidil (Pina), 100 μ mol/L P-1075, and 100 μ mol/L diazoxide (DZ) as indicated.

cies (ROS) generation, another important trigger of PTP opening, was reportedly decreased by diazoxide in hippocampal cells²⁵ and in isolated heart mitochondria.²⁶ In contrast, diazoxide increased ROS generation to result in cardiopro-




Figure 6. Proposed mechanism of neuroprotection by activation of mitoK_{ATP} channels.

tection.²⁷ These results suggest the possibility that a moderate increase of ROS by diazoxide may prevent a subsequent large increase of ROS production. As other possible mechanisms, a recent report demonstrated that diazoxide suppressed the cytochrome *c* release and translocation of Bax and increased Bcl-2 protein level in cultured hippocampal cells.⁸

Glutamate is the major excitatory neurotransmitter in the central nervous system, and, during the postnatal period, glutamate-activated *N*-methyl-D-aspartate (NMDA) receptors increase Ca²⁺ influx into cytosol to trigger changes in neuronal metabolism and gene expression that are necessary for brain development. In contrast, overstimulation of NMDA receptors by glutamate induces cell death in a process known as excitotoxicity,¹⁶ and mitochondrial Ca²⁺ overload is closely associated with this process.¹⁶ Several studies have shown that extracellular glutamate increases in brain injury,^{28,29} although the precise mechanisms are still controversial. Reversed operation of the glutamate transporter is the main candidate for the increased extracellular glutamate in ischemia,³⁰ and it may be induced by ATP depletion.³¹ In the present study we observed that H₂O₂ stimulation increased extracellular glutamate concentration, and diazoxide inhibited the increase of glutamate. These results suggest the possibility that reduced glutamate release may be one mechanism of neuroprotection by diazoxide, although the precise mechanism by which activation of mitoK_{ATP} channels prevents glutamate release remains unclear. As one possibility, diazoxide may prevent glutamate release by preserving ATP production.³² However, given the protective effects of diazoxide in cardiomyocytes,³ suppression of glutamate release may not be the primary mechanism of protection in neuronal cells. We cannot exclude the possibility that increased extracellular glutamate might simply be due to glutamate leakage from dead cells, such that the decrease in glutamate by diazoxide may be the result of (rather than the cause of) reduced cell injury. Further investigation will be needed to elucidate precisely how the suppression of glutamate release is associated with neuroprotection by mitoK_{ATP} channel openers.

The accumulation of TMRE in mitochondria depends not only on $\Delta\Psi_m$, but also on the cytosolic concentration of the probe. To exclude the possibility that the effects of diazoxide are mediated by the inhibition of multidrug resistance pump in plasma membrane, we examined the effect of MK 571,³³ a specific multidrug resistance pump inhibitor, on $\Delta\Psi_m$ in H₂O₂-stimulated cells. TMRE fluorescence did not change in MK 571-treated cells, indicating that the $\Delta\Psi_m$ -preserving effects of diazoxide or CsA do not depend on multidrug resistance inhibition (data not shown).

Many excitable cells, including some neurons, contain 2 distinct K_{ATP} channels: the classic one in the cell membrane (surface K_{ATP} channel)³⁴ and the other in the mitochondrial inner membrane (mitoK_{ATP} channel).⁵ These 2 channels are known to have important pharmacological differences in the heart. Surface K_{ATP} channels from cardiac sarcolemma are resistant to diazoxide and 5HD, whereas mitoK_{ATP} channels are sensitive to both agents.² There has been much debate as to whether surface K_{ATP} channels or mitoK_{ATP} channels predominantly contribute to cell protection mainly in the

heart, and the subject is still controversial. A newly identified mitoK_{ATP} channel opener, BMS 191095, has cardioprotective effects without activating surface K_{ATP} channels,³⁵ supporting the predominance of mitoK_{ATP} channels in the protective effects. In the present study we confirmed that a neuroprotective concentration of diazoxide did not activate surface K_{ATP} channels in cultured cerebellar granule neurons. These results indicate that the neuroprotective effects of diazoxide are due to activation of mitoK_{ATP} channels, at least in cerebellar granule cells.

When it is considered that apoptosis is a significant contributor to ischemia/reperfusion injury in the brain,^{4,8} mitoK_{ATP} channels could be a possible target for neuroprotection. Further studies are required to explore more detailed mechanisms of the observed effects.

Acknowledgments

This study was supported by the National Institutes of Health (R37 NS31238, R37 HL36957, R01 NS28208, and P50 HD240611) and the Banyu Fellowship in Cardiovascular Medicine (Dr Akao). Dr Marbán holds the Michel Mirowski, MD, Professorship of Cardiology.

References

- Garlid KD, Paucek P, Yarov-Yarovsky V, Murray HN, Darbenzio RB, D'Alonzo AJ, Lodge NJ, Smith MA, Grover GJ. Cardioprotective effect of diazoxide and its interaction with mitochondrial ATP-sensitive K⁺ channels: possible mechanism of cardioprotection. *Circ Res*. 1997;81:1072-1082.
- Liu Y, Sato T, O'Rourke B, Marbán E. Mitochondrial ATP-dependent potassium channels: novel effectors of cardioprotection? *Circulation*. 1998;97:2463-2469.
- Akao M, Ohler A, O'Rourke B, Marbán E. Mitochondrial ATP-sensitive potassium channels inhibit apoptosis induced by oxidative stress in cardiac cells. *Circ Res*. 2001;88:1267-1275.
- Li Y, Sharov VG, Jiang N, Zaloga C, Sabbah HN, Chopp M. Ultrastructural and light microscopic evidence of apoptosis after middle cerebral artery occlusion in the rat. *Am J Pathol*. 1995;146:1045-1051.
- Bajgar R, Seetharaman S, Kowaltowski AJ, Garlid KD, Paucek P. Identification and properties of a novel intracellular (mitochondrial) ATP-sensitive potassium channel in brain. *J Biol Chem*. 2001;276:33369-33374.
- Domoki F, Perciaccante JV, Veltkamp R, Bari F, Busija DW. Mitochondrial potassium channel opener diazoxide preserves neuronal-vascular function after cerebral ischemia in newborn pigs. *Stroke*. 1999;30:2713-2718.
- Shake JG, Peck EA, Marbán E, Gott VL, Johnston MV, Troncoso JC, Redmond JM, Baumgartner WA. Pharmacologically induced preconditioning with diazoxide: a novel approach to brain protection. *Ann Thorac Surg*. 2001;72:1849-1854.
- Liu D, Lu C, Wan R, Auyeung WW, Mattson MP. Activation of mitochondrial ATP-dependent potassium channels protects neurons against ischemia-induced death by a mechanism involving suppression of Bax translocation and cytochrome c release. *J Cereb Blood Flow Metab*. 2002;22:431-443.
- Whittemore ER, Loo DT, Cotman CW. Exposure to hydrogen peroxide induces cell death via apoptosis in cultured rat cortical neurons. *Neuroreport*. 1994;5:1485-1488.
- D'Mello SR, Galli C, Ciotti T, Calissano P. Induction of apoptosis in cerebellar granule neurons by low potassium: inhibition of death by insulin-like growth factor I and cAMP. *Proc Natl Acad Sci U S A*. 1993;90:10989-10993.
- Johns DC, Nuss HB, Marbán E. Suppression of neuronal and cardiac transient outward currents by viral gene transfer of dominant-negative Kv4.2 constructs. *J Biol Chem*. 1997;272:31598-31603.
- Donzanti BA, Yamamoto BK. An improved and rapid HPLC-EC method for the isocratic separation of amino acid neurotransmitters from brain tissue and microdialysis perfusates. *Life Sci*. 1988;43:913-922.
- Namura S, Zhu J, Fink K, Endres M, Srinivasan A, Tomaselli KJ, Yuan J, Moskowitz MA. Activation and cleavage of caspase-3 in apoptosis induced by experimental cerebral ischemia. *J Neurosci*. 1998;18:3659-3668.
- Kroemer G, Dallaporta B, Resche-Rigon M. The mitochondrial death/life regulator in apoptosis and necrosis. *Annu Rev Physiol*. 1998;60:619-642.
- Crompton M. The mitochondrial permeability transition pore and its role in cell death. *Biochem J*. 1999;341:233-249.
- Nicholls DG, Budd SL. Mitochondria and neuronal survival. *Physiol Rev*. 2000;80:315-360.
- Sato T, Sasaki N, O'Rourke B, Marbán E. Adenosine primes the opening of mitochondrial ATP-sensitive potassium channels: a key step in ischemic preconditioning? *Circulation*. 2000;102:800-805.
- Green DR, Reed JC. Mitochondria and apoptosis. *Science*. 1998;281:1309-1312.
- Sims NR, Anderson MF. Mitochondrial contributions to tissue damage in stroke. *Neurochem Int*. 2002;40:511-526.
- Matsumoto S, Friberg H, Ferrand-Drake M, Wieloch T. Blockade of the mitochondrial permeability transition pore diminishes infarct size in the rat after transient middle cerebral artery occlusion. *J Cereb Blood Flow Metab*. 1999;19:736-741.
- Shiga Y, Onodera H, Matsuo Y, Kogure K. Cyclosporin A protects against ischemia-reperfusion injury in the brain. *Brain Res*. 1992;595:145-148.
- Debska G, May R, Kicinska A, Szewczyk A, Elger CE, Kunz WS. Potassium channel openers depolarize hippocampal mitochondria. *Brain Res*. 2001;892:42-50.
- Murata M, Akao M, O'Rourke B, Marbán E. Mitochondrial ATP-sensitive potassium channels attenuate matrix Ca²⁺ overload during simulated ischemia and reperfusion: possible mechanism of cardioprotection. *Circ Res*. 2001;89:891-898.
- Hunter DR, Haworth RA. The Ca²⁺-induced membrane transition in mitochondria, III: transitional Ca²⁺ release. *Arch Biochem Biophys*. 1979;195:468-477.
- Goodman Y, Mattson MP. K⁺ channel openers protect hippocampal neurons against oxidative injury and amyloid beta-peptide toxicity. *Brain Res*. 1996;706:328-332.
- Ozcan C, Bienengraeber M, Dzeja PP, Terzic A. Potassium channel openers protect cardiac mitochondria by attenuating oxidant stress at reoxygenation. *Am J Physiol*. 2002;282:H531-H539.
- Forbes RA, Steenbergen C, Murphy E. Diazoxide-induced cardioprotection requires signaling through a redox-sensitive mechanism. *Circ Res*. 2001;88:802-809.
- Bullock R, Zauner A, Woodward J, Young HF. Massive persistent release of excitatory amino acids following human occlusive stroke. *Stroke*. 1995;26:2187-2189.
- Tseng EE, Brock MV, Kwon CC, Annanata M, Lange MS, Troncoso JC, Johnston MV, Baumgartner WA. Increased intracerebral excitatory amino acids and nitric oxide after hypothermic circulatory arrest. *Ann Thorac Surg*. 1999;67:371-376.
- Rossi DJ, Oshima T, Attwell D. Glutamate release in severe brain ischaemia is mainly by reversed uptake. *Nature*. 2000;403:316-321.
- Madl JE, Burgesser K. Adenosine triphosphate depletion reverses sodium-dependent, neuronal uptake of glutamate in rat hippocampal slices. *J Neurosci*. 1993;13:4429-4444.
- Ozcan C, Holmuhamedov EL, Jahangir A, Terzic A. Diazoxide protects mitochondria from anoxic injury: implications for myopreservation. *J Thorac Cardiovasc Surg*. 2001;121:298-306.
- DeCory HH, Piech-Dumas KM, Sheu SS, Federoff HJ, Anders MW. Efflux of glutathione conjugate of monochlorobimane from striatal and cortical neurons. *Drug Metab Dispos*. 2001;29:1256-1262.
- Ashford ML, Sturgess NC, Trout NJ, Gardner NJ, Hales CN. Adenosine-5'-triphosphate-sensitive ion channels in neonatal rat cultured central neurones. *Pflugers Arch*. 1988;412:297-304.
- Grover GJ, D'Alonzo AJ, Garlid KD, Bajgar R, Lodge NJ, Sleph PG, Darbenzio RB, Hess TA, Smith MA, Paucek P, Atwal KS. Pharmacologic characterization of BMS-191095, a mitochondrial K_{ATP} opener with no peripheral vasodilator or cardiac action potential shortening activity. *J Pharmacol Exp Ther*. 2001;297:1184-1192.