



## 两种品质类型花生品质形成与子叶细胞超微结构的关系

张佳蕾<sup>1</sup>, 李向东<sup>1\*</sup>, 杨传婷<sup>1</sup>, 高芳<sup>2</sup>, 张凤<sup>1</sup>, 王媛媛<sup>1</sup>, 孙莲强<sup>1</sup>

<sup>1</sup>山东农业大学农学院, 山东农业大学作物生物学国家重点实验室, 山东泰安 271018;

<sup>2</sup>山东圣丰种业科技有限公司, 山东嘉祥 272400

Relationship of quality formation and ultrastructure of cotyledon cells in two quality types of peanut

ZHANG Jia-Lei<sup>1</sup>, LI Xiang-Dong<sup>1\*</sup>, YANG Chuan-Ting<sup>1</sup>, GAO Fang<sup>2</sup>, ZHANG Feng<sup>1</sup>, WANG Yuan-Yuan<sup>1</sup>, and SUN Lian-Qiang<sup>1</sup>

<sup>1</sup>College of Agronomy, State Key Laboratory of Crop Biology, Shandong Agricultural University, Tai'an, Shandong 271018, China;

<sup>2</sup>Shofine Seed Co., Ltd., Jiaxiana, Shandong 272400, China

[摘要](#)

[图/表](#)

[参考文献](#)

[相关文章](#)

[点击分布统计](#)

[下载分布统计](#)

?