| EGU.eu | | EGU Journals | Contact |

#### Home

## Online Library eE

#### Online Library eED

- Papers in Open Discussion
- Volumes and Issues
- Special Issues
- Library Search
- Title and Author Search

## General Information



■ Volumes and Issues
■ Contents of Issue 1

eEarth Discuss., 3, 1-9, 2008 www.electronic-earth-discuss.net/3/1/2008/ © Author(s) 2008. This work is licensed under a Creative Commons License.

# Zoogeography of the bottom Foraminifera of the West-African coast

V. Mikhalevich Zoological Institute Russian Academy of Sciences, Russia

Abstract. The sediment samples from the continental shelf of West-Equatorial Africa (from the Strait of Gibraltar to the Niger estuary), depths ranging from 0 to 69 m were found to contain 176 bottom foraminiferal species. For the majority of them (126 species), their areas of occurrences were mapped and the peculiar features of the geographical range and distribution were studied. The species natural habitats were established based on the taxonomical revision of the species in study all over the World Ocean based on the collections of the Zoological Institute RAS and wide literary data. The method of perforated cards was used to mark the geographical locations of all of the species studied. In order to establish the species geographic zonal distribution (together with their depth habitat) the five characteristic groups of the species were separated: 1. pan-oceanic (cosmopolitan), 2. widely spread tropical-boreal, 3. tropicallaw boreal, 4. tropical-subtropical, 5. tropical. The percent of the species of each group among the species composition was established for the fauna of each station and for the whole region.

■ <u>Discussion Paper</u> (PDF, 2400 KB) ■ <u>Interactive Discussion</u> (Final Response, 8 Comments)

Citation: Mikhalevich, V.: Zoogeography of the bottom Foraminifera of the West-African coast, eEarth Discuss., 3, 1-9,

2008. ■ <u>Bibtex</u> ■ <u>EndNote</u> ■ <u>Reference Manager</u>



### Recent Papers

01 | eED, 29 Sep 2009: Thermogeodynamic manifestations in the Caucasus and their genesis

02 | eE, 13 Jul 2009: Holocene evolution and sedimentation rate of Alikes Lagoon, Zakynthos island, Western Greece: preliminary results

03 | eE, 08 Jul 2009: Morphology of the pore space in claystones – evidence from BIB/FIB ion beam sectioning and cryo-SEM observations