

Science News

from research organizations

New DNA study provides critical information on conserving rainforest lizards

Date: August 2, 2021

Source: University of Washington

Summary: Researchers have provided an important road map to conserving rough-nosed horned lizards in Sri Lanka.

Share: [!\[\]\(17413706fd4997a1a4bdf85c6864eee1_img.jpg\)](#) [!\[\]\(f419710cbe076aa30a9c6c031b5cbe84_img.jpg\)](#) [!\[\]\(2726020a4107bdc9042b257034f90eb3_img.jpg\)](#) [!\[\]\(9459655bf14a84f4d775e8d814cca8c9_img.jpg\)](#) [!\[\]\(de47dbdca34225b222a4a87ac0e499b3_img.jpg\)](#)

FULL STORY

A study published June 16 in *Biotropica* by a team of researchers at the University of Washington, the UW Burke Museum of Natural History and Culture, WWF Hong Kong and the University of Colombo has provided an important road map to conserving rough-nosed horned lizards in Sri Lanka.

Rough-nosed horned lizards, or *Ceratophora aspera*, are small lizards that live in Sri Lankan rainforests and aren't found anywhere else in the world. Characterized by the prominent horn in males, rough-nosed horned lizards live in moist, humid microhabitats in rainforests and palm groves. These lizards are particularly well-suited to understand the consequences of rainforest habitat destruction, climate change and the pet trade, as they are found throughout the lowland rainforests of Southwestern Sri Lanka. By analyzing the certain types of mutations in the genome of these lizards, the team was able to determine how geography and historical events impact the distribution of the remaining wild populations of rough-nosed horned lizards. Recent UW graduate Shanelle Wikramanayake led the study as part of her undergraduate work, and is now completing her master's degree at California State University, Northridge.

The team acquired DNA from the rough-nosed horned lizards by a catch-and-release approach of capturing wild lizards and taking tissue samples from the tips of their tails before releasing them back to the wild, which minimizes stress to the animals. Samples were taken across eight forests representing four forest groups from July through September 2018 and 2019, respectively. The remaining tissue samples will be housed along with photos of the lizards at the Department of Zoology and Environment Sciences at the University of Colombo. In addition, the DNA analysis from the team's work is also available online, where the samples and analysis are available for other researchers around the globe to study.

In addition to habitat destruction, nonnative species like cats and chickens eat these lizards, causing further harm to them and other prey animals. The team recommends a conservation strategy that considers the population structure, history and ecology of rough-nosed horned lizards to preserve genetic diversity.

The results from the DNA analysis found rough-nosed horned lizards are separated by distance into four forest groups: Southern Lowlands, Sabaragamuwa Hills, Central Highlands and Kithulgala. With the Sri Lankan government pledging to restore rainforest habitats, the results from the study can help inform guidelines for forest landscape restoration. For example, the similarities in populations and close geographic proximity between the lizards living in the Hiyare and Kottawa forest reserves in the Southern Lowlands forest group is a great candidate for habitat restoration. The two populations could be reconnected, increasing the chances for interbreeding, which could result in a stronger gene pool for future generations and less vulnerability to extinction.

"I hope this study brings attention to the unique and obscure biodiversity in the rainforests of Sri Lanka that require urgent conservation action," said Wikramanayake. "It's important to consider the genetic diversity in populations across their range when looking at conservation planning and strategies. This principle should also be considered in other forest ecosystems, where conservationists are planning restoration and habitat connectivity at landscape scales."

In addition to these lizards, the researchers are currently considering extending this work to other animals and other regions of Sri Lanka of high conservation value. In Sri Lanka, approximately 14% of mammals, 6.5% of birds, 75% of reptiles and 29% of flowering plant species aren't found anywhere else in the world. The methods of DNA analysis in this study not only inform conservation planning in Sri Lanka, but they also can be applied to species living in other rainforests that are at risk, including Madagascar, Borneo and the Amazon.

"I am proud and impressed with the hard work and passion that Shanelle Wikramanayake put into her project," said Adam Leaché, UW biology professor and Burke Museum curator of herpetology and genetic resources. "The Burke Museum is lucky to recruit such amazing undergraduates."

Story Source:

Materials provided by **University of Washington**. Original written by Andrea Godinez. *Note: Content may be edited for style and length.*

Journal Reference:

1. Shanelle A. Wikramanayake, Eric D. Wikramanayake, Nirmalie Pallewatta, Adam D. Leach�. **Integration of genetic structure into conservation of an endangered, endemic lizard, *Ceratophora aspera* A case study from Sri Lanka.** *Biotropica*, 2021; DOI: 10.1111/btp.12970

Cite This Page:

MLA	APA	Chicago
-----	-----	---------

University of Washington. "New DNA study provides critical information on conserving rainforest lizards." ScienceDaily. ScienceDaily, 2 August 2021. <www.sciencedaily.com/releases/2021/08/210802160707.htm>.

RELATED STORIES

Mosquito Control Program Reduces Dengue, Costs in Sri Lanka

June 5, 2019 — A public health, police, and military partnership to reduce the mosquito population in Sri Lanka resulted in a more than 50% reduction in dengue, as well as cost savings, finds a new ...

Homo Sapiens Colonized South Asian Rainforest by Hunting Small Mammals 45,000 Years Ago

Feb. 19, 2019 — A new study provides direct evidence for the hunting of tree-dwelling monkeys and other small mammals by Homo sapiens 45,000 years ago in the rainforest of Sri ...

Evidence for Persistent Forest Reliance by Indigenous Peoples in Historical Sri Lanka

Apr. 26, 2018 — Working closely with Wanniyalaeto (Vedda) elders in Sri Lanka during the repatriation of skeletal remains, a team of researchers have demonstrated that while some indigenous hunter-gatherers in Sri ...

El Niño Conditions in Pacific Precede Dengue Fever Epidemics in South Asia

Nov. 21, 2016 — Researchers have found a strong association between El Niño-Southern Oscillation conditions in the Pacific to observed weather and dengue epidemics in Sri Lanka. According to a study , el Niño ...

FROM AROUND THE WEB

ScienceDaily shares links with sites in the TrendMD network and earns revenue from third-party advertisers, where indicated.

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 [Email Newsletters](#)

 [RSS Feeds](#)

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

 [Facebook](#)

 [Twitter](#)

 [LinkedIn](#)

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 [Leave Feedback](#)

 [Contact Us](#)

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2021 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

