

[本期目录] [下期目录] [过刊浏览] [高级检索]

[打印本页] [关闭]

综述

马尔尼菲青霉的分子生物学研究进展

阙冬梅, 覃巍, 张军民

中山大学附属孙逸仙纪念医院皮肤性病科, 广州, 510120

摘要: 马尔尼菲青霉是一种主要累及免疫受损患者的条件致病性双相真菌, 近年来随着HIV感染的增多, 马尔尼菲青霉病发病率呈现逐年上升的趋势。国内外学者对马尔尼菲青霉及其致病机制进行了大量的研究, 尤其是分子生物学方法的应用, 使人们对马尔尼菲青霉的真菌学特点和马尔尼菲青霉病的发病机制有了更深入的了解, 使早期诊疗成为可能。该文主要从分子流行病学、分子遗传学和分子生物学诊断方面对马尔尼菲青霉进行综述。

关键词: 马尔尼菲青霉 深部真菌病 分子生物学

Abstract:

Keywords:

收稿日期 2010-01-19 修回日期 网络版发布日期

DOI:

基金项目:

广东省自然基金自由申请项目(8151008901000073)

通讯作者: 张军民, E-mail: junminmx@163.com

作者简介: 阙冬梅, 女(汉族), 硕士研究生在读, E-mail: domy2008@qq.com

作者Email: junminmx@163.com

参考文献:

- [1] Segretain G. *Penicillium marneffei* n.sp., agent of a mycosis of the reticuloendothelial system [J]. *Mycopathologia*, 1959, 30(11): 327-353.
- [2] Vanittanakom N, Cooper CR Jr, Fisher MC, et al. *Penicillium marneffei* infection and recent advances in the epidemiology and molecular biology aspects [J]. *Clin Microbiol Rev*, 2006, 19(1): 95-110.
- [3] Supparatpinyo K, Khamwan C, Bsosoung V, et al. Disseminated *Penicillium marneffei* infection in southeast Asia [J]. *Lancet*, 1994, 344(8915): 110-113.
- [4] Li PC, Tsui MC, Ma KF. *Penicillium marneffei*: indicator disease for AIDS in South East Asia [J]. *AIDS*, 1992, 6(2): 240-241.
- [5] 李云, 张悦, 王惠萱, 等. 马尔尼菲青霉菌的实验室诊断 [J]. 中国现代医学杂志, 2005, 15(18): 2743-2745.
- [6] 王莹, 邓卓霖, 马韵. 马尔尼菲青霉病的临床诊断进展 [J]. 中国真菌学杂志, 2007, 2(5): 312-314.
- [7] Vanittanakom N, Cooper CR Jr, Chariyalertsak S, et al. Restriction endonuclease analysis of *Penicillium*

扩展功能

本文信息

- ▶ Supporting info
- ▶ PDF (KB)
- ▶ [HTML全文]
- ▶ 参考文献[PDF]
- ▶ 参考文献

服务与反馈

- ▶ 把本文推荐给朋友
- ▶ 加入我的书架
- ▶ 加入引用管理器
- ▶ 引用本文
- ▶ Email Alert
- ▶ 文章反馈
- ▶ 浏览反馈信息

本文关键词相关文章

- ▶ 马尔尼菲青霉
- ▶ 深部真菌病
- ▶ 分子生物学

本文作者相关文章

- ▶ 阙冬梅
- ▶ 覃巍
- ▶ 张军民

PubMed

- ▶ Article by
- ▶ Article by
- ▶ Article by

- marneffei[J].J Clin Microbiol,1996,34(7):1834-1836.
- [8] Hsueh PR,Teng LJ,Hung CC,et al.Molecular evidence for strain dissemination of Penicillium marneffei: an emerging pathogen in Taiwan[J].J Infect Dis,2000,181(5):1706-1712.
- [9] Trewatcharegon S,Sirisinha S,Romsai A,et al.Molecular typing of Penicillium marneffei isolates from Thailand by NotI macrorestriction and pulsed-field gel electrophoresis[J].J Clin Microbiol,2001,39(12):4544-4548.
- [10] Imwidhaya P,Thipsuvan K,Chaiprasert A,et al.Penicillium marneffei: types and drug susceptibility [J].Mycopathologia,2001,149(3):109-115.
- [11] Lasker BA,Ran Y.Analysis of polymorphic microsatellite markers for typing Penicillium marneffei isolates[J].J Clin Microbiol,2004,42(4):1483-1490.
- [12] Fisher MC,Aanensen D,de Hoog S,et al.Muhilocs microsatellite typing system for Penicillium marneffei reveals spatially structured populations[J].J Clin Microbiol,2004,42(11):5065-5069.
- [13] Pasqualotto AC,Denning DW,Anderson MJ.A cautionary tale:lack of consistency in allele sizes between two laboratories for a published multilocus microsatellite typing system[J].J Clin Microbiol,2007,45(2):522-528.
- [14] Bomeman AR,Hynes MJ,Andrianopaidos A.The abaA homolog of Penicillium marneffei participates in two developmental programmes: conidiation and dimorphic growth[J].Mol Microbiol,2000,38(5):1034-1047.
- [15] Bomeman AR,Hynes MJ,Andrianopaulos A.An STE12 homolog from the asexual,dimorphic fungus Penicillium marneffei complements the defect in sexual development of an *Aspergillus nidulans* steA mutant[J].Genetics,2001,157(3):1003-1014.
- [16] Borneman AR,Hynes MJ,Andrianopaulos A.A basic helix-loophelix protein with similarity to the fungal morphological regulators,Phdl p,Efgl p and StuA,controls conidiation but not dimorphic growth in Penicillium marneffei[J].Mol Microbiol,2002,44(3):621-631.
- [17] Zuber S,Hynes MJ,Andrianopnalous A.G-protein signaling mediates asexual development at 25 degrees C but has no effect on yeast-like growth at 37 degrees C in the dimorphic fungus Penicillium mameffei[J].Eukaryot Cell,2002,1(3):440-447.
- [18] Zuber S,Hynes MJ,Andrianopoulos A.The G-protein alphasubunit GasC plays a major role in germination in the dimorphic fungus Penicillium marneffei[J].Genties,2003,164(2):487-499.
- [19] Boyce KJ,Hynes MJ,Andrianopoulos A.Control of morphogenesis and actin localization by the Penicillium marneffei RAC homolog[J].J Cell Sci,2003,116(7):1249-1260.
- [20] Boyce KJ,Hynes MJ,Andrianopoulos A.The Ras and Rho GT-Pases genetically interact to co-ordinately regulate cell polarity during development in Penicillium marneffei[J].Mol Microbiol,2005,55(5):1487-1501.
- [21] Boyce KJ,Andrianopoulos A.A p21-activated kinase is required for conidial germination in Penicillium marneffei[J].PLoS Pathog,2007,3(11):162.
- [22] Boyce KJ,schreider L,Andrianopoulos A.In vivo yeast cell morphogenesis is regulated by a p21-activated kinase in the human pathogen Penicillium marneffei[J].PLoS Pathos,2009,5(11):e1000678.
- [23] Pongsunk S,Andrianopoulos A,Chaiyaroj SC.Conditional lethal disruption of TATA-binding protein gene in Penicillium marneffei[J].Fungal Genet Biol,2005,42(11):893-903.
- [24] LoBuglio KF,Taylor JW.Phylogeny and PCR identification of the human pathogenic fungus Penicillium marneffei[J].J Clin Microbiol,1995,33(1):85-89.
- [25] Lindsley MD,Hurst SF,Iqbal NJ,et al.Rapid identification of dimorphic and yeast-like fungal pathogens using specific DNA probes[J].J Clin Microbiol,2001,39(10):3505-3511.
- [26] Vanittanakom N,Vanittanakom P,Hay RJ.Rapid identification of Penicillium marneffei by PCR-based detection of specific sequences on the rRNA gene[J].J Clin Microbiol,2002,40(5):1739-1742.
- [27] Prariyachatigul C,Chaiprasert A,Geenkajorn K,et al.Development and evaluation of a one-tube seminested PCR assay for the detection and identification of Penicillium marneffei[J].Mycoses,2003,46(11-12):447-454.
- [28] Tsunemi Y,Takahashi T,Tamaki T.Penicillium marneffei infection diagnosed by polymerase chain reaction from the skin specimen[J].J Am Acad Dermatol,2003,49(2):344-346.
- [29] Zeng H,Li X,Chen X,et al.Identification of Penicillium marneffei in paraffin-embedded tissue using nested PCR[J].Mycopathologia,2009,168(1):31-35.
- [30] 曹存巍,邓卓霖,罗虹,等.原位杂交技术鉴定马尔尼菲青霉[J].中国皮肤性病学杂志,2004,18(8):462-463.
- [31] Hsue HC,Huang YT,Kuo YL,et al.Rapid identification of fungal pathogens in positive blood cultures using oligonucleotide array hybridization[J].Clin Microbiol Infect,2010,16(5):493-500.
- [32] Pongpom M,Sirisanthana T,Vanittanakom N.Application of nested PCR to detect Penicillium marneffei in serum samples[J].Med Mycol,2009,47(5):549-553.
- [33] Pornprasert S,Praparattanapan J,Khamwan C,et al.Development of TaqMan real-time polymerase chain reaction for the detection and identification of Penicillium marneffei[J].Mycoses,2009,52(6):487-492.

本刊中的类似文章

1. 欧汝志, 卢祥婵, 李伟新.艾滋病合并马尔尼菲青霉病死亡的危险因素研究[J]. 中国真菌学杂志, 2011, 6(3):

158-161

2. 陈柏叡, 李若瑜.须癣毛癣菌分类进展[J]. 中国真菌学杂志, 2011, 6(1): 51-56

3. 陈柏叡, 孙毅, 胡小平, 万喆, 李若瑜.须癣毛癣菌的形态学及分子生物学鉴定[J]. 中国真菌学杂志, 2010,5(6): 321-326
4. 万力, 任强强, 王启明, 林元珠, 白逢彦.裂褶菌培养基形态学观察和DNA序列分析[J]. 中国真菌学杂志, 2010,5 (6): 327-331
5. 冯佩英, 黄怀球, 张静, 张晓辉, 孙九峰, 谢治, 鲁莎, 鲁长明, 席丽艳.小鼠巨噬细胞吞噬马尔尼菲青霉酵母相细胞的实验观察[J]. 中国真菌学杂志, 2010,5(6): 332-335
6. 王韵茹, 章强强.马拉色菌的免疫学与分子生物学研究新进展[J]. 中国真菌学杂志, 2010,5(4): 252-256
7. 姜红浩, 张宏, 余国荣, 张敬文, 张丁.1950~2007年中国深部真菌病历史地理学研究[J]. 中国真菌学杂志, 2010,5 (3): 148-153
8. 赵建荣, 樊移山, 董庭荣, 王舒.AIDS合并马尔尼菲青霉病36例分析[J]. 中国真菌学杂志, 2010,5(3): 158-161
9. 翟志芳, 王莉, 周村建, 徐艳, 杨希川, 钟白玉, 阎衡.马尔尼菲青霉病8例临床分析[J]. 中国真菌学杂志, 2010,5 (3): 162-164,169
10. 章强强, 赵颖.无绿藻的表型及其分子生物学鉴定[J]. 中国真菌学杂志, 2010,5(1): 9-12,25
11. 郭莉娜, 徐英春, 窦红涛, 谢秀丽, 李若瑜.临床相关毛孢子菌感染的研究现状[J]. 中国真菌学杂志, 2010,5 (1): 53-56

Copyright by 中国真菌学杂志