

Home
President's Welcome

NEWS CHANNELS

Arts & Architecture
Environment
Health & Medicine
Humanities & Social Sciences
International
Science & Engineering
Campus

MULTIMEDIA

Netcasts Videos
RSS Photos

Calendar of Events
Yale Bulletin
ELIne Newsletter
Poynter Fellowship
Yale in the News

Contact Public Affairs
About Yale
Yale University

FOUND IN: [Health & Medicine](#)

Brain Energy Use Key to Understanding Consciousness, Yale Researchers Find

Published: June 15, 2009

New Haven, Conn. — High levels of brain energy are required to maintain consciousness, a finding which suggests a new way to understand the properties of this still mysterious state of being, Yale University researchers report.

At its simplest, consciousness can be defined as the ability to respond meaningfully to external stimuli. Most studies of consciousness have used imaging technology to try to pinpoint areas of brain activity during tasks such as memorization or problem solving.

There are two problems with such an approach, said [Robert G. Shulman](#), Sterling Professor Emeritus of molecular biophysics and biochemistry at Yale and lead author of the paper, to be published this week in the online edition of the journal *Proceedings of the National Academy of Sciences*. First, functional magnetic resonance imaging has shown that many areas of the brain, not just one or two, are recruited during tasks such as memory tests and are scant help in studying the state of being conscious. Second, the amount of energy used in such tasks is minute, about one percent of baseline energy available to the brain.

" Neuroimaging has been looking at the tip of the iceberg," Shulman said. " We looked at the rest of the iceberg."

What is the other 99 percent of energy consumption doing?

Shulman and colleagues have proposed that it is needed to maintain a person in a state of consciousness. Heavily anesthetized people are known to show approximately 50 percent reductions in cerebral energy consumption. When the paws of lightly anesthetized rats with rather high baseline energy levels were stroked, fMRI signals were received in the sensory cortex and in many other areas of the brain. In heavily anesthetized rats the signal stopped at the sensory cortex. Both the total energy and the fMRI signals changed when the person or animal lost consciousness.

" What we propose is that a conscious person requires a high level of brain energy," Shulman said.

The finding has profound implications for our understanding of the connection between the brain and consciousness, Shulman said. " You can think of consciousness not as a property of the brain, but of the person."

Anesthesiologists consider a person to be in a behavioral state of consciousness when he or she can respond to simple stimuli. Properties of this state, such as the high energy and the delocalized fMRI signals, allow the person to perform the interconnected activities that make up our everyday lives. Shulman suggests that these more energetic properties of the brain support human behavior and should be considered when interpreting the much weaker signals that are typically recorded during fMRI studies.

Other Yale researchers involved in the study are professors Fahmeed Hyder and Douglas L. Rothman.

The study was funded by the National Institutes of Health.

PRESS CONTACT: [Bill Hathaway](#) 203-432-1322

Print E-Mail Share

PRESS CONTACT

Bill Hathaway
203-432-1322

OTHER ARTICLES

[Researchers Suggests Gene Inhibition May Help Normalize Type 2 Diabetes](#)

[The Season of Ticks: Could Climate Change Worsen Lyme Disease?](#)

[Health Experts Make Case for Tax on Sugar-Sweetened Beverages](#)

[Yale Stem Cell Researchers Reap State Grants](#)

RELATED LINKS

[Yale Cancer Center](#)

[Yale School of Medicine](#)

[Yale University School of Nursing](#)

[Yale School of Public Health](#)

[Yale-New Haven Hospital](#)

[BACK TO TOP OF PAGE ^](#)

