



Nuclear Experiment

# Anisotropic flow at the LHC measured with the ALICE detector

Raimond Snellings, for the ALICE Collaboration

(Submitted on 30 Jun 2011 (v1), last revised 10 Jul 2011 (this version, v2))

The ALICE detector at the LHC recorded first Pb-Pb collisions at 2.76 TeV in November and December of 2010. We report on the measurements of anisotropic flow for charged and identified particles. From the comparison with measurements at lower energies and with model predictions we find that the system created at these collision energies is described well by hydrodynamical model calculations and behaves like an almost perfect fluid.

Comments: 8 pages, 16 figures, plenary talk at Quark Matter 2011, May 23rd-28th 2011, Annecy, France

Subjects: **Nuclear Experiment (nucl-ex)**; High Energy Physics - Experiment (hep-ex)

Cite as: [arXiv:1106.6284v2](#) [nucl-ex]

## Submission history

From: Raimond Snellings [[view email](#)]

[v1] Thu, 30 Jun 2011 16:14:09 GMT (303kb,D)

[v2] Sun, 10 Jul 2011 19:06:16 GMT (303kb,D)

*Which authors of this paper are endorsers?*

## Download:

- [PDF](#)
- [Other formats](#)

Current browse context:

nucl-ex

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[hep-ex](#)

## References & Citations

- [SLAC-SPIRES HEP](#)  
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark ([what is this?](#))

