

H-ZSM-5 分子筛上环己烯芳构化反应历程的理论研究

左士颖, 周丹红*, 任珏, 王凤娇

辽宁师范大学化学化工学院功能材料化学研究所, 辽宁大连 116029

ZUO Shiyong, ZHOU Danhong*, REN Jue, WANG Fengjiao

Institute of Chemistry for Functionalized Materials, College of Chemistry and Chemical Engineering, Liaoning Normal University, Dalian 116029, Liaoning, China

- 摘要
- 参考文献
- 相关文章

Download: PDF (420KB) [HTML](#) (1KB) Export: BibTeX or EndNote (RIS) Supporting Info

摘要 基于 76T 簇模型, 采用量子力学和分子力学联合的 ONIOM2 (B3LYP/6-31G(d, p):UFF) 方法研究了 H-ZSM-5 分子筛上环己烯芳构化反应历程。结果表明, 环己烯首先吸附在分子筛酸性位上, 与酸性质子共同脱除一个 H₂ 分子后, 在分子筛骨架氧上生成烷氧配合物中间体; 然后再脱质子得到环己二烯, 同时酸性位复原; 再经历脱氢和脱质子历程, 最后得到产物苯, 并吸附在复原的分子筛酸性位上。计算得到脱氢的活化能依次为 279.64 和 260.21 kJ/mol, 脱质子的活化能依次为 74.64 和 59.14 kJ/mol。所有脱氢反应都是吸热过程, 生成表面烷氧活性中间体, 随后的脱质子反应能垒较低, 而且是放热过程。此外, 比较了环己烯在分子筛酸性位上的三个竞争反应, 即脱氢、质子化和氢交换反应的活化能垒, 证明环己烯优先发生脱氢反应。

关键词: H-ZSM-5 分子筛 环己烯 芳构化 脱氢 量子力学 / 分子力学

Abstract: By using the QM/MM hybrid ONIOM2 (B3LYP/6-31G(d,p):UFF) method, the dehydro-aromatization reaction of cyclohexene over H-ZSM-5 zeolite was studied. The calculation was based on the 76T cluster model with Al substituted at the T12 site. The results indicated that cyclohexene was adsorbed on the Brønsted acid (B-acid) site of H-ZSM-5 zeolite. The dehydrogenation reaction consisted of cleavage of a C-H bond by acidic proton. After release of dihydrogen, the carbonium moiety was bonded with the basic framework oxygen, resulting in surface cyclohexene alkoxide intermediate. Then the alkoxide released the proton to the framework oxygen which led to generation of B-acid site, and the produced cyclohexene was adsorbed on the B-acid site. After subsequent dehydrogenation and deprotonation steps, a benzene molecule was produced as final product. The activation energy for dehydrogenation steps was 279.64 and 260.21 kJ/mol, respectively, and the activation energy for deprotonation process was 74.64 and 59.14 kJ/mol, respectively. All dehydrogenation steps are endothermic to lead to surface alkoxide intermediates, and the subsequent deprotonation have lower energy barriers and are exothermic. In addition, the three competitive reactions of cyclohexene, i.e. protonation, hydrogen exchange, and dehydrogenation, were also compared. It was found that cyclohexene preferred to dehydrogenate with zeolite proton.

Keywords: H-ZSM-5 zeolite, cyclohexene, aromatization, dehydrogenation, QM/MM

收稿日期: 2012-03-23; 出版日期: 2012-07-31

引用本文:

左士颖, 周丹红, 任珏等. H-ZSM-5 分子筛上环己烯芳构化反应历程的理论研究[J] 催化学报, 2012, V33(8): 1367-1373

ZUO Shi-Ying, ZHOU Dan-Hong, REN Jue etc. Theoretical Study of Reaction Mechanism of Cyclohexene Aromatization over H-ZSM-5 Zeolite[J] Chinese Journal of Catalysis, 2012, V33(8): 1367-1373

链接本文:

<http://www.chxb.cn/CN/10.3724/SP.J.1088.2012.20346> 或 <http://www.chxb.cn/CN/Y2012/V33/I8/1367>

- [1] Corma A. Chem Rev, 1995, 95: 559
- [2] Tukur N M, Al-Khattaf S. Chem Eng Process, 2005, 44: 1257
- [3] Xu Y D, Shu Y Y, Liu S T, Huang J S, Guo X X. Catal Lett, 1995, 35: 233
- [4] Solymosi F, Erdöhelyi A, Szöke A. Catal Lett, 1995, 32: 43
- [5] 邢双英, 周丹红, 曹亮, 李新. 催化学报 (Xing Sh Y, Zhou D H, Cao L, Li X. Chin J Catal), 2010, 31: 415
- [6] 张佳, 周丹红, 倪丹. 催化学报 (Zhang J, Zhou D H, Ni D. Chin J Catal), 2008, 29: 715
- [7] 曹亮, 周丹红, 邢双英, 李新. 催化学报 (Cao L, Zhou D H, Xing Sh Y, Li X. Chin J Catal), 2010, 31: 645
- [8] Derouane E G, Fripiat J G. J Phys Chem, 1987, 91: 145

Service

- ▶ 把本文推荐给朋友
- ▶ 加入我的书架
- ▶ 加入引用管理器
- ▶ Email Alert
- ▶ RSS

作者相关文章

- ▶ 左士颖
- ▶ 周丹红
- ▶ 任珏
- ▶ 王凤娇

- [9] Fripiat J G, Berger-Andre F, Andre J M, Deroane E G. Zeolites, 1983, 3: 306
- [10] Rozanska X, van Santen R A, Hutschka F. J Phys Chem B, 2002, 106: 4652
- [11] Chatterjee A, Bhattacharya D, Chatterjee M, Iwasaki T. Microporous Mesoporous Mater, 1999, 32: 189
- [12] Chu Y Y, Han B, Fang H J, Zheng A M, Deng F. Microporous Mesoporous Mater, 2012, 151: 241
- [13] Zheng A M, Deng F, Liu Sh B. Catal Today, 2011, 164: 40
- [14] Vreven T, Byun K S, Komáromi I, Dapprich S, Montgomery J A Jr, Morokuma K, Frisch M J. J Chem Theory Comput, 2006, 2: 815
- [15] Maseras F, Morokuma K. J Comput Chem, 1995, 16: 1170
- [16] Humbel S, Sieber S, Morokuma K. J Chem Phys, 1996, 105: 1959
- [17] Vreven T, Morokuma K. J Comput Chem, 2000, 21: 1419
- [18] Boronat M, Viruela P M, Corma A. J Am Chem Soc, 2004, 126: 3300
- [19] Joshi Y V, Thomson K T. J Catal, 2005, 230: 440
- [20] Jansang B, Nanok T, Limtrakul J. J Phys Chem B, 2006, 110: 12626
- [21] Rungsisakun R, Jansang B, Pantu P, Limtrakul J. J Mol Struct, 2005, 733: 239
- [22] 李惊鸿, 周丹红, 任珏. 物理化学学报 (Li J H, Zhou D H, Ren J. Acta Phys-Chim Sin), 2011, 27: 1393
- [23] Sun Y X, Yang J, Zhao L F, Dai J X, Sun H. J Phys Chem C, 2010, 114: 5975
- [24] Derouane E G, Chang C D. Microporous Mesoporous Mater, 2000, 35-36: 425
- [25] Derouane E G. J Mol Catal A, 1998, 134: 29
- [26] Kasuriya S, Namuangruk S, Treesukol P, Tirtowidjojo M, Limtrakul J. J Catal, 2003, 219: 320
- [27] Namuangruk S, Pantu P, Limtrakul J. Chem Phys Chem, 2005, 6: 1333
- [28] Derouane E G, Fripiat J G. Zeolites, 1985, 5: 165
- [29] Lonsinger S R, Chakraborty A K, Theodorou D N, Bell A T. Catal Lett, 1991, 11: 209
- [30] Zheng X B, Blowers P. J Phys Chem A, 2006, 110: 2455
- [31] Kazansky V B. Catal Today, 1999, 51: 419
- [32] Chatterjee A, Vetrivel R. Microporous Mesoporous Mater, 1994, 3: 211
- [33] Van Koningveld H, Van Bekkum H, Jansen J C. Acta Crystallogr B, 1987, 43: 127
- [34] Frisch M J, Trucks G W, Schlegel H B, Scuseria G E, Robb M A, Cheeseman J R, Montgomery J A Jr, Vreven T, Kudin K N, Burant J C, et al. Gaussian 03, Revision C.02 Ed. Wallingford, CT: Gaussian, Inc., 2004
- [35] Panjan W, Limtrakul J. J Mol Struct, 2003, 654: 35
- [36] Correa R J, Mota C J A. PhysChemChemPhys, 2002, 4: 375
- [37] Cuán A, Martínez-Magadán J M, García-Cruz I, Galván M. J Mol Catal A, 2005, 236: 194
- [38] Furtado E A, Milas I, Milam De Albuquerque Lins J O, Chaer Nascimento M A. Phys Status Solid A, 2001, 187: 275
- [39] Gounder R, Iglesia E, J Am Chem Soc, 2009, 31: 1958

- [1] 王迪勇, 王金渠, 杨建华, 鲁金明, 殷德宏, 张艳. 蒸气相法 ZSM-5 分子筛的合成及其负载的 Mo 催化剂在甲烷芳构化中的应用[J]. 催化学报, 2012,33(8): 1383-1388
- [2] 高旭锋, 谌春林, 任士远, 张建, 苏党生. 氧化铈的结构对其热稳定性及催化丙烷氧化脱氢反应性能的影响[J]. 催化学报, 2012,33(7): 1069-1074
- [3] 孙海杰, 潘雅洁, 王红霞, 董英英, 刘仲毅, 刘寿长. 二乙醇胺作添加剂 Ru-Zn 催化剂上苯选择加氢制环己烯[J]. 催化学报, 2012,33(4): 610-620
- [4] 刘文芳, 侯本象, 侯延慧, 赵之平. 中空纤维膜固定化甲酸脱氢酶催化 CO₂ 合成甲酸[J]. 催化学报, 2012,33(4): 730-735
- [5] 施梅勤, 陈宁宁, 马淳安, 李瑛, 魏爱平. 双功能 WC/HZSM-5 催化剂上正己烷芳构化反应性能[J]. 催化学报, 2012,33(3): 570-575
- [6] 周功兵, 刘建良, 许可, 裴燕, 乔明华, 范康年. 载体焙烧温度对 Ru/MgAl₂O₄ 催化剂液相苯部分加氢性能的影响[J]. 催化学报, 2011,32(9): 1537-1544
- [7] 俞佳枫, 方雯, 葛庆杰, 徐恒泳. 助剂形态对 Pt/V-Al₂O₃ 催化剂抗积炭性能的影响[J]. 催化学报, 2011,32(8): 1364-1369
- [8] 孙果宋, 黄青则, 李会泉, 柳海涛, 张钊, 王兴瑞, 王秋萍, 王金淑. 不同载体负载的 Cr 基催化剂催化 CO₂ 氧化异丁烷脱氢制异丁烯[J]. 催化学报, 2011,32(8): 1424-1429
- [9] 李玉宁, 任丽萍, 李亚男, 金照生, 滕加伟, 杨为民. 无粘结剂成型的 Zn/ZSM-5 催化剂上混合碳四烃类芳构化反应性能[J]. 催化学报, 2011,32(6): 992-996
- [10] 李巧灵 1, 张元华 1, 陈世萍 1, 方维平 1,2, 杨意泉 1,2. 反映工艺条件对管式反应器催化反应影响的转化率方程[J]. 催化学报, 2011,32(3): 446-450
- [11] 杨建华, 于素霞, 胡慧晔, 初乃波, 鲁金明, 殷德宏, 王金渠. 无第二模板剂法合成多级结构 ZSM-5 分子筛微球及其在甲烷无氧芳构化反应中的应用[J]. 催化学报, 2011,32(2): 362-367
- [12] 孙海杰, 周小莉, 陈志浩, 郭伟, 刘仲毅, 刘寿长. 单层分散型 Ru-Zn 催化剂及其催化苯选择加氢制环己烯的性能[J]. 催化学报, 2011,32(2): 224-230
- [13] 王晓丽, 吴功德, 王芳, 魏伟, 孙予罕. 非均相 Schiff 碱铬 (III) 配合物催化环己烯环氧化性能的研究[J]. 催化学报, 2011,32(12): 1812-1821
- [14] 秦燕飞, 薛伟, 李芳, 王延吉, 魏碧芳. Ru-[bmim]BF₄ 的制备及催化苯选择加氢反应性能[J]. 催化学报, 2011,32(11): 1727-1732

