Scientific Research

December 2001.



Search Keywords, Title, Author, ISBN, ISSN

Home	Journals	Books	Conferences	News	About Us	s Job	
Home > Journal > Earth & Environmental Sciences > JEP					Open Special Issues		
Indexing View Papers Aims & Scope Editorial Board Guideline Article Processing Charges					Published Special Issues		
JEP> Vol.2 No.4, June 2011					Special Issues Guideline		
OPEN@ACCESS Radioactivity in Surface Soils Around the Proposed Sites for						JEP Subscription	
Titanium Mining Project in Kenya						Most popular papers in JEP	
PDF (Size: 366KB) PP. 460-464 DOI: 10.4236/jep.2011.24053 Author(s)						About JEP News	
M. K Osoro, I. V. S. Rathore, M. J Mangala, A. O Mustapha ABSTRACT Radioactivity measurements were carried out around Maumba and Nguluku villages, two of the proposed sites for titanium mining in the coastal area of Kenya. Samples of surface soils were analyzed using a HPGe gamma spectrometer. The average activity concentrations for ²²⁶ Ra, ²³² Th and ⁴⁰ K are 20.9 ± 7.6, 27.6 ± 9.1 and 69.5 ± 16.5 Bqkg ^{-1<} , respectively. The absorbed dose rates in air, calculated on the basis of the measured activity concentrations, range from 9.8 to 50.0 nGyh ⁻¹ , with an average of 29.2 nGyh ⁻¹ . These values are below the global population-weighted mean, and they should be considered when planning appropriate monitoring and surveillance programmes during the mining operation, as well as the reclamation and restoration programmes after mining. KEYWORDS Radioactivity Measurements, Titanium Mining, Absorbed Dose Rates, Natural Background Radiation					Frequently Asked Questions		
					Recommend to Peers		
					Recommend to Library		
					Contact Us		
					Downloads:	301,497	
					Visits:	673,121	
Cite this paper M. Osoro, I. Rathore, M. Mangala and A. Mustapha, "Radioactivity in Surface Soils Around the Proposed Sites for Titanium Mining Project in Kenya," <i>Journal of Environmental Protection</i> , Vol. 2 No. 4, 2011, pp. 460-464.					Sponsors, Associates, a Links >>		
 doi: 10.4236/jep.2011.24053. References [1] IAEA (International Atomic Energy Agency), " Extent of Environmental Contamination by Naturally Occurring Radioactive Material (NORM) and Technological Options for Mitigation," Technical report series, No. 419, IAEA, Vienna, 2003. 					The International Conference Pollution and Treatment Technology (PTT 2013)		
			ection," NORM Industries iation IRPA11, Madrid, 23				
·	RI (Tiomin Resources Inc.), Technical report on Kwale, 2004. Available at ttp://www.tiomin.com/s/NewsReleases.asp						
	(Kwale District)," In A		Metal Prospecting in the - Austria mineral explora				
			nental Issues: The Case (IFD) Project Launch mee	•			

[6] IAEA (International Atomic Energy Agency), " Soil sampling for environmental contaminants," IAEA-TECDOC- 1415, IAEA, Vienna, 2004.

[7] A. O. Mustapha, J. P. Patel, and I. V. S. Rathore, "Assessment of Human Exposures to Natural Sources of Radiation in Kenya," Radiation Protection Dosimetry, Vol 82, No. 4, 1999, pp. 285-292.

[8] N. O. Hashim, I. V. S. Rathore, A. M. Kinyua and A. O. Mustapha, "Natural and Artificial Radioactivity Levels in Sediments Along the Kenyan Coast," Radiation Physics and Chemistry, Vol. 71, No. 3-4, 2004, pp. 805-806. doi:10.1016/j.radphyschem.2004.04.101