



F₁代法检测棉铃虫种群对Bt棉的抗性等位基因频率变化

潘利东, 施明, 张凯, 陈进, 高聪芬*

南京农业大学植物保护学院农药科学系/教育部病虫监测与治理重点开放实验室, 南京 210095

Analysis of Frequency of Resistance Allele to Transgenic Bt Cotton in Field Populations of *Helicoverpa armigera*(Hübner) (Lepidoptera: Noctuidae) with F₁ Screening Method

PAN Li-dong, SHI Ming, ZHANG Kai, CHEN Jin, GAO Cong-fen**

Department of Pesticide Science, College of Plant Protection, Nanjing Agricultural University / Key Laboratory of Integrated Management of Crop Diseases and Pests, Ministry of Education, Nanjing 210095, China

摘要

参考文献

相关文章

Download: PDF (807KB) HTML 1KB Export: BibTeX or EndNote (RIS) Supporting Info

摘要 在具有室内高抗Bt棉棉铃虫品系的前提下, F₁代法是检测棉铃虫对Bt棉抗性等位基因频率的最简捷、快速的生物测定方法。2010—2012年采用F₁代法检测了河北省邱县棉铃虫田间种群对Bt棉的抗性等位基因频率变化。结果表明, 2010—2012年采集的122、141及124头田间雄虫中, 分别有32、22及43头携带抗性基因, 抗性等位基因频率分别为0.131(95%置信限CI:0.101~0.162)、0.078(95% CI:0.034~0.122)及0.199(95% CI:0.124~0.274)。2012年采用带毒饲料法测定河北邱县等4个地区Bt棉田间棉铃虫对Cry1Ac的抗性。结果表明, 河北邱县棉铃虫种群的抗性最高(抗性倍数RR为19.2倍), 明显高于湖北荆州、湖北枣阳和安徽萧县种群的抗性(RR为4.9~9.3倍)。该地区长期大面积种植转基因Bt棉导致田间棉铃虫抗性进化, 需要尽快采取有效的抗性治理措施。

关键词: 转Bt棉 棉铃虫 F₁代法 抗性基因频率 抗性检测

Abstract: F₁ screening method is the best bioassay measure to detect the resistance allele frequency of *Helicoverpa armigera* (Hübner) to transgenic Bt cotton if there is a high Bt-resistant strain. From 2010 to 2012, the frequency of allele conferring resistance to Bt cotton in Qiuxian County, Hebei Province, was estimated by using the F₁ screening method. The results showed that from 2010 to 2012, 32 out of 122, 22 out of 141 and 22 out of 141 field-collected male moths were detected to carry resistance alleles and the resistance allele frequency was estimated as 0.131 (95% confidence interval (CI): 0.101-0.162), 0.078 (95% CI: 0.034-0.122) and 0.199 (95% CI: 0.124-0.274), respectively. Bioassay of *Helicoverpa armigera* populations on Bt diet was adopted to detect resistance to Cry1Ac in 2012. The results showed that the resistance of field population from Qiuxian County was the highest, the resistance ratio reached 19.2 fold, compared with susceptible strain, and the resistance ratio of other three populations which ranged between 4.9 and 9.3 fold. Long-term adoption of planting uni-toxin-producing Bt cotton, and lack of conventional cotton refuge system may accelerate the resistance evolution in this region, and it is necessary to establish and implement effective resistance management strategy as soon as possible.

Keywords: *Helicoverpa armigera* transgenic Bt cotton F₁ screening method resistance allele frequency resistance monitoring

Received 2013-02-25;

Corresponding Authors: gaocongfen@njau.edu.cn

About author: 潘利东(1987-), 男, 硕士研究生

引用本文:

潘利东, 施明, 张凯, 陈进, 高聪芬. F₁代法检测棉铃虫种群对Bt棉的抗性等位基因频率变化[J] 棉花学报, 2013, V25(3): 240-246

PAN Li-Dong, SHI Ming, ZHANG Kai, CHEN Jin, GAO Cong-Fen. Analysis of Frequency of Resistance Allele to Transgenic Bt Cotton in Field Populations of *Helicoverpa armigera*(Hübner) (Lepidoptera: Noctuidae) with F₁ Screening Method[J] Cotton Science, 2013, V25(3): 240-246

链接本文:

http://journal.cricaas.com.cn:8082/mhxb/CN/1002-7807(2013)03-0240-07 或 http://journal.cricaas.com.cn:8082/mhxb/CN/Y2013/V25/I3/240

Service

- ▶ 把本文推荐给朋友
- ▶ 加入我的书架
- ▶ 加入引用管理器
- ▶ Email Alert
- ▶ RSS

作者相关文章

- ▶ 潘利东
- ▶ 施明
- ▶ 张凯
- ▶ 陈进
- ▶ 高聪芬