

[Available Issues](#) | [Japanese](#)>> [Publisher Site](#)Author: Keyword: 

Search

ADVANCED

[TOP](#) > [Available Issues](#) > [Table of Contents](#) > [Abstract](#)

ONLINE ISSN : 1882-0484

PRINT ISSN : 0031-9473

Japanese Journal of Phytopathology

Vol. 72 (2006) , No. 4 pp.200-201

[\[PDF \(716K\)\]](#) [\[References\]](#)

## Black spot on flowering cabbage caused by *Alternaria brassicicola* (Schwein.) Wiltshire in Japan

J. NISHIKAWA<sup>1)</sup>

1) Kakegawa Research Center, Sakata Seed Co.

(Received February 24, 2006)

(Accepted May 15, 2006)

### ABSTRACT

A black spot disease on flowering cabbage (*Brassica oleracea* L. var. *acephala* DC.) was found in Shizuoka Prefecture, Japan in March 2003. A fungus was isolated and used to inoculate cabbage leaves to reproduce the black spot symptoms. It was identified as *Alternaria brassicicola*. This is the first record of the disease in Japan and is named black spot (Kokuhan-byo) on flowering cabbage.

**Key words:** flowering cabbage, *Alternaria brassicicola*, black spot

[\[PDF \(716K\)\]](#) [\[References\]](#)Download Meta of Article [\[Help\]](#)[RIS](#)[BibTeX](#)

To cite this article:

J. NISHIKAWA (2006). Black spot on flowering cabbage caused by *Alternaria brassicicola* (Schwein.) Wiltshire in Japan . Japanese Journal of Phytopathology 72: 200-201 .

doi:10.3186/jjphytopath.72.200

JOI JST.JSTAGE/jjphytopath/72.200



---

[Japan Science and Technology Information Aggregator, Electronic](#)

