

[本期目录](#) | [下期目录](#) | [过刊浏览](#) | [高级检索](#)[\[打印本页\]](#) [\[关闭\]](#)**农学—研究报告****大茂山地区苏云金芽孢杆菌分布与多样性研究**宋健¹,杜立新²,王容燕²,魏利民^{3,3},曹伟平²,宋健²,王金耀²,冯书亮²

1. 河北大学生命科学学院
 2. 河北省农林科学院植物保护研究所
 3.

摘要:

为了对大茂山地区苏云金芽孢杆菌的分布与多样性进行研究,本试验利用温度-抗生素筛选法从大茂山不同海拔地区采集的784份土壤样品中分离得到76株苏云金芽孢杆菌,发现在海拔900~1200 m之间为土壤最佳采集区域。对这76株菌株进行光学显微观察,发现其伴孢晶体呈现球形、菱形、方形、不规则形等多种形态;PCR-RFLP对分离获得的76株菌株基因鉴定发现,主要包含cry1、cry2等7种基因类型,其中有28株未鉴定出基因型;其SDS-PAGE晶体蛋白分析发现,这些菌株主要表达130 kDa、97 kDa、66 kDa和45 kDa蛋白。对铜绿丽金龟进行生物活性测定,17株具有高毒力的菌株中的16株含有cry8Ca基因,另一株未鉴定出已有基因型。从晶体形态、基因类型、SDS-PAGE检测以及生物活性方面均说明苏云金芽孢杆菌在河北省大茂山地区的分布具有良好的多样性。

关键词: 苏云金芽孢杆菌; 海拔; PCR-RFLP; SDS-PAGE

Research on the Distribution and Diversity of *Bacillus thuringiensis* from Damaoshan Mountains

Abstract:

By the temperature-antibiotic method, the author isolated 74 strains *Bacillus thuringiensis* from totally 764 soil samples which were collected from different altitude Damaoshan mountains of Hebei province. The results showed the optimal collecting area at an altitude of 900 to 1200 m. Light microscope observation showed that the crystals of these Bt strains were in shape of spherical, bipyramidal, square, irregular and so on; 7 kinds of cry gene were found by PCR-RFLP system, and 11 Bt strains harbour unknown cry genes; The results of SDS-PAGE analysis showed that these Bt strains mainly expressed 130 kDa, 97 kDa, 66 kDa and 45 kDa proteins. 17 strains were shown to be highly toxic to scarabaeoidae larvae, 16 of them contained the gene of cry8Ca, one of them could not be found any known cry gene. All of these showed the great diversity of Bt strains in Damaoshan mountains of Hebei province.

Keywords: *Bacillus thuringiensis* altitude RCP-RFLP SDS-PAG**收稿日期** 2010-11-24 **修回日期** 2010-12-02 **网络版发布日期** 2011-02-18**DOI:****基金项目:**

国家863;国家支撑;国家支撑

通讯作者: 冯书亮 河北省农业有害生物综合防治工程技术研究中心, 河北保定071000**作者简介:**

作者Email: fshliang@163.com

扩展功能
本文信息
Supporting info
PDF(1177KB)
[HTML全文]
参考文献[PDF]
参考文献
服务与反馈
把本文推荐给朋友
加入我的书架
加入引用管理器
引用本文
Email Alert
文章反馈
浏览反馈信息
本文关键词相关文章
苏云金芽孢杆菌; 海拔; PCR-RFLP; SDS-PAGE
本文作者相关文章
宋健
杜立新
王容燕
魏利民
曹伟平
宋健
王金耀
冯书亮
PubMed
Article by Song,J
Article by Du,L.X
Article by Yu,R.Y
Article by Wei,L.M
Article by Cao,W.B
Article by Song,J
Article by Yu,J.Y
Article by Feng,S.L

参考文献：

- [1] Schnepf E,Crickmore N,Van Rie J,et al.Bacillus thuringiensis and its pesticidal crystal proteins.Microbiol Mol Biol Rev,1998,62: 775-806.
- [2] Feitelson J S.The Bacillus thuringiensis family tree.In: Kim,L.(Ed.) Advanced Engineered Pesticides.New York: Marcel Dekker,1993: 63-72.
- [3] Suzuki,N,Hori,H,Ogiwara,K.,et al.Insecticidal spectrum of a novel Isolate of Bacillus thuringiensis serovar japonensis.Biological control,1992,2: 138-142.
- [4] Pinto L M,Azambuja A O,Diehl E and Fiúza L M.Pathogenicity of Bacillus thuringiensis isolated from two species of Acromyrmex (Hymenoptera,Formicidae).Braz.J.Biol.,2003,63(2): 6-301.
- [5] 苏旭东,马雯,马晓燕,等.一株苏云金芽孢杆菌的分离与杀虫基因的鉴定.华北农学报.2007,22(4): 184-188.
- [6] Kuo W S,Chak K F.Identification of novel cry-type genes from Bacillus thuringiensis strains on the basis of restriction fragment length polymorphism of the PCR amplified DNA.Appl Environ Microbiol,1996,62(4): 1369-1377.
- [7] 宋福平,张杰,黄大昉,等.苏云金芽孢杆菌cry基因PCR-RFLP鉴定体系的建立.中国农业科学,1998,31(3): 13-18.
- [8] J.萨姆布鲁克,D.V.拉塞尔著《分子克隆实验指南》.黄培堂译.第三版.北京:科学出版社,2002: 418-424.
- [9] 冯书亮,付韵琴,范秀华等.几株高效苏云金杆菌菌株对玉米螟、棉铃虫、粘虫和黄地老虎的毒力测定,中国生物防治,1995,11(1): 22-25.
- [10] 潘晓鸿,王飞鹏,姚荣英等.不同生境Bt分离株晶体含量与靶标昆虫生物活性关系分析.中国农学通报,2009,25(11): 156-160.
- [11] Leckie S E,Prescott C E,Grayston J,et al.Characterization of humus microbial communities in adjacent forest types that differ in nitrogen availability.Microbm Ecol,2004,48(1): 29-40.
- [12] 王津红,吴光辉,陈月华,等.中国苏云金芽孢杆菌的分布与cry基因的多样性[J].微生物通报,2001,28(3): 50-55.
- [13] 胡晓婷,宋健,王容燕,等.对铜绿丽金龟幼虫有活性的苏云金芽孢杆菌菌株的分离及鉴定,华北农学报,2008,23(6): 81-83.

本刊中的类似文章

Copyright by 中国农学通报