

植物保护—研究报告

不同农药单剂及混配微乳剂的环境毒性研究

高越¹,张润祥^{2,2},王振^{2,2},封云涛^{2,2},范仁俊¹

1. 山西省农科院植保所

2.

摘要:

为了研究高效氯氰菊酯、阿维菌素、甲维盐等9种单剂或混配微乳剂的安全性,笔者测试了其对环境生物的毒性大小。根据《化学农药环境安全评价试验准则》,对斑马鱼、鹌鹑、蜜蜂、桑蚕分别采用半静水式鱼毒测定法、经口染毒发、接触染毒法、食下毒叶法进行测试。在鱼毒性方面,0.2%高渗甲维盐、3%啶虫脒和20%阿维?杀虫单微乳剂表现为低或中毒,其余药剂均表现为高毒或剧毒;在鸟毒性方面,4.5%高效氯氰菊酯微乳剂表现为高毒,其余药剂均表现为低毒或中毒;在蜜蜂毒性方面,0.2%高渗甲维盐和3%啶虫脒微乳剂表现为低毒或中毒,其余均表现为高毒;在桑蚕毒性方面,所有药剂均表现为高毒或剧毒。说明啶虫脒和低含量甲维盐对鱼、鸟、蜂、蚕毒性较低,而拟除虫菊酯类杀虫剂、阿维菌素及高含量甲维盐风险性较高,在施用时应远离鱼塘、蜂场及桑树。

关键词: 环境毒性

Environmental Toxicity of Several Different Pesticides Single or Mixed Micro-Emulsion

Abstract:

In order to study the safety of beta-cypermethrin, avermectin, emamectin benzoate, et al, 9 single or mixed micro-emulsion, the authors tested their environmental toxicity. Using semi-hydrostatic-type, oral exposure method, food intake method, contact exposure method to test the toxicity to fish, bird, honeybees, silkworm, respectively according to 'The guidelines of chemical pesticides environmental safety evaluation'. On the toxicity to fish, 0.2% hypertonic emamectin benzoate, 3% acetamiprid and 20% avermectin. monosultap were evaluated as 'low toxic class' or 'moderate toxic class', others were 'high toxic class' or 'hypertoxic class'; on the bird toxicity, 4.5% beta-cypermethrin was evaluated as 'high toxic class', others were 'low toxic class' or 'moderate toxic class'; on the toxicity to honeybees, 0.2% hypertonic emamectin benzoate and 3% acetamiprid were evaluated as 'low toxic class' or 'moderate toxic class', others were 'high toxic class'; on the toxicity to silkworm, all the pesticides were evaluated as 'high toxic class' or 'hypertoxic class'. Shows that acetamiprid and low content emamectin benzoate behaved low toxicity to fish, bird, honeybees, silkworm, but pyrethroid insecticides, vavermectin and high content emamectin benzoate behaved high toxicity, so these insecticides should be far away from fish pond, bee yard and mulberry while applying.

Keywords: environmental toxicity

收稿日期 2010-11-30 修回日期 2010-12-27 网络版发布日期 2011-05-27

DOI:

基金项目:

山西省科技创新重大项目《农药水基化制剂新技术的研究创制》;山西省科技攻关项目《高效氟氯氰菊酯水乳剂制剂优化及应用技术研究》

通讯作者: 范仁俊

作者简介:

作者Email: rjfan@163.com

扩展功能

本文信息

- ▶ Supporting info
- ▶ PDF(719KB)
- ▶ [HTML全文]
- ▶ 参考文献[PDF]
- ▶ 参考文献

服务与反馈

- ▶ 把本文推荐给朋友
- ▶ 加入我的书架
- ▶ 加入引用管理器
- ▶ 引用本文
- ▶ Email Alert
- ▶ 文章反馈
- ▶ 浏览反馈信息

本文关键词相关文章

- ▶ 环境毒性

本文作者相关文章

- ▶ 高越
- ▶ 张润祥
- ▶ 王振
- ▶ 封云涛
- ▶ 范仁俊

PubMed

- ▶ Article by Gao,h
- ▶ Article by Zhang,R.X
- ▶ Article by Yu,z
- ▶ Article by Feng,Y.S
- ▶ Article by Fan,R.J

参考文献:

- [1]华纯.拟除虫菊酯类农药的进展和剂型[J].世界农药,2009,(5):39-44 [2]马志卿,陈安良,冯俊涛,李广泽,张兴.阿维菌素和氟虫腈对粘虫的毒杀作用比较[J].农药学报,2003,5(4):37-41 [3]杨向黎;陈春丽;李杰;王鹏.2.5%高效氟氯氰菊酯乳油防治棉铃虫药效试验总结[J].山东省农业管理干部学院学报,2005,23(5):175-176 [4]Mandal K, Chahil GS, Sahoo SK, Battu RS, Singh B.Dissipation kinetics of beta-cyfluthrin and imidacloprid in brinjal and soil under subtropical conditions of Punjab, India[J].Bull Environ Contam Toxicol,2010,84(2):225-229 [5]Mukherjee I, Gopal M, Mathur DS.Behavior of beta-cyfluthrin after foliar application on chickpea (*cicer aretinium* L) and pigeon pea (*cajanus cajan* L.)[J]. Bull Environ Contam Toxicol,2007,78(1):85-89 [6]毕富春,徐风波.甲氨基阿维菌素苯甲酸盐研究概述[J].农药科学与管理,2002,23(3):31-33 [7]王亚廷,刘亚敏,李波,王凤芝,邹向东.4.3%高氯?甲维盐微乳剂的研制[J].现代农药,2007,6(4):21-23 [8]魏书娟,宗建平,王海迎,罗万春.啮虫脘不同施药方法在甘蓝中的残留及其对蚜虫的防效[J].农药研究与应用,2009,13(2):25-28 [9]蒋迎,陈刘汪.杀虫单在水稻及稻田中的残留动态研究[J].浙江农业科学,2009,(2):410-414 [10]国家环境保护局.化学农药环境安全评价试验准则[J].,1989,:- [11]张午中,白锡川,吴明亮.阿维菌素对家蚕毒性试验[J].江苏蚕业,2007,(3):54-56 [12]吴声敢,王强,赵学平,吴长兴,陈丽萍,徐浩,章虎.阿维菌素和氟虫腈对家蚕毒性与安全性评价研究[J].浙江农业学报,2004,16(5):309-312 [13]王冬兰,刘贤进,张存政,余向阳,凌良建.阿维菌素和吡虫啉对意大利蜂的联合毒性作用[J].江苏农业科学,2006,(1):146-148 [14]魏方林,朱金文,李少南,朱国念.甲氨基阿维菌素苯甲酸盐乳油对环境生物的急性毒性研究[J].农药科学与管理,2008,29(3):19-24 [15]苍涛,赵学平,吴长兴,吴声敢,陈丽萍,王强,张志恒.甲氨基阿维菌素苯甲酸盐对4种非靶生物毒性及安全性评价[J].农药,2007,46(7):481-483

本刊中的类似文章