

[本期目录](#) | [下期目录](#) | [过刊浏览](#) | [高级检索](#)

[[打印本页](#)] [[关闭](#)]

园艺—研究进展

豆类芽苗菜生产工艺的研究进展

李振华¹,段玉¹,康玉凡²

1. 内蒙古农牧业科学院植物营养与分析研究所

2. 中国农业大学

摘要:

豆类芽苗菜指豆类种子经过萌发之后,短时间内生长为豆芽或者食用幼苗的一类特色蔬菜。与大宗蔬菜相比,豆类芽苗菜风味独特、口感脆嫩、生育周期短、对光照要求低,是一类经济价值较高的蔬菜。国内,豆类芽苗菜的基本生产技术已经初步形成,但芽苗用品种资源的筛选、浸种技术、栽培管理等研究进展缓慢。笔者从豆类芽苗菜种质资源研究与利用现状、高产优质生产工艺2个层面进行综述,供中国豆类芽苗菜科研工作者参考,并希望为豆类芽苗菜生产者提供一些理论依据和技术支持。

关键词: 栽培技术

Research progresses about production process of Bean sprout

Abstract:

Bean sprout is an edible vegetable, which comes from legume seed after a period of germination. Compared with common vegetables, bean sprouts is a kind of vegetables of high economic value which tastes unique, crisp, and its birth cycle is short, required of low illumination. In domestic, the production technology of legumes sprouts has been formed initially, but the development of germplasm resources for sprouts production, production technology and cultivation is backward. This paper was viewed from the current situation of the research and utilization of sprouts beans germplasm and from the process of high-yield, high-quality production. Hope it was helpful for researchers in Chinese beans sprouts and could provide some theoretical and technical support for bean sprouts producers.

Keywords: cultivation techniques

收稿日期 2010-11-04 修回日期 2010-12-30 网络版发布日期 2011-05-06

DOI:

基金项目:

国家食用豆产业技术体系建设专项资金资助

通讯作者: 李振华

作者简介:

作者Email: lixing_19841014@126.com

参考文献:

- [1] 王德槟,张德纯.芽苗菜蔬菜生产技术图册.北京:中国农业出版社,1997:1-15.
- [2] Rozan P, Y H Kuo, F Lambein. Nonprotein amino acids in edible lentil and garden pea seedlings. *Phytochemistry*, 2001,58: 281-289.
- [3] 张余洋,胡全凌,李汉霞.不同处理对豌豆和萝卜芽苗菜生长、产量及品质的影响.华中农业大学学报,2008,2(27):289-293.
- [4] 肖伶俐,康玉凡,陶礼明,等.不同大豆品种芽用特性比较.大豆科学,2008,27(6):23-26.
- [5] 罗珊,康玉凡,濮绍京,等.黑河地区55份大豆品种资源农艺性状和营养成分的聚类分析.大豆科学,2009,28(3):421-425.
- [6] 程须珍,王素华.不同绿豆品种豆芽产量与质量的差异分析.中国绿豆科技运用论文集.北京:中国农业出版社,1995:144-148..

扩展功能

本文信息

Supporting info

[PDF \(546KB\)](#)

[\[HTML全文\]](#)

[参考文献\[PDF\]](#)

[参考文献](#)

服务与反馈

[把本文推荐给朋友](#)

[加入我的书架](#)

[加入引用管理器](#)

[引用本文](#)

[Email Alert](#)

[文章反馈](#)

[浏览反馈信息](#)

本文关键词相关文章

栽培技术

本文作者相关文章

[李振华](#)

[段玉](#)

[康玉凡](#)

PubMed

[Article by Li,Z.H](#)

[Article by Duan,y](#)

[Article by Kang,Y.F](#)

- [7] 李振华,康玉凡,程须珍,等.绿豆品种芽用特性的初步评价.中国农业大学学报,2010,15(5):31-36
- [8] 李振华.绿豆品种资源芽用特性的综合评价.中国农业大学图书馆:中国农业大学硕士学位论文,2010
- [9] 李振华,康玉凡,濮绍京,等.不同地区绿豆品种芽用特性的研究.中国农学通报,2010,15:361-364.
- [10] Kwon S H, Lee Y I, Kim J R. Evaluation of important sprouting characteristics of edible soybean sprout cultivars. Korean J Breed, 1981, 13:202-206
- [11] Lee S C, Choi G G, Chang Y N . Studies on the performance characteristics of collected native sprout soybean. I. Variation of characteristics in collected native sprout soybean. Korean J Breed, 1992, 24:87-95
- [12] Park M H, Kim D C, Kim B S . Studies on pollution-free soybean sprout production and circulation market improvement. Korea Soybean Dig, 1995, 12: 51 - 67
- [13] 花登峰,赵团结,张黎萍.小粒专用大豆品种遗传改良研究进展.杂粮作物,2005,25(5):311-313.
- [14] 吴叶青.播种密度对豌豆芽生产的影响.长江蔬菜,2006,02:51-52.
- [15] Lee J D, Hwang Y H, Cho H Y. Comparison of characteristics related with soy bean sprouts between Glycine max and G soja. Korean Journal of Crop Science, 2002, 47(3): 189-195.
- [16] Lee S H, K Y Park, H S Lee. Genetic mapping of QTLs conditioning soybean sprout yield and quality. Theoretical and Applied Genetics, 2001, 103(5): 702-709.
- [17] 张雷闰,洪睿鹿,文成梁.黑河17号的特征特性及高产栽培技术.耕作与栽培,2005,2:61.
- [18] 马启慧.2000年黑龙江省审定推广的大豆品种.大豆科学,2000,19(2): 193-196.
- [19] 王秋玲,谷传彦,郭凌云,等.黄淮夏大豆主要农艺性状与单株产量关系分析.山东农业科学,1999,3: 26-28.
- [20] 梁燕,杨又迪.大粒绿豆农艺性状与产量的相关关系.西北农业学报, 1998, 7(3): 98-99.
- [21] 韩粉霞,李桂英.绿豆主要农艺性状的相关关系.华北农学报, 1998, 13(4): 66-69.
- [22] Danisova,C., Holotnakova,E., Hozova,B., Buchtova,V. Effect of germination on a range of nutrients of selected grains and legumes. Acta Alimentaria, 1995, 23 (3):287-298.
- [23] Urbano,G., Lopez-Jurado,M., Hernandez,J., Fernandez,M., Moren,M.C., Frias,J., Diaz-Pollan,C., Prodanov,M., Vidal-Valverde,C. Nutritional assessment of raw, heated and germination lentils Journal of Agricultural and Food Chemistry, 1995, 43:1871-877.
- [24] Ayet, G., Burbano, C., Cuadrado, C., Pedrosa, M.M., Robredo, L.M., Muzquiz, M., Cuadra, C., Castaño, A., Osagis, A. Effect of germination, under different environmental conditions, on saponins, physic acid and tannins in lentils (*Lens culinaris*). Journal of the Science of Food and Agriculture, 1997, 74 (2):273-278.
- [25] Vidal-Valverde, C., Frías, J., Sierra, I., Blazquez, I., Lambien, F., Kuo,Y-H.. New functional legume food by germination. Effect on the nutritive value of beans, lentils and peas. European Food Research and Technology, 2002, 215: 472-476.
- [26] Vidal-Valverde, C., Frías, J., Hernández, A., Martín-Alvarez, P.J., Sierra, I., Rodríguez, C., Blazquez, I., Vicente, G. Assessment of nutritive compounds and ant nutritive factors in pea (*Pisum sativum*) seeds. Journal of the Science of Food and Agriculture, 2003, 83: 1-4.
- [27] López-Amorós, M.L., Hernández, T., Estrella, I. 2001. The germination process in relation to the nutritional value of legumes. In: Amadó, R., Aby, B., Bravo, L., Goñi, I., Saura-Calixto, F. (Eds.), Bioactive Compounds in Plant Foods. Health Effects and Perspectives for the Food Industry. Cost 916. European Commission, Luxembourg, pp. 173-176.
- [28] Frias, J., Fernández-Orozco, R., Zielinski, H., Piskula, M., Kozlowska, H., Vidal-Valverde, C. Effect of germination on the content of vitamin C and E of lentils. Polish Journal of Food and Nutrition Science ,2002, 52(11): 76.
- [29] Bau H M, Villaume C, Nicolas J P. Effect of germination on chemical composition, biochemical constituents and anti-nutritional factors of Soya bean (Glycine) seeds. Journal of the Science of Food and Agriculture, 1997, 73:1-9.
- [30] Rozan P,Kuo Y H, Lambein F. Free amino acids present in commercially available seedlings sold for human consumption. A potential hazard for consumers. Journal of agricultural and food chemistry ,2000, 48(3): 716-23.
- [31] Whalley W, Bengough A, Dexter A. Water stress induced by PEG decreases the maximum growth pressure of the roots of pea seedlings. Journal of Experimental Botany, 1998, 49:1689-1694.
- [32] 张永清,顾振新,张颖,邢邯.豆芽生产中大豆浸泡条件与吸水率和发芽率的关系研究.食品研究与开发,2007,28 (1):26-29.
- [33] 吴小勇,曾庆孝,田金河,柳明鲜.绿豆的浸泡工艺及其对绿豆种子萌发的影响研究.工艺技术,2004 (4) : 107-109.
- [34] 许彬,张应华,范眸天.不同处理对豌豆芽苗菜生长和产量的影响.云南农业大学学报,2004,19(5):613-615.
- [35] 刘福霞,刘乃森,何莉,孙长利.浸种时间对黑豆芽苗菜产量及蛋白质和Vc含量的影响.安徽农业科学,2007,35 (31):9855-9860.
- [36] 曹健,李植良,黄河勋,石尧清,谭青,赖穗春,何焕清.浸种时间对小豆种子萌发及生长的影响.广东农业科学,1999(3):25.
- [37] K A Taiwo. Regression Relationships for the Soaking and Cooking Properties of Two Cowpea Varieties. Food Engineering, 1998, 33 (37):1-344.
- [38] Li Z H, Liu Z J. Effects of Benzyl adenine and Naphthalene Acetic Acid on Growth and Camptothecin accumulation in Camptotheca Acuminata Seedlings J Plant Growth Regul, 2003, 22: 205-216.

- [39] Karin Kollàrovà, Mária Henselovà, Desana Li?ková. Effect of auxins and plant oligosaccharides on root formation and Elongation growth of mung bean hypocotyls. Plant Growth Regulation, 2005, 46: 1-9.
- [40] Polanco M C, Peláez M I, Ruiz M L. Factors that affect plant regeneration from in vitro culture of immature seeds in four lentil cultivars. Plant Cell Tissue Organ Cult 1988, 15: 175-182.
- [41] Jin M C, Ren Y P, Chen X H. Determination of 6-Benzyladenine in Bean Sprout by LC - ESI-IT-MS - MS . Chromatographia, 2007, 66: 407 - 410.
- [43] Ryu Yeong Seop, Hong Dong. Effects of watering method and BA concentration on growth and Morphological characteristics of mungbean sporouts. Korean Journal of Crop science, 2006, 51(3): 204-208.
- [44] Li Q , Zhou Q , Zhao X R . Application of chitosan and chitosan derivatives into processing of mung bean sprout. Journal of Anhui Agricultural University, Hefei, China: 2005. 32(3), 402-405.
- [45] Ro J S, Chung I M. Method for manufacturing green elemental bean sprouts using yellow light (2005). Patent No. 428997, Korea.
- [46] 吴正景, 郭大龙, 王少先, 高文, 汪伦记. 有色薄膜覆盖对豌豆芽苗生长及富硒的影响. 河南科技大学学报, 2008, 29(5): 70-72.
- [47] Hwang Y H, Lee J D, Cho H Y . Sprout characteristics of improved and indigenous soybeans in Korea. Agric Res Bull Kyungpook Natl Univ, 2002, 20: 99 - 105.
- [48] Yoon D J, Lee J D, Kang D J. Effect of electroglyzed acidic water on the growth of soybean sprout. J Life Sci, 2004, 14: 809 - 814.
- [49] Lee J D , Shannon J G , Jeong Y S. Young-Hyun Hwang. A simple method for evaluation of sprout characters in soybean. Euphytica , 2007, 153: 171 - 180.
- [50] 张丽, 吴小刚, 张力群, 等. 豆芽烂芽的病原菌分离鉴定及致病性研究. 长江蔬菜, 2010, 2: 71-74.

本刊中的类似文章

- 孙其宝, 徐义流, 俞飞飞, 孙俊, 陆丽娟. 葡萄优质高效避雨栽培技术研究[J]. 中国农学通报, 2006, 22(11): 477-477
- 张绪璋. 北虫草新菌株选育及栽培技术研究[J]. 中国农学通报, 2010, 26(19): 206-210
- 付文词, 张京顺, 马宗荣, 袁珺. 无公害石竹茶的人工栽培与制作技术[J]. 中国农学通报, 2004, 20(3): 67-67
- 阮培均, 马俊. 优质高产杂交玉米毕单4号栽培技术模式研究[J]. 中国农学通报, 2003, 19(4): 78-78
- 郝继伟, 王连翠. 爆裂玉米的经济价值与栽培技术[J]. 中国农学通报, 2003, 19(6): 97-97
- 杨海棠, 王伟, 马东波. 中国北方地区花生栽培技术的研究进展[J]. 中国农学通报, 2004, 20(4): 169-169
- 刘伟明. 两优培九在浙江台州作单季晚稻栽培的特征特性及其栽培技术研究[J]. 中国农学通报, 2005, 21(8): 162-162
- 马聪, 吴振民, 姚延宾, 李光奇, 马京民. 气候过渡型烟区优质烤烟标准化生产关键技术*[J]. 中国农学通报, 2004, 20(3): 258-258
- 王嘉祥. 薄荷间作高产高效栽培技术[J]. 中国农学通报, 2004, 20(3): 204-204
- 王俊峰. Research and Application on Combination of Comprehensive Agronomic Measures in High-yielding Cotton Cultivation[J]. 中国农学通报, 2005, 21(4): 156-156
- zouyingbin@.net. 超级稻“三定”栽培法研究 I 概念与理论依据[J]. 中国农学通报, 2006, 22(5): 158-158
- 马春红, 陈霞, 翟彩霞, 郭秀林, 崔四平, 李广敏. 薄荷的快速繁殖与栽培技术的研究[J]. 中国农学通报, 2004, 20(4): 230-230
- 周秋峰, 黄长志, 王保林, 赵建国. 栽培技术对小麦品质的调节效应[J]. 中国农学通报, 2009, 25(08): 137-141
- 卢春生. 加工咸酥花生专用型品种“台南选9号”高产栽培技术研究[J]. 中国农学通报, 2005, 21(11): 140-140
- 赖永红. 不同栽培技术措施对福建龙岩红芽芋产量相关性状的影响[J]. 中国农学通报, 2009, 25(08): 169-173