



首页 期刊介绍 编委会 编辑部 投稿须知 期刊订阅 广告服务 联系我们 留

动物营养学报 » 2012, Vol. 24 » Issue (8) :1397-1403 DOI: 10.3969/j.issn.1006-267x.2012.08.002

综述

[最新目录](#) | [下期目录](#) | [过刊浏览](#) | [高级检索](#)

[<< Previous Articles](#) | [Next Article >>](#)

微生物饲料添加剂的主要功能及其研究进展

徐鹏, 董晓芳, 佟建明

中国农业科学院北京畜牧兽医研究所, 北京 100193

Microbial Feed Additives: Major Functions and Its Research Advances

XU Peng, DONG Xiaofang, TONG Jianming

Institute of Animal Sciences, Chinese Academy of Agricultural Sciences, Beijing 100193, China

- 摘要
- 参考文献
- 相关文章

Download: PDF (967KB) [HTML](#) (1KB) Export: BibTeX or EndNote (RIS) Supporting Info

摘要 益生菌是当摄入量足够时能对机体产生有益作用的活性微生物。人们常把应用于畜牧业生产上的益生菌称为微生物饲料添加剂。以往研究表明,微生物饲料添加剂具有维护动物肠道健康、缓解不良应激、改善畜舍环境、调节机体脂肪代谢和改善畜产品品质的功能。还有研究者认为,微生物饲料添加剂具有替代抗生素功能的作用。本文旨在就微生物饲料添加剂的主要功能及其研究进展进行综述,为今后在畜牧业生产上的科学应用及相关研究提供参考。

关键词: [益生菌](#) [微生物饲料添加剂](#) [功能](#)

Abstract: Probiotics are live microorganisms which when administered in adequate amounts confer a health benefit on the host. The probiotics used in livestock production are usually called 'microbial feed additives'. A large quantity of previous studies showed that microbial feed additives had many desirable functions, including maintaining intestinal health, relieving unpleasant stress, ameliorating animal housing environment and improving the quality of animal by-products. And some articles also indicated that microbial feed additives had a potential ability of replacing antibiotics. The present paper reviewed the recent advances about major functions of microbial feed additives.

Keywords: [probiotics](#), [microbial feed additives](#), [functions](#)

收稿日期: 2012-03-10;

基金资助:国家蛋鸡产业技术体系建设专项经费(CARS-41-K16);"十一五"科技支撑计划项目课题(2006BAD12B05)

通讯作者 董晓芳,副研究员,硕士生导师,E-mail: xiaofangd1124@sina.com

引用本文:

徐鹏, 董晓芳, 佟建明 . 微生物饲料添加剂的主要功能及其研究进展[J]. 动物营养学报, 2012,V24(8): 1397-1403

XU Peng, DONG Xiaofang, TONG Jianming . Microbial Feed Additives: Major Functions and Its Research Advances[J]. Chinese Journal of Animal Nutrition, 2012,V24(8): 1397-1403.

链接本文:

http://118.145.16.228/Jweb_dwyy/CN/10.3969/j.issn.1006-267x.2012.08.002 或 http://118.145.16.228/Jweb_dwyy/CN/Y2012/V24/I8/

Service

- ▶ 把本文推荐给朋友
- ▶ 加入我的书架
- ▶ 加入引用管理器
- ▶ Email Alert
- ▶ RSS

作者相关文章

- ▶ 徐鹏
- ▶ 董晓芳
- ▶ 佟建明

- [1] FAO,WHO.Report of a joint FAO/WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria[M].Cordoba:Argentina,2001.
- [2] HANSEN I G,MØLLGAARD H.Investigations of the effect of lactic acid on the metabolism of calcium and phosphorus[J].Acta Physiologica Scandinavica,1947,14(1/2): 158-170.
- [3] PARKER R B.Probiotics,the other half of the antibiotic story[J].Animal Nutrition and Health,1974,29(2): 4-8.

- [4] FULLER R.History and development of probiotics . :Probiotics:the scientific basis,1992:1-8.
- [5] YASON C V,SUMMERS B A,SCHAT K A.Pathogenesis of rotavirus infection in various age groups of chickens and turkeys:pathology[J].Am Journal of Veterinary Research,1987,48(6):927-938.
- [6] PAGAN J,SEERLEY B,COLE D,et al.How do mannanoligosaccharides work[J].Feeding Times,1999,4(1):7-9.
- [7] AWAD W A,BOHM J,RAZZAZI-FAZELI E,et al.Effect of addition of a probiotic microorganism to broiler diets contaminated with deoxynivalenol on performance and histological alterations of intestinal villi of broiler chickens[J].Poultry Science,2006,85(6):974-979.
- [8] BONTEMPO V,DI GIANCAMILLO A,SAVOINI G,et al.Live yeast dietary supplementation acts upon intestinal morpho-functional aspects and growth in weanling piglets[J].Animal Feed Science and Technology,2006,129(3/4):224-236.
- [9] GIANG H H,VIET T Q,OGLE B,et al.Growth performance,digestibility,gut environment and health status in weaned piglets fed a diet supplemented with a complex of lactic acid bacteria alone or in combination with *Bacillus subtilis* and *Saccharomyces boulardii*[J].Livestock Science,2012,143(2/3):132-141.
- [10] MARTEAU P R,VRESE M,CELLIER C J,et al.Protection from gastrointestinal diseases with the use of probiotics[J].The American Journal of Clinical Nutrition,2001,73(2):430-436.
- [11] CHOI J Y,SHINDE P L,INGALE S L,et al.Evaluation of multi-microbe probiotics prepared by submerged liquid or solid substrate fermentation or antibiotics in weaning pigs[J].Livestock Science,2011,138(1/2/3):144-151.
- [12] LE BON M,DAVIES H E,GLYNN C,et al.Influence of probiotics on gut health in the weaned pig[J].Livestock Science,2010,133(1/2/3):179-186.
- [13] 黄俊文,林映才,冯定远,等.益生菌、甘露寡糖对早期断奶仔猪生长、免疫和抗氧化机能的影响[J].动物营养学报,2005,17(4):16-20.
- [14] KODALI V P,SEN R.Antioxidant and free radical scavenging activities of an exopolysaccharide from a probiotic bacterium[J].Biotechnology Journal,2008,3(2):245-251. 
- [15] SUTTON A L,KEPHART K B,VERSTEGEN M W,et al.Potential for reduction of odorous compounds in swine manure through diet modification [J].Journal of Animal Science,1999,77(2):430-439.
- [16] JIMENEZ-COLMENERO F,CARBALLO J,COFRADES S.Healthier meat and meat products: their role as functional foods[J].Meat Science,2001,58(1):5-13. 
- [17] SALMA U,MIAH A G,MAKI T,et al.Effect of dietary *Rhodobacter capsulatus* on cholesterol concentration and fatty acid composition in broiler meat[J].Poultry Science,2007,86(9):1920-1926.
- [18] SALMA U,MIAH A G,TAREQ K M,et al.Effect of dietary *Rhodobacter capsulatus* on egg-yolk cholesterol and laying hen performance[J].Poultry Science,2007,86(4):714-719.
- [19] SALMA U,MIAH A G,TSUJII H,et al.Effect of dietary *Rhodobacter capsulatus* on lipid fractions and egg-yolk fatty acid composition in laying hens[J].Journal of Animal Physiology and Animal Nutrition,2011.doi: 10.1111/j.1439-0396.2011.01224.x.
- [20] LI L,XU C L,JI C,et al.Effects of a dried *Bacillus subtilis* culture on egg quality[J].Poultry Science,2006,85(2):364-368.
- [21] 康白.微生态学[M].大连:大连出版社,1988.
- [22] THRELFALL E J,WARD L R,FROST J A,et al.The emergence and spread of antibiotic resistance in food-borne bacteria[J].International Journal of Food Microbiology,2000,62(1/2):1-5.
- [23] SILBERGELD E K,GRAHAM J,PRICE L B.Industrial food animal production,antimicrobial resistance,and human health[J].Annual Review of Public Health,2008,29:151-169.
- [24] GRIGGS J P,JACOB J P.Alternatives to antibiotics for organic poultry production[J].The Journal of Applied Poultry Research,2005,14(4):755-756.
- [25] MOUNTZOURIS K C,TSIRTSIKOS P,KALAMARA E,et al.Evaluation of the efficacy of a probiotic containing *Lactobacillus*,*Bifidobacterium*,*Enterococcus*,and *Pediococcus* strains in promoting broiler performance and modulating cecal microflora composition and metabolic activities[J].Poultry Science,2007,86(2):309-317.
- [26] YEO J,KIM K I.Effect of feeding diets containing an antibiotic,a probiotic,or yucca extract on growth and intestinal urease activity in broiler chicks[J].Poultry Science,1997,76(2):381-385.
- [1] 刘立恒,闵力,瞿明仁,戈婷婷,钟志勇.甘露寡糖、果寡糖和大豆寡糖组合对锦江黄牛瘤胃液细菌多样性的影响[J].动物营养学报,2012,24(8): 1583-1588
- [2] 刘虎传,张敏红,李素霞,冯京海,姜海龙,杨家军,殷瑞娟.益生菌制剂对早期断奶仔猪肠道pH、黏膜形态结构和挥发性脂肪酸含量的影响 [J]. 动物营养学报, 2012,24(7): 1329-1335
- [3] 田金可, Ahmad Hussain, 李伟, 高尚, 王恬.不同硒源及水平对肉鸡组织硒含量及抗氧化功能的影响[J]. 动物营养学报, 2012,24(6): 1030-1037
刘虎传, 张敏红, 冯京海, 姜海龙, 杨晓岚, 姜礼文, 陈燕.益生菌制剂对早期断奶仔猪生长性能和免疫指标的影响[J]. 动物营养学报, 2012,24(6): 1124