

[本期目录](#) | [下期目录](#) | [过刊浏览](#) | [高级检索](#)[\[打印本页\]](#) [\[关闭\]](#)**兽医—研究报告****油樟叶挥发油的皮肤和眼睛毒理学研究**张笑乐¹,向楠¹,殷中琼¹,尚永祥²1. 四川农业大学动物医学院
2. 四川农业大学生命科学与理学院**摘要:**

目的 研究油樟叶挥发油对皮肤和眼睛的刺激性及其光毒性。方法 通过一次给药和多次给药皮肤刺激试验、一次给药和多次给药的眼睛刺激试验和皮肤光毒性试验，观察皮肤反应和眼睛反应，评价其皮肤刺激性、眼睛刺激性和光毒性。结果 一次和多次用药高、中剂量油樟叶挥发油对破损皮肤有可逆性刺激，低剂量无刺激；三个剂量对完整皮肤均无刺激。一次用药对眼睛有轻微可逆刺激，多次用药刺激个别个体不可逆。油樟叶挥发油无光毒性。结论 中剂量以下油樟叶挥发油皮肤用药较为安全，眼睛用药要慎重。

关键词： 光毒性

Research on Dermal Toxicity and Eye Toxicity of the Volatile Oil from the Leaf of Cinnamomum Longepaniculatum

Abstract:

Objective To evaluate the skin irritation, the eye irritation and the phototoxicity of the volatile oil from the leaf of *Cinnamomum Longepaniculatum*. **Method** Evaluate the security of the volatile oil from the leaf of *Cinnamomum Longepaniculatum* by carrying on the skin irritation, the eye irritation test and the phototoxicity test and observing the response. **Result** The high and the middle concentrations had slight and reversible irritation to the damaged skin, but the low concentration didn't. Those three all had no irritation to the intact skin. When the volatile oil from the leaf of *Cinnamomum Longepaniculatum* was given to the eyes once, it had slight and reversible irritation, yet it became irreversible when given several times to a few rabbits. The volatile oil from the leaf of *Cinnamomum Longepaniculatum* had no phototoxicity. **Conclusion** As a whole, it's safe to apply the volatile oil from the leaf of *Cinnamomum Longepaniculatum* to the skin, while the application to the eyes must be careful.

Keywords: phototoxicity**收稿日期** 2011-04-08 **修回日期** 2011-04-29 **网络版发布日期** 2011-09-21**DOI:****基金项目:**

教育部《长江学者和创新团队发展计划》创新团队项目

通讯作者: 殷中琼**作者简介:**

作者Email: yinzhongq@163.com

参考文献:

- [1]中国科学院中国植物志编辑委员会.中国植物志第三卷[J].北京科学出版社,1982,:184-186
- [2]罗中杰,李维一,魏琴,等.宜宾油樟的现状及未来[J].四川师范大学学报(自然科学版),2001,24(3):317-317
- [3] Joshi SC, Padalia RC, Bisht DS,等.Terpenoid diversity in the leaf essential oils of Himalayan Lauraceae[J].speciesChem Biodivers,2009,6(9):1364-1373
- [4]陈寿仁.樟科药用植物药理研究及开发[J].中国中医药信息杂志,1996. 3 (2) : 13- 15.,1996,3(2):13-15
- [5]王涛,游玲,魏琴,等.油樟内生细菌的多样性及抑制植物病原菌初步研究[J].西北林学院学报,2009,

扩展功能
本文信息
Supporting info
PDF(1570KB)
[HTML全文]
参考文献[PDF]
参考文献
服务与反馈
把本文推荐给朋友
加入我的书架
加入引用管理器
引用本文
Email Alert
文章反馈
浏览反馈信息
本文关键词相关文章
光毒性
本文作者相关文章
张笑乐
向楠
殷中琼
尚永祥
PubMed
Article by Zhang,X.L
Article by Xiang,n
Article by Yin,Z.Q
Article by Chang,Y.X

[6]Yeh RY, Shiu YL, Shei SC, 等. Evaluation of the antibacterial activity of leaf and twig extracts of stout camphor tree, *Cinnamomum kanehirae*, and the effects on immunity and disease resistance of white shrimp, *Litopenaeus vannamei*[J].*Fish Shellfish Immunol*,2009,27(1):26-32

[7]Soon-II Kim, Young-Eun Na, Ji-Hwan Yi, 等.Contact and fumigant toxicity of oriental medicinal plant extracts against *Dermanyssus gallinace* (Acari: Dermyssidae)[J].*Veterinary Parasitology*,2007,145: 377-382

[8]Gilbert S. Bunker, Christopher.T. Rhodes. Modern Pharmaceutics, Fourth edition [J].NewYork: Informa Healthcare,2002,: 266-274

[9]楼宜嘉.药品毒理学[J].北京: 人民卫生出版社,2007,: 192-204

[10]中华人民共和国卫生部.化学品毒性鉴定技术规范[J].北京: 卫生部法制与监督司编印,2005,: 26-36

[11]杨美玲, 张紫虹, 胡永成, 等.一种聚维酮碘消毒液的毒理学安全性评价[J].中国卫生检验杂志,2007,17

(7):1286-1288

[12] 龙正海, 杨再昌, 杨雄志. 油茶树嫩枝挥发油皮肤毒理及其促透作用研究[J]. 中国中药杂志, 2007. 32 (17) : 1780- 178

[13]黄正, 杜之渝, 黄易.汉防己甲素对兔眼毒性作用的初步研究[J].眼科新进展,2008,28(3):195-198

[14]张丽霞, 宋丽华, 史勤俭,等.光毒性试验方法研究[J].环境与健康杂志,2002,19: 60-61

本刊中的类似文章