🔁 Current Issue 🥖 Browse Issues	Acta Medica Iranica 2009;47(4) : 179-185	
Search	"11Hydroxylas deficient form of congenital adrenal hyperplasia with bilateral testicular tumors and true precocious puberty (A case report) "	
🦪 About this Journal	"Moayeri H "	
🔏 Instruction to Authors	Abstract:	
 Online Submission Subscription Contact Us 	The development of testicular masses in male patients with congenital adrenal hyperplasia due to 21 hydroxylase deficiency has been recognized for many years. We present here the eighth and ninth reported patient with bilateral testicular tumors associated with 11 hydroxylase deficiency. They were two brothers aged 7.5 and 5 yr. who had bilateral testicular tumors and diagnosed because of signs and symptoms of true precocious puberty and high blood pressure. The patients testicular enlargement is suggestive of bilateral testicular adrenal rest.	
S Feed	Keywords: Testicular tumor ، Adrenal hyperplasia ، 11 Hydroxylase deficiency	
	TUMS ID: 1381	
	Full Text HTML 🔊 Full Text PDF 🖄 1477 KB	,

Home - About - Contact Us

TUMS E. Journals 2004-2009 Central Library & Documents Center Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024*768 Resolutions