

[Available Issues](#) | [Japanese](#)Author:  [ADVANCED](#) | Volume  Page Keyword:    [TOP](#) > [Available Issues](#) > [Table of Contents](#) > **Abstract****Tropical Medicine and Health**

Vol. 36 (2008) , No. 4 p.181

**Disuse atrophy of small muscles due to effusion of the synovial compartments of the dorsal wrist in CHIKV infection**[Lourduraj John De Britto](#)<sup>1)</sup>, [Yuvaraj Jayaraman](#)<sup>1)</sup> and [Lalitha](#)

1) Division of Clinical Epidemiology &amp; Chemotherapy Vector Control Center, Tamil Nadu Veterinary, Animal and Fisheries Sciences University, Chennai, India

(Accepted August 8, 2008)

**Abstract:** Inflammation of the joints is generally considered to be a common and prolonged symptoms in CHIKV infection. Disuse atrophy of the small muscles and effusion of the synovial compartments of the dorsal wrist in CHIKV infection has not been reported so far. Studying these atypical cases may shed light on the pathogenesis of CHIKV infection and appropriate individual case management.

**Key words:** [Night Blood Survey](#), [W.bancrofti](#), [filariasis](#), [hydrocele](#), [Dance Sign](#)

To cite this article:

Lourduraj John De Britto, Yuvaraj Jayaraman and Lalit Kumar Da  
muscles due to effusion of the synovial compartments of the dorsal  
A case report”. Tropical Medicine and Health, Vol. **36**, pp.181-18

---

doi:10.2149/tmh.2008-14

JOI JST.JSTAGE/tmh/2008-14

Copyright (c) 2008 by The Japanese Society of Tropic

---