



 **Current Issue**

 **Browse Issues**

 **Search**



 **About this Journal**

 **Instruction to Authors**

 **Online Submission**

 **Subscription**

 **Contact Us**



 **RSS Feed**

## Acta Medica Iranica

2009;47(4) : 183-186

### Low Magnesium Concentration in Erythrocytes of Children with Acute Asthma

Mostafa Sedighi, Zahra Pourpak, Behrouz Bavarian, Reza Safar Alizadeh, Ahad Zare, Mostafa Moin

#### Abstract:

Magnesium (Mg) is the second most abundant intracellular cation and is involved in numerous physiological functions, including protein folding, intracellular signaling and enzyme catalysis. It has been shown that magnesium deficiency exacerbates pulmonary airways hyperreactivity. Several studies suggest that magnesium level has no effect on asthma but others had shown a contributory effect. Because of its intracellular abundance, the aim of this study was to determine if there was any difference in plasma and intracellular Mg concentrations of children with acute asthma compared to non asthmatic children. Twenty nine patients with acute asthma aged 2 to11 years admitted to the emergency department of hospital and 37 non asthmatic children with the same age were included in our study. 0.5 mL of heparinized whole blood samples of patients who were meeting inclusion criteria at the onset of admission with bronchoconstriction and before using any medication was drawn and it was immediately sent to the laboratory. Plasma and erythrocytes were separated and stored at  $-20^{\circ}\text{C}$  and later their Mg levels were quantified with atomic absorption spectrophotometry method. The average plasma and intracellular magnesium levels in patients were  $(0.79 \pm 0.098 \text{ mmol/L})$  and  $(1.17 \pm 0.27 \text{ mmol/L})$  respectively. Results of 37 non asthmatic persons [plasma  $(0.85 \pm 0.1 \text{ mmol/L})$  and erythrocytes  $(1.33 \pm 0.21 \text{ mmol/L})$ ] showed that there was no significant difference between plasma Mg levels in two groups ( $p 0.06$ ) but intracellular magnesium level was significantly lower ( $p 0.03$ ) in patients group. These results indicate that intracellular Mg level may be a more accurate method to assess Mg level in patients with asthma. Hence, determination of Mg concentration in erythrocytes may be used in evaluation of asthma pathophysiology. There are recommendations for using intravenous Mg sulfate in acute asthma, and this study supports the rational for using it in emergency departments for acute severe asthma.

#### Keywords:

Children Magnesium

TUMS ID: 2921

Full Text HTML  Full Text PDF  188 kB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009  
Central Library & Documents Center  
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024\*768 Resolutions