

缺血性脑损伤治疗新策略:促进血管新生和神经再生

立即下载

New strategies for ischemic cerebral injury therapy: promoting angiogenesis and neurogenesis

丁利静;石京山;李菲;黄彬;杨丹莉;龚其海;

摘要:

缺血性脑损伤病因及发病机制仍不完全清楚。目前促进血管新生和神经再生在其发生和发展中的作用日益受到关注。本文综述了近年来关于促进血管新生和神经再生的研究进展,旨在为将来研发缺血性脑损伤治疗药物提供新策略。

关键词: 脑缺血;;血管生成调节剂;;神经再生

基金项目: 国家自然科学基金(81060349); 贵州省科技厅联合资金(黔科合J字LKZ[2011]27号); 教育部新世纪优秀人才支持计划(NCET-11-0927); 教育部创新团队(IRT1197)

通讯作者: 丁利静;

Email:

参考文献:

- [1]ARAI K,JIN G,NAVARATNA D,et al.Brain angiogenesis indevelopmental and pathological processes:neurovascular injuryand angiogenic recovery after stroke[J].FEBS J,2009,276(17):4644-4652.
- [2]ADAMS HP Jr,del ZOPPO G,ALBERTS MJ,et al.Guidelinesfor the early management of adults with ischemic stroke[J].Circulation,2007,115(20):e478-e534.
- [3]LOK J,GUPTA P,GUO S,et al.Cell-cell signaling in the neurovascular unit[J].Neurochem Res,2007,32(12):2032-2045.
- [4]高梅,刘睿,杜冠华.缺血性中风治疗新策略——靶向神经血管单元[J].中国临床药理学与治疗学,2008,13(7):813-821.
- [5]FONT MA,ARBOIX A,KRUPINSKI J,et al.Angiogenesis,neurogenesis and neuroplasticity in ischemic stroke[J].Curr CardiolRev,2010,6(3):238-244.
- [6]ZHANG RL,ZHANG ZG,CHOPP M,et al.Ischemic strokeand neurogenesis in the subventricular zone[J].Neuropharmacology,2008,55(3):345-352.
- [7]ZHANG RL,ZHANG ZG,ZHANG L,et al.Proliferation and differentiation of progenitor cells in the cortex and the subventricular zone in the adult rat after focal cerebral ischemia[J].Neuroscience,2001,105(1):33-41.
- [8]LIU XS,ZHANG ZG,ZHANG RL,et al.Stroke induces gene profile changes associated with neurogenesis and angiogenesis in adult subventricular zone progenitor cells[J].J Cereb Blood

扩展功能

● 本文信息

[PDF\(332K\)](#)[参考文献](#)

● 服务与反馈

[引用本文](#)

● 本文关键词相关文章

[脑缺血](#)[血管生成调节](#)[剂](#)[神经再生](#)

● 本文作者相关文章

[丁利静](#)[石京山](#)[李菲](#)[黄彬](#)[杨丹莉](#)[龚其海](#)

● 中国知网

[丁利静](#)

FlowMetab,2007,27(3):564-574.

[9]LING L,HOU Q,XING S,et al.Exogenous kallikrein enhances neurogenesis and angiogenesis in the subventricular zone and the peri-infarction region and improves neurological function after focal cortical infarction in hypertensive rats[J].Brain Res,2008,1206:89-97.

[10]BROWN JP,COUILLARD-DESPRÉS SS,COOPER-KUHN CM,et al.Transient expression of doublecortin during adult neurogenesis[J].J Comp Neurol,2003,467(1):1-10.

[11]ZHANG RL,ZHANG ZG,LU M,et al.Reduction of the cell cycle length by decreasing G(1) phase and cell cycle reentry expand neuronal progenitor cells in the subventricular zone of adult rat after stroke[J].J Cereb Blood Flow Metab,2006,26(6):857-863.

[12]NAGAO M,SUGIMORI M,NAKAFUKU M.Cross talk between notch and growth factor/cytokine signaling pathways in neural stem cells[J].Mol Cell Biol,2007,27(11):3982-3994.

[13]ZHANG RL,ZHANG ZG,CHOPP M.Ischemic stroke and neurogenesis in the subventricular zone[J].Neuropharmacology,2008,55(3):345-352.

[14]LEVENTHAL C,RAFII S,SHAHAR A,et al.Endothelial trophic support of neuronal production and recruitment from the adult mammalian subependyma[J].Mol Cell Neurosci,1999,13(6):450-464.

[15]OHAB JJ,FLEMING S,BLESCH A,et al.A neurovascular niche for neurogenesis after stroke[J].J Neurosci,2006,26(50):13007-13016.

[16]BENRAISS A,CHMIELNICKI E,LERNER K,et al.Adenoviral brain-derived neurotrophic factor induces both neostriatal and olfactory neuronal recruitment from endogenous progenitor cells in the adult forebrain[J].J Neurosci,2001,21(17):6718-6731.

[17]ARVIDSSON A,COLLIN T,KIRIK D,et al.Neuronal replacement from endogenous precursors in the adult brain after stroke[J].Nat Med,2002,8(9):963-970.

[18]李欢欢,何旭,刘克丁.内皮祖细胞与缺血性脑卒中[J].中国组织工程研究与临床康复,2010,14(19):3569-3572.

[19]TAGUCHI A,SOMA T,TANAKA H,et al.Administration of CD34+ cells after stroke enhances neurogenesis via angiogenesis in a mouse model[J].J Clin Invest,2012,122(3):330-338.

[20]NOCHI R,KATO T,KANEKO J,et al.Involvement of metabotropic glutamate receptor 5 signaling in activity-related proliferation of adult hippocampal neural stem cells[J].Eur J Neurosci,2012,36(3):2273-2283.

[21]INOUE T,SUGIYAMA M,HATTORI H,et al.Stem cells from human exfoliated deciduous tooth-derived conditioned medium enhance recovery of focal cerebral ischemia in rats[J].Tissue Eng Part A,2013,19(1-2):24-29.

[22]ZHANG RL,ZHANG ZG,ZHANG L,et al.Delayed treatment with sildenafil enhances neurogenesis and improves functional recovery in aged rats after focal cerebral ischemia[J].J Neurosci Res,2006,83(7):1213-1219.

[23]HICKEL M,SCHLENGER K,DOCTROW S,et al.Therapeutic angiogenesis[J].Arch Surg,1993,128(4):423-429.

[24]KRUPINSKI J,KALUZA J,KUMAR P,et al.Role of angiogenesis in patients with cerebral ischemic stroke[J].Stroke,1994,25(9):1794-1798.

[25]ZAN L,WU H,JIANG J,et al.Temporal profile of Src, Src kinase, and angiogenic factors after focal cerebral ischemia: correlations with angiogenesis and cerebral edema[J].Neurochem Int,2011,58(8):872-879.

[26]TENG H,ZHANG ZG,WANG L,et al.Coupling of angiogenesis and neurogenesis in cultured endothelial cells and neural progenitor cells after stroke[J].J Cereb Blood Flow Metab,2008,28(4):764-771.

[27]XIE YC,LI CY,LI T,et al.Effect of mild hypothermia on angiogenesis in rats with focal cerebral

ischemia[J].Neurosci Lett,2007,422(2):87-90.

[28]LIAO SJ,LIN JW,PEI Z,et al.Enhanced angiogenesis with dl-3n-butylphthalide treatment after focal cerebral ischemia inRHRSP[J].Brain Res,2009,1289:69-78.

[29]LUO C,YI B,FAN W,et al.Enhanced angiogenesis andastrocyte activation by ecdysterone treatment in a focal cerebralischemia rat model[J].Acta Neurochir Suppl,2011,110(pt1):151-155.

[30]ZHANG R,WANG L,ZHANG L,et al.Nitric oxide enhancesangiogenesis via the synthesis of vascular endothelial growthfactor and cgmp after stroke in the rat[J].Circ Res,2003,92(3):308-313.

[31]ZHANG ZG,ZHANG L,JIANG Q,et al.VEGF enhancesangiogenesis and promotes blood-brain barrierleakage in theischemic brain[J].J Clin Invest,2000,106(7):829-838.

[32]LU RY,LUO DF,XIAO SH,et al.Kallikrein gene transferinduces angiogenesis and further improves regional cerebralblood flow in the early period after cerebral ischemia/reperfusionin rats[J].CNS Neurosci Ther,2012,18(5):395-399.

[33]DING G,JIANG Q,LI L,et al.Magnetic resonance imaginginvestigation of axonal remodeling and angiogenesis after embolicstroke in sildenafil-treated rats[J].J Cereb Blood Flow Metab,2008,28(8):1440-1448.

[34]LI L,JIANG Q,ZHANG L,et al.Angiogenesis and improvedcerebral blood flow in the ischemic boundary area detected byMRI after administration of sildenafil to rats with embolic stroke[J].Brain Res,2007,1132(1):185-192.

[35]WEI L,FRASER JL,LU ZY,et al.Transplantation of hypoxiapreconditioned bone marrow mesenchymal stem cells enhancesangiogenesis and neurogenesis after cerebral ischemia in rats [J].Neurobiol Dis,2012,46(3):635-645.

[36]王杰华,刘楠,杜厚伟,等.脂肪来源的干细胞移植对大鼠脑缺血后微血管生成及bFGF和VEGF表达的影响[J].细胞与分子免疫学杂志,2008,24(10):958-961.

[37]CHEN J,CUI X,ZACHAREK A,et al.eNOS mediatesTO90317 treatment-induced angiogenesis and functional outcomeafter stroke in mice[J].Stroke,2009,40(7):2532-2538.

[38]ZHANG L,ZHANG RL,WANG Y,et al.Functional recoveryin aged and young rats after embolic stroke treatment with aphosphodiesterase type 5 inhibitor[J].Stroke,2005,36(14):847-852.

[39]WANG Y,KILIC E,KILIC U,et al.VEGF overexpressioninduces post-ischaemic neuroprotection,but facilitateshaemodynamic steal phenomena[J].Brain,2005,128(Pt1):52-63.

[40]KOJIMA T,HIROTA Y,EMA M,et al.Subventricular zone-derivedneural progenitor cells migrate along a blood vesselscaffold toward the post-stroke striatum[J].Stem Cells,2010,28(3):545-554.

[41]CHEN J,ZHANG ZG,LI Y,et al.Statins induce angiogenesis,neurogenesis,and synaptogenesis after stroke[J].Ann Neurol,2003,53(6):743-751.

[42]YAO RQ,ZHANG L,WANG W,et al.Cornel iridoid glycosidepromotes neurogenesis and angiogenesis and improves neurolo-gical function after focal cerebral ischemia in rats[J].Brain ResBull,2009,79(1):69-76.

[43]PHLMER TD,WILLHOITE AR,GAGE FH,et al.Vascular niche for adult hippocampal neurogenesis [J].J Comp Neurol,2000,425(4):479-494.