



中华内分泌代谢杂志 » 2013, Vol. 29 » Issue (6): 494-497 DOI: 10.3760/cma.j.issn.1000-6699.2013.06.012

临床研究

[最新目录](#) | [下期目录](#) | [过刊浏览](#) | [高级检索](#)

## 低浓度性激素结合球蛋白与代谢综合征的发生相关

缪琳,石娟,洪洁,张翼飞,顾卫琼,李丽娟,王柏凯,王睿,张娟娟,王计秋,李小英,王卫庆,宁光

200025 上海交通大学医学院附属瑞金医院内分泌代谢病科,上海市内分泌代谢病临床医学中心,上海市内分泌代谢病研究所

Low level of serum sex hormone binding globulin is associated with the occurrence of metabolic syndrome.

MAO Lin, SHI Juan, HONG Jie, ZHANG Yi-fei, GU Wei-qiong, LI Li-juan, WANG Bo-kai, WANG Rui, ZHAN Ying, WANG Wei-qing, NI NG Guang. Department of Endocrine and Metabolic Diseases, Shanghai Clinical Center for Endocrine and Metabolic Disease, Shanghai Institute of Endocrinology and Metabolic Diseases, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

Department of Endocrine and Metabolic Diseases, Shanghai Clinical Center for Endocrine and Metabolic Disease, Shanghai Jiaotong University School of Medicine, Shanghai 200025, China

[摘要](#)

[参考文献](#)

[相关文章](#)