

综述

黑色素与常见病原真菌致病性的关系

周真, 杜妍娴, 李希清

中山大学孙逸仙纪念医院皮肤性病科, 广州, 510120

摘要: 黑色素是一类广泛存在于微生物及动植物的生物大分子聚合物,被认为是多数病原真菌的毒力因子,具有抗氧化、吸收紫外线、抵御外界不良环境及宿主免疫等作用。本文就黑色素的生物学特点、作用及其与常见病原真菌致病性的关系等方面作一综述。

关键词: 黑色素 病原真菌 致病性

Abstract:

Keywords:

收稿日期 2011-06-28 修回日期 网络版发布日期

DOI:

基金项目:

NSFC-广东联合基金重点项目(U0932003)

通讯作者: 李希清,E-mail:xiqing.li@gmail.com

作者简介: 周真,女(汉族),硕士研究生在读.E-mail: zdzz19881025@126.com

作者Email: xiqing.li@gmail.com

参考文献:

- [1] Revankar SG,Sutton DA.Melanized fungi in human disease[J]. Clin microbiol Rev,2010,23(4):884-928.
- [2] Dadachova E,Casadevall A.Ionizing radiation:how fungi cope,adapt,and exploit with the help of melanin[J]. Curr Opin Microbiol,2008,11 (6):525-531.
- [3] Ruan L,Yu Z,Fang B.Melanin pigment formation and increased UV resistance in Bacillus thuringiensis following high temperature induction[J]. Syst Appl Microbiol,2004,27 (3):286-289.
- [4] Bell AA,Wheeler MH.Biosynthesis and functions of fungal melanins[J]. Ann Rev Phytopathol,1986,24(1):411-451.
- [5] Howard RJ,Valent B.Breaking and entering:host penetration by the fungal rice blast pathogen Magnaporthe grisea[J]. Annu Rev Microbiol,1996,50(1):491-512.
- [6] Butler MJ,Day AW,Henson JM.Pathogenic properties of fungal melanins[J]. Mycologia,2001,93(1):1-8.
- [7] Garcia-Rivera J,Tucker SC,Feldmesser M.Laccase expression in murine pulmonary Cryptococcus neoformans infection [J]. Infect Immun,2005,73(5):3124-3127.
- [8] Jacobson ES.Pathogenic roles for fungal melanins[J]. Clin Microbiol Rev,2000,13(4):708-817.
- [9] Hamilton A,Gomez B.Melanins in fungal pathogens[J]. J Med Microbiol,2002,51 (3):189-191.
- [10] Langfelder K,Streibel M,Jahn B.Biosynthesis of fungal melanins and their importance for human pathogenic fungi[J]. Fungal Genet Biol,2003,38(2):143-158.
- [11] Lanisnik Rizner T,Wheeler MH.Melanin biosynthesis in the fungus Curvularia lunata (teleomorph: Cochliobolus lunatus) [J]. Can J Microbiol,2003,49(2):110-119.
- [12] 曹志艳,杨胜勇,董金皋.植物病原真菌黑色素与致病性关系的研究进展[J]. 微生物学通报,2006,33(1):154-158.
- [13] Nurudeen TA,Abeam DG.Regulation of melanin production by Cryptococcus neoformans[J]. J Clin Microbiol,1979,10

扩展功能

本文信息

- Supporting info
- PDF(391KB)
- [HTML全文]
- 参考文献[PDF]
- 参考文献

服务与反馈

- 把本文推荐给朋友
- 加入我的书架
- 加入引用管理器
- 引用本文
- Email Alert
- 文章反馈
- 浏览反馈信息

本文关键词相关文章

- 黑色素
- 病原真菌
- 致病性

本文作者相关文章

- 周真
- 杜妍娴
- 李希清

PubMed

- Article by
- Article by
- Article by

(5): 724-729.

- [14] Nosanchuk JD, Cómez BL, Youngchim S. Histoplasma capsulatum synthesizes melanin-like pigments in vitro and during mammalian infection[J]. Infect Immun, 2002, 70(9): 5124-5131.
- [15] Morris-Jones R, Gomez BL, Diez S. Synthesis of melanin pigment by Candida albicans in vitro and during infection[J]. Infect Immun, 2005, 73(9): 6147-6150.
- [16] Alviano DS, Franzen AJ, Travassos LR. Melanin from Fonsecaea pedrosoi induces production of human antifungal antibodies and enhances the antimicrobial efficacy of phagocytes[J]. Infect Immun, 2004, 72(1): 229-237.
- [17] Costa JM, Corbellini VA, Scrofemeker ML. Study of different nitrogen sources on glucose uptake and production of melanin precursors and fungal mass of Fonsecaea pedrosoi cultured in tricyclazole[J]. Process Biochem, 2004, 39(5): 633-636.
- [18] Cunha MM, Franzen AJ, Alviano DS. Inhibition of melanin synthesis pathway by tricyclazole increases susceptibility of Fonsecaea pedrosoi against mouse macrophages[J]. Microsc Res Tech, 2005, 68(6): 377-384.
- [19] Franzen AJ, Cunha MM, Batista EJ. Effects of tricyclazole (5-methyl-1,2,4-triazol 3,4 berzothiazole), a specific DHN-melanin inhibitor, on the morphology of Fonsecaea pedrosoi conidia and sclerotic cells[J]. Microsc Res Tech, 2006, 69(9): 729-737.
- [20] Bocca AL, Brito PP, Figueiredo F. Inhibition of nitric oxide production by macrophages in chromoblastomycosis: a role for Fonsecaea pedrosoi melanin[J]. Mycopathologia, 2006, 161(4): 195-203.
- [21] Cunha MM, Franzen AJ, Seabra SH. Melanin in Fonsecaea pedrosoi: a trap for oxidative radicals[J]. BMC Microbiol, 2010, 10: 80.
- [22] Steenbergen IN, Casadevall A. The origin and maintenance of virulence for the human pathogenic fungus Cryptococcus neoformans[J]. Microbes Infect, 2003, 5(7): 667-675.
- [23] Zhu X, Williamson PR. Role of laccase in the biology and virulence of Cryptococcus neoformans[J]. FEMS yeast Re, 2004, 5(1): 1-10.
- [24] Noverr MC, Williamson PR, Fajardo RS. CNLAC 1 is required for extrapulmonary dissemination of Cryptococcus neoformans but not pulmonary persistence[J]. Infect Immun, 2004, 72(3): 1693-1699.
- [25] Missall TA, Moran JM, Corbett JA. Distinct stress responses of two functional laccases in Cryptococcus neoformans are revealed in the absence of the thiol-specific antioxidant Tsa1[J]. Eukaryot Cell, 2005, 4(1): 202-208.
- [26] Latge JP. Aspergillus fumigatus and aspergillosis[J]. Clin microbiol Rev, 1999, 12(2): 310-350.
- [27] Jahn B, Koch A, Schmidt A. Isolation and characterization of a pigmentless-conidium mutant of Aspergillus fumigatus with altered conidial surface and reduced virulence[J]. Infect Immun, 1997, 65(12): 5110-5117.
- [28] Tsai HF, Chang YC, Washburn RG. The developmentally regulated alb I gene of Aspergillus fumigatus: its role in modulation of conidial morphology and virulence[J]. J Bacteriol, 1998, 180(12): 3031-3038.
- [29] Pihet M, Vandeputte P, Tronchin G. Melanin is an essential component for the integrity of the cell wall of Aspergillus fumigatus conidia[J]. BMC Microbiol, 2009, 9: 177.
- [30] Sugareva V, Hartl A, Brock M. Characterisation of the laccase-encoding gene abr2 of the dihydroxynaphthalene-like melanin gene cluster of Aspergillus fumigatus[J]. Arch Microbiol, 2006, 186(5): 345-355.
- [31] Chai LY, Netea MG, Sugui J. Aspergillus fumigatus conidial melanin modulates host cytokine response[J]. Immunobiology, 2010, 215(11): 915-920.
- [32] Butler M, Day A. Fungalmelanins: a review[J]. Can J Microbiol, 1998, 44(12): 1115-1136.
- [33] Romero-Martinez R, Wheeler M, Guerrero-Plata A. Biosynthesis and functions of melanin in Sporothrix schenckii[J]. Infect Immun, 2000, 68(6): 3696-3703.
- [34] Morris-Jones R, Youngchim S, Gomez BL. Synthesis of melanin-like pigments by Sporothrix schenckii in vitro and mammalian infection[J]. Infect Immun, 2003, 71(7): 4026-4033.
- [35] Almeida-Paes R, Frases S, Fialho Monteiro PCF. Growth conditions influence melanization of Brazilian clinical Sporothrix schenckii isolates[J]. Microbes Infect, 2009, 11(5): 554-562.
- [36] Teixeira PA, De Castro RA, Ferreira FR. L-DOPA accessibility in culture medium increases melanin expression and virulence of Sporothrix schenckii yeast cells[J]. Med Mycol, 2010, 48(5): 687-695.
- [37] Montejo M. Epidemiology of invasive fungal infection in solid organ transplant[J]. Rev Iberoamericana Micol, 2011, 28(3): 120-123.
- [38] Walker CA, Gomez BL, Mora-Montes HM. Melanin externalization in Candida albicans depends on cell wall chitin structures[J]. Eukaryot Cell, 2010, 9(9): 1329-1342.
- [39] Youngchim S, Hay RJ, Hamilton AJ. Melanization of Penicillium marneffeii in vitro and in vivo[J]. Microbiology, 2005, 151(Pt1): 291-299.
- [40] Woo PC, Tam EW, Chong KT. High diversity of polyketide synthase genes and the melanin biosynthesis gene cluster in Penicillium marneffeii[J]. FEBS J, 2010, 277(18): 3750-3758.
- [41] Hort W, Mayser P. Malassezia virulence determinants[J]. Curr Opin Infect Dis, 2011, 24(2): 100-105.
- [42] Gaitanis G, Chnsapi V, Velegri A. Novel application of the masson-fontana stain for demonstrating Malassezia species melanin-like pigment production in vitro and in vitro and in clinical specimens[J]. J Clin Microbiol, 2005, 43(8): 4147-4151.

本刊中的类似文章

1. 张金芳, 万力, 林元珠. 普通裂褶菌真菌学及实验室研究进展[J]. 中国真菌学杂志, 2011, 6(4): 252-256