



		Sign in
The Bulletin of TOKYO DENTAL COLLEGE	Published by Tokyo	o Dental College, Japan
Available Issues Japanese		>> <u>Publisher Site</u>
	ANCED Volume Page arch	Go
Add to Favorite/Citation Articles Alerts	Add to Favorite Publications	Register
TOP > Available Issues > Table of Cont	nts > Abstract	
		PRINT ISSN : 0040-8891
The Bulletin of Tokyo Dental College		
Vol. 48 (2007), No. 2:47-56		

Effect of Cevimeline on Radiation-Induced Salivary Gland Dysfunction and AQP5 in Submandibular Gland in Mice

<u>Katsuhiro Takakura</u>¹⁾, <u>Sachiko Takaki</u>¹⁾, <u>Ienaka Takeda</u>²⁾, <u>Nobuyuki Hanaue</u>¹⁾, <u>Yasuhiro Kizu</u>¹⁾, <u>Morio Tonogi</u>¹⁾ and <u>Gen-yuki Yamane</u>¹⁾

- 1) Department of Oral Medicine, Oral and Maxillofacial Surgery, Tokyo Dental Collage
- 2) Department of Radiation Oncology, Cross Cancer Institute, University of Alberta

(Received January 17, 2007) (Accepted May 18, 2007)

[PDF (266K)] [References]

Abstract: The aim of this study was to clarify the effects of the muscarinic receptor agonist, cevimeline, on saliva flow and expression of aquaporin5 (AQP5) in submandibular gland after X-ray irradiation. Using a previously established radiation-induced xerostomia model mouse, saliva flow from at 7 days before irradiation to at 28 days after irradiation was investigated in mice that were treated with cevimeline before or after irradiation. Radiation caused a significant decrease in saliva flow compared with nonirradiated salivary glands. Cevimeline post-treatment also caused a significant decrease in saliva flow. In contrast, cevimeline pre-treatment did not significantly decrease saliva flow. Expression of AQP5 fluorescent intensity and mRNA were also analyzed. Irradiation significantly decreased expression of AQP5 in submandibular gland. However, pre-treatment with cevimeline prevented this decrease in AQP5 expression. These data suggest that pretreatment with cevimeline prevents radiation-induced xerostomia and radiation-induced decrease in expression of AQP5 in submandibular gland.

Key words: Cevimeline, Aquaporin5, Radiation, Xerostomia, Salivary gland

[PDF (266K)] [References]

To cite this article:

Katsuhiro Takakura, Sachiko Takaki, Ienaka Takeda, Nobuyuki Hanaue, Yasuhiro Kizu, Morio Tonogi and Gen-yuki Yamane: "Effect of Cevimeline on Radiation-Induced Salivary Gland Dysfunction and AQP5 in Submandibular Gland in Mice". The Bulletin of Tokyo Dental College, Vol. **48**: 47-56 (2007) .

doi:10.2209/tdcpublication.48.47

JOI JST.JSTAGE/tdcpublication/48.47

Copyright (c) 2007 by Tokyo Dental College, Japan











Japan Science and Technology Information Aggregator, Electronic

