

中药复方抗癌汤联合化疗治疗晚期口腔癌疗效观察

宋琰华; 宋树田; 薛乐勋;

河南省直属机关第一医院; 郑州市二七人民医院; 郑州大学医学院医学实验中心 450003 郑州;

Observe the effects of Chinese herb complex decoction on tumour combination chemotherapy on advanced mouth squamous cell carcinoma

SONG Yan-hua; SONG Shu-tian; XUE Le-xun;

- 摘要
- 参考文献
- 相关文章

全文: PDF (79 KB) HTML (0 KB) 输出: BibTeX | EndNote (RIS) 背景资料

摘要 目的 观察中药复方抗癌汤联合化疗晚期口腔鳞状细胞癌的疗效。方法 将 12例晚期口腔癌患者随机分为治疗组 7例,对照组 5例,治疗 2个月,观察病灶大小、生存率及毒副反应,并观察治疗后食欲、食量及体重的变化。结果 治疗组和对照组总有效(CR +PR)率分别为 85.71%及 60.0% (P <0.05),12个月生存率分别为 100%及 40%,18个月生存率分别为 57.14%及 0% (P <0.01)。治疗组患者毒副反应轻微、食欲、食量及体重均有不同程度增加,而对照组变化不明显。结论 中药复方抗癌汤配合化疗能明显提高晚期口腔鳞状细胞癌的疗效,并有减轻化疗毒性的作用

关键词: 中药复方抗癌汤 口腔癌 化学疗法

Abstract: Objective To observe the effects of Chinese herb complex decoction on tumour combination chemotherapy on advanced mouth squamous cell carcinoma. Methods 12 patients with advanced mouth squamous cell carcinoma were randomly divided into two groups, 7 patients serves as treated group and 5 patients as controls. After 2 months, observe the size of carcinoma, survival rate, adverse effects and the appetite, quantities of eating, body weight of all patients. Results The total effective rate has statistical difference between the two groups. The total effective rate of the treatment group and control group were 85.71% and 60.0% (P < 0.05), the survival rate of 12 months were 100% and 40%, the survival rate of 18 months were 57.14% and 0% (P < 0.01). The adverse effects of the treatment group were mild, the appetite, quantities of eating and body weight of the treatment group had different degrees of increase, while the control group had no obvious change. Conclusion Chinese herb complex decoction combined with chemotherapy can significantly improve the efficacy of advanced mouth squamous cell carcinoma, and has the effect of reducing the toxicity of chemotherapy.

Key words: Chinese herb complex decoction on tumour Advanced mouth squamous cell carcinoma Chemotherapy

收稿日期: 2002-02-19;

通讯作者: 宋琰华

引用本文:

宋琰华, 宋树田, 薛乐勋. 中药复方抗癌汤联合化疗治疗晚期口腔癌疗效观察[J]. 肿瘤防治研究, 2002, 29(4): 337-338.

SONG Yan-hua, SONG Shu-tian, XUE Le-xun. Observe the effects of Chinese herb complex decoction on tumour combination chemotherapy on advanced mouth squamous cell carcinoma[J]. CHINA RESEARCH ON PREVENTION AND TREATMENT, 2002, 29(4): 337-338.

没有本文参考文献

[1] 张振华; 吴敬波. 脂质体阿霉素热化疗对食管癌细胞的毒性实验研究[J]. 肿瘤防治研究, 2011, 38(7): 736-739.

[2] 许涛; 景红霞; 曹凤军; 李林均; 邓守恒; 雷金华; 俞远东; 陈萍. 局部晚期非小细胞肺癌三维适形放疗加培美曲塞卡铂同期化疗的疗效分析 [J]. 肿瘤防治研究, 2011, 38(7): 848-849.

[3] 李刚; 谭晓虹. RNA干扰survivin对口腔表皮样癌细胞株 KB生长的抑制作用[J]. 肿瘤防治研究, 2011, 38(3): 257-260.

[4] 朱海生; 杨云利; 赵丹; 刘颖新; 朱小东. 171例老年鼻咽癌临床及预后因素分析[J]. 肿瘤防治研究, 2011, 38(3): 281-285.

[5] 李嵘; 刘秀峰; 何泽明; 华海清; 王琳; 陈映霞. 伊立替康联合替加氟持续静脉滴注治疗晚期结肠癌的临床观察[J]. 肿瘤防治研究, 2011, 38(12): 1419-1422.

[6] 张彬; 乔田奎. 肺癌同期放化疗中放射性肺损伤的相关因素分析[J]. 肿瘤防治研究, 2010, 37(5): 578-581.

服务

- 把本文推荐给朋友
- 加入我的书架
- 加入引用管理器
- E-mail Alert
- RSS

作者相关文章

- 宋琰华
- 宋树田
- 薛乐勋

- [7] 陈坤; 蒯强; 赵彦南; 毕艳华; 王娜; 刘月娥. 吉西他滨与紫杉醇分别联合顺铂治疗非小细胞肺癌的对比研究[J]. 肿瘤防治研究, 2010, 37(06): 709-711.
- [8] 吴琼雅; 周道安; 李爱武; 徐瑛; 吴红宇; 蔡勇; 刘宇. 老年局部晚期非小细胞肺癌放化疗综合治疗的疗效分析 [J]. 肿瘤防治研究, 2008, 35(05): 352-354.
- [9] 薛兴欢; 张淑群; 姜建涛; 王西京; 薛锋杰; 刘晓旭. 靶向survivin siRNA与5-Fu协同抑制MCF-7细胞增殖[J]. 肿瘤防治研究, 2008, 35(04): 255-257.
- [10] 宋德刚; 刘杰; 王哲海; 宋宝; 李长征. XRCC1单核苷酸多态性预测晚期NSCLC铂类药物化疗的敏感性[J]. 肿瘤防治研究, 2007, 34(11): 845-847.
- [11] 曹立明; 胡成平; 杨红忠; 苏晓莉; 杨华平. 98例小细胞肺癌化疗疗效分析[J]. 肿瘤防治研究, 2007, 34(10): 785-787.
- [12] 张素欣; 王士杰; 段玉芹; 许彦枝. 口腔癌患者外周血树突状细胞的体外诱导培养及鉴定[J]. 肿瘤防治研究, 2007, 34(10): 811-812.
- [13] 高建飞; 朱宇泽; 饶智国; 章必成; 欧武陵; 杜光祖. 草酸铂联合异长春花碱治疗晚期NSCLC [J]. 肿瘤防治研究, 2006, 33(8): 593-595.
- [14] 陈志明; 丘希辉; 杨钰贤; 庄晓文; 江艺; 张盛奇; 林丹霞. 地榆升白片预防NP方案化疗患者白细胞减少症 [J]. 肿瘤防治研究, 2006, 33(7): 527-528.
- [15] 唐求; 杨坤禹; 王鹏; 胡绍; 胡从良. 大分割三维适形放疗联合低剂量密集化疗治疗III期非小细胞肺癌72例临床分析 [J]. 肿瘤防治研究, 2006, 33(6): 453-454.