

膀胱不同充盈状态下RapidArc与IMRT对早期宫颈癌靶区剂量的影响

王 瑜^①, 曲雅勤^①, 贾晓晶^②, 夏文明^①, 王立波^①, 蒋鑫萍^①, 郝 毅^①

作者单位: ①吉林大学白求恩第一医院放疗科(长春市130021); ②吉林大学白求恩第二医院放疗科

Effect of RapidArc and IMRT target doses at various bladder filling status on early cervical cancer

Yu WANG¹, Yaqin QU¹, Xiaojing JIA², Wenming XIA¹, Libo WANG¹, Xinping JIANG¹, Yi HAO¹

1Radiotherapy Department, Cancer Center, The First Norman Bethune Hospital of Jilin University and

2Radiotherapy Department, The Second Norman Bethune Hospital of Jilin University, Changchun130021, China.

[摘要](#)[图/表](#)[参考文献\(0\)](#)[相关文章\(15\)](#)

友情链接



中国科学技术协会



中国抗癌协会



天津市肿瘤医院

cnki 中国知网
www.cnki.net维普网
仓储式在线出版平台万方数据
WANFANG DATA

版权所有 ©2013 《中国肿瘤临床》编辑部

地址: 天津市河西区体院北环湖西路肿瘤医院内 300060

电话/传真: (022)23527053 E-mail: cjco@cjco.cn cjcotj@sina.com 津ICP备1200315号